

09/668,021.

PATENT

I hereby certify that on the date specified below, this correspondence is being deposited with the United States Postal Service as first-class mail in an envelope addressed to: Certificate of Correction Branch, Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.

March 8, 2005
Date


Jessica Gaunt

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Inventors : Mary E. Brunkow et al.
Patent No. : 6,803,453 B1
Issue Date : October 12, 2004
Title : ANTIBODIES ASSOCIATED WITH ALTERATIONS IN BONE DENSITY

Docket No. : 60117-32
Date : March 8, 2005

Certificate of Correction Branch
Commissioner for Patents
P.O. Box 1450
Alexandria, VA 22313-1450

Certificate
MAR 15 2005

of Correction

REQUEST FOR REPRINTING OF PATENT

Sir:

A detailed review of U.S. Patent No. 6,803,453, which issued on October 12, 2004, has shown that it contains more than 270 individual errors and entire word substitutions or deletions. Only nine of the errors in the printed version of the patent correspond to minor typographical errors in the application as filed. As a result, at least 260 changes are the responsibility of the Patent Office. Many of the errors and changes significantly alter the meaning of terms in the patent, and would lead readers to miss important content. For example, in several cases, the word "epitope" was replaced with "epitome." "Epitope" is an important term in this patent because it defines an immunological construct. In contrast, the word "epitome" is meaningless in that context as that word means "a brief summary" or "an ideal example." The name of the disease

MAR 15 2005

to which the invention relates, sclerosteosis, is spelled incorrectly on multiple occasions. This would preclude a word search of the text by someone using the Patent Office's own database to find relevant patents.

The inaccurate text in Examples 2-9 at columns 41 through 49 is of particular concern as the Examples now contain an inordinate number of technical errors, including errors in the oligonucleotide sequences. These are unacceptable, as correct oligonucleotide sequences are required for a person of ordinary skill in the art to practice the invention. Unlike some word substitutions, errors in DNA sequences would not be obvious from a reading of the text. Many errors are also found in the technical terms, names of compounds used to practice the experiments, and authors of cited references. Unlike simple typographical errors, such errors clearly are not self-evident from a reading of the text.


A few examples are as follows. In one paragraph (column 35, lines 53-67) the word "transgenic" appears three times, but on each appearance it is incorrectly, and inconsistently, spelled: at line 54, *transcience*; at line 58, *transonic*; and at line 60, *transgress*. If one were searching the Patent Office records using the key word "transgenic" its accurate contextual use in this patent would not be found. Other examples in which meaningless words are substituted include "its" for patients (col. 2, line 30); "smudgy" for strategy (col. 2, line 49); "polygonal" for polyclonal (col. 28, line 33); "sexes" for secretes (col. 29, line 11); "misinformation" for misincorporation (col. 32, line 14); "son" for expression (col. 39, line 3); and "purificanon" for purification (col. 42, line 28). This is just a very small but representative sample of the numerous changes in terms and words. In many cases words are omitted altogether.

Applicants respectfully but strongly submit that the excessive number of errors present in this patent cannot effectively be rectified by a Certificate of Correction. Even with a list of corrections at hand, a person reading the patent would have an enormous burden when trying to understand the text, particularly for those paragraphs that have 5, 10, or more separate errors. Given the sheer number of corrections required, and the technical significance of the incorrect words and polynucleotide sequences in the context of understanding the invention, applicants request that the Patent Office re-print the patent rather than issue a Certificate of Correction. Issuance

of the patent in its current condition reflects badly on the high quality of the Patent Office. The invention and the patent in which it is claimed reflect an extensive investment of time by both the applicants and the Examiner. Applicants expect that this patent will be referred to and read often and it is an embarrassment.

Filed herewith is a marked-up copy of the patent, showing the errors, changes and omitted words.

Respectfully submitted,
Mary E. Brunkow et al.
DAVIS WRIGHT TREMAINE LLP

By 
Jane E. R. Potter
Registration No. 33,332

2600 Century Square
1501 Fourth Avenue
Seattle, WA 98101-1688
Phone: (206) 628-7650
Facsimile: (206) 628-7699



US006803453B1

(12) **United States Patent**
Brunkow et al.

(10) **Patent No.:** **US 6,803,453 B1**
(45) **Date of Patent:** **Oct. 12, 2004**

(54) **ANTIBODIES ASSOCIATED WITH
ALTERATIONS IN BONE DENSITY**

(75) **Inventors:** Mary E. Brunkow, Seattle, WA (US);
David J. Galas, Mercer Island, WA
(US); Brian Kovacevich, Renton, WA
(US); John T. Mulligan, Seattle, WA
(US); Bryan W. Paepfer, Seattle, WA
(US); Jeffrey Van Ness, Seattle, WA
(US); David G. Winkler, Seattle, WA
(US)

(73) **Assignee:** Darwin Discovery Ltd., Bothell, WA
(US)

(*) **Notice:** Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 151 days.

(21) **Appl. No.:** 09/668,021

(22) **Filed:** Sep. 21, 2000

Related U.S. Application Data

(62) Division of application No. 09/449,218, filed on Nov. 24,
1999, now Pat. No. 6,395,511.

(60) Provisional application No. 60/110,283, filed on Nov. 27,
1998.

(51) **Int. Cl.⁷** C07K 16/00

(52) **U.S. Cl.** 530/388.23; 435/331; 424/184.1;
530/387.1; 530/388.1; 530/388.2; 530/389.2

(58) **Field of Search** 435/331; 424/184.1;
530/387.1, 388.1, 388.2, 388.23, 389.2

(56) **References Cited**

U.S. PATENT DOCUMENTS

5,453,492 A	9/1995	Bützow et al.	530/413
5,780,263 A	7/1998	Hastings et al.	435/69.1
5,811,238 A	9/1998	Stemmer et al.	435/6
5,830,721 A	11/1998	Stemmer et al.	435/172.1
5,837,458 A	11/1998	Minshull et al.	435/6

FOREIGN PATENT DOCUMENTS

WO	WO 91/13152	9/1991
WO	WO 92/06693	4/1992

OTHER PUBLICATIONS

Colman et al., Research in Immunology, vol. 145 pp. 33-36,
1994.*

Hay et al. (ATCC Cell Lines and Hybridomas, 8th ed. 1994;
pp. 149, 258, and 428).*

Harlow et al. (Antibodies: A Laboratory Manual, Cold
Spring Harbor Laboratory, Cold Spring Harbor, NY, 1988,
pp. 141-157.*

Bendayan (The Journal of Histochemistry and Cytochemis-
try, 1995. 43(9): 881-886).*

Bost et al. (Immunological Investigations, 1988. 17(6&7):
577-586).*

Birren et al., EMBL Sequence Database, Accession No.
AC003098, Nov. 14, 1997.

Bonaldo et al., "Normalization and Subtraction: Two
Approaches to Facilitate Gene Discovery," *Genome Res.*
6(9):791-1996.

Bonaldo et al., EMBL Sequence Database, Accession No.
AI113131, Sep. 4, 1998.

Hillier et al., EMBL Sequence Database, Accession No.
AA393939, May 19, 1997.

Hsu et al., "The Xenopus dorsalizing factor gremlin identi-
fies a novel family of secreted proteins that antagonize BMP
activities," *Molecular Cell* 1:673-683, 1998.

Iemura et al., "Direct binding of follistatin to a complex of
bone-morphogenetic protein and its receptor inhibits ventral
and epidermal cell fates in early Xenopus embryo", *Proc.*
Natl. Acad. Sci. USA 95:9337-9342, 1998.

Khosla and Riggs, "Concise review for primary-care physi-
cians. Treatment options for osteoporosis," *Mayo Clinic Proc*
70:978-982, 1995.

Riggs, "Overview of osteoporosis," *West J. Med.* 154:63-77,
1991.

* cited by examiner

Primary Examiner—Patrick J. Nolan


(74) **Attorney, Agent, or Firm**—Seed Intellectual Property
Law Group PLLC

(57) **ABSTRACT**





Compositions and methods for increasing bone density
using antibodies directed to a novel class or family of TGF- β
binding proteins are provided. The disclosed compositions
are useful, for example, in the diagnosis, prevention and/or
treatment of diseases associated with a loss of bone density,
for example osteoporosis.




12 Claims, 6 Drawing Sheets


Fig. 1


*Fig. 2*

RNA In Situ Hybridization of Mouse Embryo Sections


Fig. 3A*Fig. 3B**Fig. 3C**Fig. 3D*

*Fig. 4A**Fig. 4B**Fig. 4C*

*Fig. 5*

BMP-5/Beer Dissociation Constant Characterization**.75 1.5 7.5 15 30 60 120 nM BMP-5*****Anti-FLAG immunoprecipitation*****Anti-BMP-5 western blot****Ionic Disruption of BMP-5/Beer Binding**

NaCl(mM)	500	150	150	BMP-5
Beer	+	+	-	western
BMP-5	+	+	+	control

***Anti-FLAG immunoprecipitation*****Anti-BMP-5 western blot*****Fig. 6***

Filed

US 6,803,453 B1

1
**ANTIBODIES ASSOCIATED WITH
ALTERATIONS IN BONE DENSITY**

**CROSS-REFERENCE TO RELATED
APPLICATIONS**

This application is a divisional of U.S. patent application Ser. No. 09/449,218 now U.S. Pat. No. 6,395,511 filed Nov. 24, 1999; which application claims priority from U.S. Provisional Patent Application No. 60/110,283, filed Nov. 27, 1998.

TECHNICAL FIELD

The present invention relates generally to pharmaceutical products and methods and, more specifically, to methods and compositions suitable for increasing the mineral content of bone. Such compositions and methods may be utilized to treat a wide variety of conditions, including for example, osteopenia, osteoporosis, and other disorders in which low bone mineral density are a hallmark of the disease.

BACKGROUND OF THE INVENTION

Two or three distinct phases of changes to bone mass occur over the life of an individual (see Riggs, *West J. Med.* 154:63-77, 1991). The first phase occurs in both men and women, and proceeds to attainment of a peak bone mass. This first phase is achieved through linear growth of the endochondral growth plates, and radial growth due to a rate of periosteal apposition. The second phase begins around age 30 for trabecular bone (flat bones such as the vertebrae and pelvis) and about age 40 for cortical bone (e.g., long bones found in the limbs) and continues to old age. This phase is characterized by slow bone loss, and occurs in both men and women. In women, a third phase of bone loss also occurs, most likely due to postmenopausal estrogen deficiencies. During this phase alone, women may lose an additional 10% of bone mass from the cortical bone and 25% from the trabecular compartment (see Riggs, *supra*).

Loss of bone mineral content can be caused by a wide variety of conditions, and may result in significant medical problems. For example, osteoporosis is a debilitating disease in humans characterized by marked decreases in skeletal bone mass and mineral density, structural deterioration of bone including degradation of bone microarchitecture and corresponding increases in bone fragility and susceptibility to fracture in afflicted individuals. Osteoporosis in humans is preceded by clinical osteopenia (bone mineral density that is greater than one standard deviation but less than 2.5 standard deviations below the mean value for young adult bone), a condition found in approximately 25 million people in the United States. Another 7-8 million patients in the United States have been diagnosed with clinical osteoporosis (defined as bone mineral content greater than 2.5 standard deviations below that of mature young adult bone). Osteoporosis is one of the most expensive diseases for the health care system, costing tens of billions of dollars annually in the United States. In addition to health care-related costs, long-term residential care and lost working days add to the financial and social costs of this disease. Worldwide approximately 75 million people are at risk for osteoporosis.

The frequency of osteoporosis in the human population increases with age, and among Caucasians is predominant in women (who comprise 80% of the osteoporosis patient pool in the United States). The increased fragility and susceptibility to fracture of skeletal bone in the aged is aggravated by the greater risk of accidental falls in this population.

2

More than 1.5 million osteoporosis-related bone fractures are reported in the United States each year. Fractured hips, wrists, and vertebrae are among the most common injuries associated with osteoporosis. Hip fractures in particular are extremely uncomfortable and expensive for the patient, and for women correlate with high rates of mortality and morbidity.

Although osteoporosis has been defined as an increase in the risk of fracture due to decreased bone mass, none of the presently available treatments for skeletal disorders can substantially increase the bone density of adults. There is a strong perception among all physicians that drugs are needed which could increase bone density in adults, particularly in the bones of the wrist, spinal column and hip that are at risk in osteopenia and osteoporosis.

Current strategies for the prevention of osteoporosis may offer some benefit to individuals but cannot ensure resolution of the disease. These strategies include moderating physical activity (particularly in weight-bearing activities) with the onset of advanced age, including adequate calcium in the diet, and avoiding consumption of products containing alcohol or tobacco. For patients presenting with clinical osteopenia or osteoporosis, all current therapeutic drugs and strategies are directed to reducing further loss of bone mass by inhibiting the process of bone absorption, a natural component of the bone remodeling process that occurs constitutively.

For example, estrogen is now being prescribed to retard bone loss. There is, however, some controversy over whether there is any long term benefit to it and whether there is any effect at all on patients over 75 years old. Moreover, use of estrogen is believed to increase the risk of breast and endometrial cancer.

High doses of dietary calcium, with or without vitamin D has also been suggested for postmenopausal women. However, high doses of calcium can often have unpleasant gastrointestinal side effects, and serum and urinary calcium levels must be continuously monitored (see Khosla and Riggs, *Mayo Clin. Proc.* 70:978-982, 1995).

Other therapeutics which have been suggested include calcitonin, bisphosphonates, anabolic steroids and sodium fluoride. Such therapeutics however, have undesirable side effects (e.g., calcitonin and steroids may cause nausea and provoke an immune reaction, bisphosphonates and sodium fluoride may inhibit repair of fractures, even though bone density increases modestly) that may prevent their usage (see Khosla and Riggs, *supra*).

No currently practiced therapeutic strategy involves a drug that stimulates or enhances the growth of new bone mass. The present invention provides compositions and methods which can be utilized to increase bone mineralization, and thus may be utilized to treat a wide variety of conditions where it is desired to increase bone mass. Further, the present invention provides other, related advantages.

SUMMARY OF THE INVENTION

As noted above, the present invention provides a novel class or family of TGF-beta binding-proteins, as well as assays for selecting compounds which increase bone mineral content and bone mineral density, compounds which increase bone mineral content and bone mineral density and methods for utilizing such compounds in the treatment or prevention of a wide variety of conditions.

Within one aspect of the present invention, isolated nucleic acid molecules are provided, wherein said nucleic

fractures
three

fragility

patients

Riggs

strategy

acid molecules are selected from the group consisting of: (a) an isolated nucleic acid molecule comprising sequence ID Nos. 1, 5, 7, 9, 11, 13, or 15, or complementary sequence thereof; (b) an isolated nucleic acid molecule that specifically hybridizes to the nucleic acid molecule of (a) under conditions of high stringency; and (c) an isolated nucleic acid that encodes a TGF-beta binding-protein according to (a) or (b). Within related aspects of the present invention, isolated nucleic acid molecules are provided based upon hybridization to only a portion of one of the above-identified sequences (e.g., for (a) hybridization may be to a probe of at least 20, 25, 50, or 100 nucleotides selected from nucleotides 156 to 539 or 555 to 687 of Sequence ID No. 1). As should be readily evident, the necessary stringency to be utilized for hybridization may vary based upon the size of the probe. For example, for a 25-mer probe high stringency conditions could include: 60 mM Tris pH 8.0, 2 mM EDTA, 5xDenhardt's, 6xSSC, 0.1% (w/v) N-laurylsarcosine, 0.5% (w/v) NP-40 (nonidet P-40) overnight at 45 degrees C., followed by two washes with 0.2xSSC/0.1% SDS at 45-50 degrees. For a 100-mer probe under low stringency conditions, suitable conditions might include the following: 5xSSPE, 5xDenhardt's and 0.5% SDS overnight at 42-50 degrees, followed by two washes with 2xSSPE (or 2xSSC)/0.1% SDS at 42-50 degrees.

Within related aspects of the present invention, isolated nucleic acid molecules are provided which have homology to Sequence ID Nos. 1, 5, 7, 9, 11, 13, or 15, at a 50%, 60%, 75%, 80%, 90%, 95%, or 98% level of homology utilizing a Wilbur-Lipman algorithm. Representative examples of such isolated molecules include, for example, nucleic acid molecules which encode a protein comprising Sequence ID NOs. 2, 6, 10, 12, 14, or 16, or have homology to these sequences at a level of 50%, 60%, 75%, 80%, 90%, 95%, or 98% level of homology utilizing a Lipman-Pearson algorithm.

Isolated nucleic acid molecules are typically less than 100 kb in size, and, within certain embodiments, less than 50 kb, 25 kb, 10 kb, or even 5 kb in size. Further, isolated nucleic acid molecules, within other embodiment; do not exist in a "library" of other unrelated nucleic acid molecules (e.g., a subclone BAC such as described in GenBank Accession No. AC003099 and EMB No. AQ171546). However, isolated nucleic acid molecules can be found in libraries of related molecules (e.g., for shuffling, such as is described in U.S. Pat. Nos. 5,837,458; 5,830,721; and 5,811,238). Finally, isolated nucleic acid molecules as described herein do not include nucleic acid molecules which encode Dan, Cerberus, Gremlin, or SCGF (U.S. Pat. Nos. 5,780,263).

Also provided by the present invention are cloning vectors which contain the above-noted nucleic acid molecules, and expression vectors which comprise a promoter (e.g., a regulatory sequence) operably linked to one of the above-noted nucleic acid molecules. Representative examples of suitable promoters include tissue-specific promoters, and viral-based promoters (e.g., CMV-based promoters such as CMV I-E, SV40 early promoter, and MuLV LTR). Expression vectors may also be based upon, or derived from viruses (e.g., a "viral vector"). Representative examples of viral vectors include herpes simplex viral vectors, adenoviral vectors, adenovirus-associated viral vectors and retroviral vectors. Also provided are host cells containing or comprising any of above-noted vectors (including for example, host cells of human, monkey, dog, rat, or mouse origin).

Within other aspects of the present invention, methods of producing TGF-beta binding-proteins are provided, comprising the step of culturing the aforementioned host cell

containing vector under conditions and for a time sufficient to produce the TGF-beta binding protein. Within further embodiments, the protein produced by this method may be further purified (e.g., by column chromatography, affinity purification, and the like). Hence, isolated proteins which are encoded by the above-noted nucleic acid molecules (e.g., Sequence ID NOs. 2, 4, 6, 8, 10, 12, 14, or 16) may be readily produced given the disclosure of the subject application.

It should also be noted that the aforementioned proteins, or fragments thereof, may be produced as fusion proteins. For example, within one aspect fusion proteins are provided comprising a first polypeptide segment comprising a TGF-beta binding-protein encoded by a nucleic acid molecule as described above, or a portion thereof of at least 10, 20, 30, 50, or 100 amino acids in length, and a second polypeptide segment comprising a non-TGF-beta binding-protein. Within certain embodiments, the second polypeptide may be a tag suitable for purification or recognition (e.g., a polypeptide comprising multiple anionic amino acid residues—see U.S. Pat. No. 4,851,341), a marker (e.g., green fluorescent protein, or alkaline phosphatase), or a toxic molecule (e.g., ricin).

Within another aspect of the present invention, antibodies are provided which are capable of specifically binding the above-described class of TGF-beta binding proteins (e.g., human BEER). Within various embodiments, the antibody may be a polyclonal antibody, or a monoclonal antibody (e.g., of human or murine origin). Within further embodiments, the antibody is a fragment of an antibody which retains the binding characteristics of a whole antibody (e.g., an F(ab')₂, F(ab)₂, Fab', Fab, or Fv fragment, or even a CDR). Also provided are hybridomas and other cells which are capable of producing or expressing the aforementioned antibodies.

Within related aspects of the invention, methods are provided detecting a TGF-beta binding protein, comprising the steps of incubating an antibody as described above under conditions and for a time sufficient to permit said antibody to bind to a TGF-beta binding protein, and detecting the binding. Within various embodiments the antibody may be bound to a solid support to facilitate washing or separation, and/or labeled. (e.g., with a marker selected from the group consisting of enzymes, fluorescent proteins, and radioisotopes).

Within other aspects of the present invention, isolated oligonucleotides are provided which hybridize to a nucleic acid molecule according to Sequence ID NOs. 1, 3-5, 7, 9, 11, 13, 15, 17, or 18 or the complement thereto, under conditions of high stringency. Within further embodiments, the oligonucleotide may be found in the sequence which encodes Sequence ID Nos. 2, 4, 6, 8, 10, 12, 14, or 16. Within certain embodiments, the oligonucleotide is at least 15, 20, 30, 50, or 100 nucleotides in length. Within further embodiments, the oligonucleotide is labeled with another molecule (e.g., an enzyme, fluorescent molecule, or radioisotope). Also provided are primers which are capable of specifically amplifying all or a portion of the above-mentioned nucleic acid molecules which encode TGF-beta binding-proteins. As utilized herein, the term, "specifically amplifying" should be understood to refer to primers which amplify the aforementioned TGF-beta binding-proteins, and not other TGF-beta binding proteins such as Dan, Cerberus, Gremlin, or SCGF (U.S. Pat. No. 5,780,263).

Within related aspects of the present invention, methods are provided for detecting a nucleic acid molecule which

THIS

5

encodes a TGF-beta binding protein, comprising the steps of incubating an oligonucleotide as described above under conditions of high stringency, and detecting hybridization of said oligonucleotide. Within certain embodiments, the oligonucleotide may be labeled and/or bound to a solid support.

Within other aspects of the present invention, ribozymes are provided which are capable of cleaving RNA which encodes one of the above-mentioned TGF-beta binding-proteins (e.g., Sequence ID NOs. 2, 6, 8, 10, 12, 14, or 16). Such ribozymes may be composed of DNA, RNA (including 2'-O-methyl ribonucleic acids), nucleic acid analogs (e.g., nucleic acids having phosphorothioate linkages) or mixtures thereof. Also provided are nucleic acid molecules (e.g., DNA or cDNA) which encode these ribozymes, and vectors which are capable of expressing or producing the ribozymes. Representative examples of vectors include plasmids, retrotransposons, cosmids, and viral-based vectors (e.g., viral vectors generated at least in part from a retrovirus, adenovirus, or, adeno-associated virus). Also provided are host cells (e.g., human, dog, rat, or mouse cells) which contain these vectors. In certain embodiments, the host cell may be stably transformed with the vector.

Within further aspects of the invention, methods are provided for producing ribozymes either synthetically, or by in vitro or in vivo transcription. Within further embodiments, the ribozymes so produced may be further purified and/or formulated into pharmaceutical compositions (e.g., the ribozyme or nucleic acid molecule encoding the ribozyme along with a pharmaceutically acceptable carrier or diluent). Similarly, the antisense oligonucleotides and antibodies or other selected molecules described herein may be formulated into pharmaceutical compositions.

Within other aspects of the present invention, antisense oligonucleotides are provided comprising a nucleic acid molecule which hybridizes to a nucleic acid molecule according to Sequence ID NOs 1, 3-5, 7, 9, 11, 13, or 15, or the complement thereto, and wherein said oligonucleotide inhibits the expression of TGF-beta binding protein as described herein (e.g., human BEER). Within various embodiments, the oligonucleotide is 15, 20, 25, 30, 35, 40, or 50 nucleotides in length. Preferably, the oligonucleotide is less than 100, 75, or 60 nucleotides in length. As should be readily evident, the oligonucleotide may be comprised of one or more nucleic acid analogs, ribonucleic acids, or deoxyribonucleic acids. Further, the oligonucleotide may be modified by one or more linkages, including for example, covalent linkage such as a phosphorothioate linkage, a phosphotriester linkage, a methyl phosphonate linkage, a methylene(methylimino) linkage, a morpholino linkage, an amide linkage, a polyamide linkage, a short chain alkyl intersugar linkage, a cycloalkyl intersugar linkage, a short chain heteroatomic intersugar linkage and a heterocyclic intersugar linkage. One representative example of a chimeric oligonucleotide is provided in U.S. Pat. No. 5,989,912.

Within yet another aspect of the present invention, methods are provided for increasing bone mineralization, comprising introducing into a warm-blooded animal an effective amount of the ribozyme as described above. Within related aspects, such methods comprise the step of introducing into a patient an effective amount of the nucleic acid molecule or vector as described herein which is capable of producing the desired ribozyme, under conditions favoring transcription of the nucleic acid molecule to produce the ribozyme.

Within other aspects of the invention transgenic, non-human animals are provided. Within one embodiment a

6

transgenic animal is provided whose germ cells and somatic cells contain a nucleic acid molecule encoding a TGF-beta binding-protein as described above which is operably linked to a promoter effective for the expression of the gene, the gene being introduced into the animal, or an ancestor of the animal, at an embryonic stage, with the proviso that said animal is not a human. Within other embodiments, transgenic knockout animals are provided, comprising an animal whose germ cells and somatic cells comprise a disruption of at least one allele of an endogenous nucleic acid molecule which hybridizes to a nucleic acid molecule which encodes a TGF-binding protein as described herein, wherein the disruption prevents transcription of messenger RNA from said allele as compared to an animal without the disruption, with the proviso that the animal is not a human. Within various embodiments, the disruption is a nucleic acid deletion, substitution, or, insertion. Within other embodiments the transgenic animal is a mouse, rat, sheep, pig, or dog.

Within further aspects of the invention, kits are provided for the detection of TGF-beta binding-protein gene expression, comprising a container that comprises a nucleic acid molecule, wherein the nucleic acid molecule is selected from the group consisting of (a) a nucleic acid molecule comprising the nucleotide sequence of SEQ ID NOs: 1, 3, 5, 7, 9, 11, 13, or 15; (b) a nucleic acid molecule comprising the complement of the nucleotide sequence of (a); (c) a nucleic acid molecule that is a fragment of (a) or (b) of at least 15, 20, 30, 50, 75, or, 100 nucleotides in length. Also provided are kits for the detection of a TGF-beta binding-protein which comprise a container that comprise one of the TGF-beta binding protein antibodies described herein.

For example, within one aspect of the present invention methods are provided for determining whether a selected molecule is capable of increasing bone mineral content, comprising the steps of (a) mixing one or more candidate molecules with TGF-beta-binding-protein encoded by the nucleic acid molecule according to claim 1 and a selected member of the TGF-beta family of proteins (e.g., BMP 5 or 6), (b) determining whether the candidate molecule alters the signaling of the TGF-beta family member, or alters the binding of the TGF-beta binding-protein to the TGF-beta family member. Within certain embodiments, the molecule alters the ability of TGF-beta to function as a positive regulator of mesenchymal cell differentiation. Within this aspect of the present invention, the candidate molecule(s) may alter signaling or binding by, for example, either decreasing (e.g., inhibiting), or increasing (e.g., enhancing) signaling or binding.

Within yet another aspect, methods are provided for determining whether a selected molecule is capable of increasing bone mineral content, comprising the step of determining whether a selected molecule inhibits the binding of TGF-beta binding-protein to bone, or an analogue thereof. Representative examples of bone or analogues thereof include hydroxyapatite and primary human bone samples obtained via biopsy.

Within certain embodiments of the above-recited methods, the selected molecule is contained within a mixture of molecules and the methods may further comprise the step of isolating one or more molecules which are functional within the assay. Within yet other embodiments, TGF-beta family of proteins is bound to a solid support and the binding of TGF-beta binding-protein is measured or TGF-beta binding-protein are bound to a solid support and the binding of TGF-beta proteins are measured.

Utilizing methods such as those described above, a wide variety of molecules may be assayed for their ability to

animals

from

an

aspect

the

provided

increase bone mineral content by inhibiting the binding of the TGF-beta binding-protein to the TGF-beta family of proteins. Representative examples of such molecules include proteins or peptides, organic molecules, and nucleic acid molecules.

Within other related aspects of the invention, methods are provided for increasing bone mineral content in a warm-blooded animal, comprising the step of administering to a warm-blooded animal a therapeutically effective amount of a molecule identified from the assays recited herein. Within another aspect, methods are provided for increasing bone mineral content in a warm-blooded animal, comprising the step of administering to a warm-blooded animal a therapeutically effective amount of a molecule which inhibits the binding of the TGF-beta binding-protein to the TGF-beta super-family of proteins, including bone morphogenic proteins (BMs). Representative examples of suitable molecules include antisense molecules, ribozymes, ribozyme genes, and antibodies (e.g., a humanized antibody) which specifically recognize and alter the activity of the TGF-beta binding-protein.

Within another aspect of the present invention, methods are provided for increasing bone mineral content in a warm-blooded animal, comprising the steps of (a) introducing into cells which home to the bone a vector which directs the expression of a molecule which inhibits the binding of the TGF-beta binding in to the TGF-beta family of proteins and bone morphogenic proteins (BMPs), and (b) administering the vector-containing cells to a warm-blooded animal. As utilized herein, it should be understood that cells "home to bone" if they localize within the bone matrix after peripheral administration. Within one embodiment, such methods further comprise, prior to the step of introducing, isolating cells from the marrow of bone which home to the bone. Within a further embodiment, the cells which home to bone are selected from the group consisting of CD34+ cells and osteoblasts.

Within other aspects of the present invention, molecules are provided (preferably isolated) which inhibit the binding of the TGF-beta binding-protein to the TGF-beta super-family of proteins.

Within further embodiments, the molecules may be provided as a composition, and can further comprise an inhibitor of bone resorption. Representative examples of such inhibitors include calcitonin, estrogen, bisphosphonates, a growth factor having anti-resorption activity and tamoxifen.

Representative examples of molecules which may be utilized in the afore-mentioned therapeutic contexts include, e.g., ribozymes, ribozyme genes, antisense molecules, and/or antibodies (e.g., humanized antibodies). Such molecules may depending upon their selection, used to alter, antagonize, or agonize the signalling or binding of a TGF-beta binding-protein family member as described herein.

Within various embodiments of the invention, the above-described molecules and methods of treatment or prevention may be utilized on conditions such as osteoporosis, osteomalasia, periodontal disease, scurvy, Cushing's Disease, bone fracture and conditions due to limb immobilization and steroid usage.

These and other aspects of the present invention will become evident upon reference to the following detailed description and attached drawings. In addition, various references are set forth herein which describe in more detail certain procedures or compositions (e.g., plasmids, etc.), and are therefore incorporated by reference in their entirety.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 is a schematic illustration comparing the amino acid sequence of Human Dan; Human Gremlin; Human Cerberus and Human Beer. Arrows indicate the Cysteine backbone.

FIG. 2 summarizes the results obtained from surveying a variety of human tissues for the expression of a TGF-beta binding-protein gene, specifically, the Human Beer gene. A semi-quantitative Reverse Transcription-Polymerase Chain Reaction (RT-PCR) procedure was used to amplify a portion of the gene from first-strand cDNA synthesized from total RNA (described in more detail in EXAMPLE 2A).

FIG. 3 summarizes the results obtained from RNA in situ hybridization of mouse embryo sections, using a cRNA probe that is complementary to the mouse Beer transcript (described in more detail in EXAMPLE 2B). Panel A is a transverse section of 10.5 dpc embryo. Panel B is a sagittal section of 12.5 dpc embryo and panels C and D are sagittal sections of 15.5 dpc embryos.

FIG. 4 illustrates, by western blot analysis, the specificity of three different polyclonal antibodies for their respective antigens (described in more detail in EXAMPLE 4). FIG. 4A shows specific reactivity of an anti-H. Beer antibody for H. Beer antigen, but not H. Dan or H. Gremlin. FIG. 4B shows reactivity of an anti-H. Gremlin antibody for H. Gremlin antigen, but not H. Beer or H. Dan. FIG. 4C shows reactivity of an anti-H. Dan antibody for H. Dan, but not H. Beer or H. Gremlin.

FIG. 5 illustrates, by western blot analysis, the selectivity of the TGF-beta binding-protein, Beer, for BMP-5 and BMP-6, but not BMP-4 (described in more detail in EXAMPLE 5).

FIG. 6 demonstrates that the ionic interaction between the TGF-beta binding-protein, Beer, and BMP-5 has a dissociation constant in the 15-30 nM range.

DETAILED DESCRIPTION OF THE INVENTION

Definition

Prior to setting forth the invention in detail, it may be helpful to an understanding thereof to set forth definitions of certain terms and to list and to define the abbreviations that will be used hereinafter.

"Molecule" should be understood to include proteins or peptides (e.g., antibodies, recombinant binding partners, peptides with a desired binding affinity), nucleic acids (e.g., DNA, RNA, chimeric nucleic acid molecules, and nucleic acid analogues such as PNA); and organic or inorganic compounds.

"TGF-beta" should be understood to include any known or novel member of the TGF-beta super-family, which also includes bone morphogenic proteins (BMPs).

"TGF-beta receptor" should be understood to refer to the receptor specific for a particular member of the TGF-beta super-family (including bone morphogenic proteins (BMPs)).

"TGF-beta binding-protein" should be understood to refer to a protein with specific binding affinity for a particular member or subset of members of the TGF-beta super-family (including bone morphogenic proteins (BMPs)). Specific examples of TGF-beta binding-proteins include proteins encoded by Sequence ID Nos. 1, 5, 7, 9, 11, 13) and 15.

Inhibiting the "binding of the TGF-beta binding-protein to the TGF-beta family of proteins and bone morphogenic proteins (BMPs)" should be understood to refer to molecules which allow the activation of TGF-beta or bone morphogenic proteins (BMPs), or allow the binding of TGF-beta family members including bone morphogenic proteins (BMPs) to their respective receptors, by removing or preventing TGF-beta from binding to TGF-binding-protein.

bisphosphonate
resorptive

Such inhibition may be accomplished, for example, by molecules which inhibit the binding of the TGF beta binding-protein to specific members of the TGF-beta super-family.

"Vector" refers to an assembly which is capable of directing the expression of desired protein. The vector must include transcriptional promoter elements which are operably linked to the gene(s) of interest. The vector may be composed of either deoxyribonucleic acids ("DNA"), ribonucleic acids ("RNA"), or a combination of the two (e.g., a DNA-RNA chimeric). Optionally, the vector may include a polyadenylation sequence, one or more restriction sites, as well as one or more selectable makers such as neomycin phosphotransferase or hygromycin phosphotransferase. Additionally, depending on the host cell chosen and the vector employed, other genetic elements such as an origin of replication, additional nucleic acid restriction sites, enhancers, sequences conferring inducibility of transcription, and selectable markers, may also be incorporated into the vectors described herein.

An "isolated nucleic acid molecule" is a nucleic acid molecule that is not integrated in the genomic DNA of an organism. For example, a DNA molecule that encodes a TGF-binding protein that has been separated from the genomic DNA of a eukaryotic cell is an isolated DNA molecule. Another example of an isolated nucleic acid molecule is a chemically-synthesized nucleic acid molecule that is not integrated in the genome of an organism. The isolated nucleic acid molecule may be genomic DNA, cDNA, RNA, or composed at least in part of nucleic acid analogs.

An "isolated polypeptide" is a polypeptide that is essentially free from contaminating cellular components, such as carbohydrate, lipid, or other proteinaceous impurities associated with the polypeptide in nature. Within certain embodiments, a particular protein preparation contains an isolated polypeptide if it appears nominally as a single band on SDS-PAGE gel with Coomassie Blue staining. "Isolated" when referring to organic molecules means that the compounds are greater than 90 percent pure utilizing methods which are well known in the art (e.g., NMR, melting point).

"Sclerosteosis" Sclerosteosis is a term that was applied by Hansen (1967) (Hansen, H. G., Sklerosteose. In: Opitz, H.; Schmid, F., Handbuch der Kinderheilkunde. Berlin: Springer (pub.) 6 1967. Pp. 351-355) to a disorder similar to van Buchem hyperostosis corticalis generalisata but possibly differing in radiologic appearance of the bone changes and in the presence of asymmetric cutaneous syndactyly of the index and middle fingers in many cases. The jaw has an unusually square appearance in this condition.

"Humanized antibodies" are recombinant proteins in which murine complementary determining regions of monoclonal antibodies have been transferred from heavy and light variable chains of the murine immunoglobulin into a human variable domain.

As used herein, an "antibody fragment" is a portion of an antibody such as F(ab')₂, F(ab)₂, F(ab'), Fab, and the like. Regardless of structure, an antibody fragment binds with the same antigen that is recognized by the intact antibody. For example, an anti-TGF-beta binding-protein monoclonal antibody fragment binds with an epitope of TGF-beta binding-protein.

The term "antibody fragment" also includes any synthetic or genetically engineered protein that acts like an antibody by binding to a specific antigen to form a complex. For example, antibody fragments include isolated fragments

consisting of the light chain variable region, "Fv" fragments consisting of the variable regions of the heavy and light chains, recombinant single chain polypeptide molecules in which light and heavy variable regions are connected by a peptide linker ("sFv proteins"), and minimal recognition units consisting of the amino acid residues that mimic the hypervariable region.

A "detectable label" is a molecule or atom which can be conjugated to an antibody moiety to produce a molecule useful for diagnosis. Examples of detectable labels include chelators, photoactive agents, radioisotopes, fluorescent agents, paramagnetic ions, enzymes, and other marker moieties.

As used herein, an "immunoconjugate" is a molecule comprising an anti-TGF-beta binding-protein antibody, or an antibody fragment, and a detectable label. An immunoconjugate has roughly the same, or only slightly reduced, ability to bind TGF-beta binding-protein after conjugation as before conjugation.

Abbreviations: TGF-beta—"Transforming Growth Factor-beta"; TGF-bBP—"Transforming Growth Factor-beta binding-protein" (one representative TGF-bBP is designated "H. Beer"); BMP—"bone morphogenic protein"; PCR—"polymerase chain reaction"; RT-PCR—PCR process in which RNA is first transcribed into DNA at the first step using reverse transcriptase (RT); cDNA—any DNA made by copying an RNA sequence into DNA form.

As noted above, the present invention provides a novel class of TGF-beta binding-proteins, as well as methods and compositions for increasing bone mineral content in warm-blooded animals. Briefly, the present inventions are based upon the unexpected discovery that a mutation in the gene which encodes a novel member of the TGF-beta binding-protein family results in a rare condition (sclerosteosis) characterized by bone mineral contents which are one- to four-fold higher than in normal individuals. Thus, as discussed in more detail below this discovery has led to the development of assays which may be utilized to select molecules which inhibit the binding of the TGF-beta binding-protein to the TGF-beta family of proteins and bone morphogenic proteins (BMPs), and methods of utilizing such molecules for increasing the bone mineral content of warm-blooded animals (including for example, humans).

Discussion of the Disease Known as Sclerosteosis

Sclerosteosis is a term that was applied by Hansen (1967) (Hansen, H. G., Sklerosteose. In: Opitz, H.; Schmid, F., Handbuch der Kinderheilkunde. Berlin: Springer (pub.) 6 1967. Pp. 351-355) to a disorder similar to van Buchem hyperostosis corticalis generalisata but possibly differing in radiologic appearance of the bone changes and in the presence of asymmetric cutaneous syndactyly of the index and middle fingers in many cases.

Sclerosteosis is now known to be an autosomal semi-dominant disorder which is characterized by widely disseminated sclerotic lesions of the bone in the adult. The condition is progressive. Sclerosteosis also has a developmental aspect which is associated with syndactyly (two or more fingers are fused together). The Sclerosteosis Syndrome is associated with large stature and many affected individuals attain a height of six feet or more. The bone mineral content of homozygotes can be 1 to 6 fold over normal individuals and bone mineral density can be 1 to 4 fold above normal values (e.g., from unaffected siblings).

The Sclerosteosis Syndrome occurs primarily in Afrikaners of Dutch descent in South Africa. Approximately

interest
sequence
inducibility

Sklerosteose

structure
epitope

useful

heterozygotes

1/140 individuals in the Afrikaaner population are carriers of the mutated gene (heterozygotes). The mutation shows 100% penetrance. There are anecdotal reports of increased of bone mineral density in heterozygotes with no associated pathologies (syndactyly or skull overgrowth).

It appears at the present time that there is no abnormality of the pituitary-hypothalamus axis in Sclerosteosis. In particular, there appears to be no over-production of growth hormone and cortisone. In addition, sex hormone levels are normal in affected individuals. However, bone turnover markers (osteoblast specific alkaline phosphatase, osteocalcin, type 1 procollagen C' propeptide (PICP), and total alkaline phosphatase; (see Cornier, C., *Curr. Opin. in Rheu.* 7:243, 1995) indicate that there is hyperosteoblastic activity associated with the disease but that there is normal to slightly decreased osteoclast activity as measured by markers of bone resorption (pyridinoline, deoxypyridinoline, N-telopeptide, urinary hydroxyproline, plasma tartrate-resistant acid phosphatases and galactosyl hydroxyllysine (see Cornier, supra)).

Sclerosteosis is characterized by the continual deposition of bone throughout the skeleton during the lifetime of the affected individuals. In homozygotes the continual deposition of bone mineral leads to an overgrowth of bone in areas of the skeleton where there is an absence of mechanoreceptors (skull, jaw, cranium). In homozygotes with Sclerosteosis, the overgrowth of the bones of the skull leads to cranial compression and eventually to death due to excessive hydrostatic pressure on the brain stem. In all other parts of the skeleton there is a generalized and diffuse sclerosis. Cortical areas of the long bones are greatly thickened resulting in a substantial increase in bone strength. Trabecular connections are increased in thickness which in turn increases the strength of the trabecular bone. Sclerotic bones appear unusually opaque to x-rays.

As described in more detail in Example 1, the rare genetic mutation that is responsible for the Sclerosteosis syndrome has been localized to the region of human chromosome 17 that encodes a novel member of the TGF-beta binding-protein family (one representative example of which is designated "H. Beer"). As described in more detail below, based upon this discovery, the mechanism of bone mineralization is more fully understood, allowing the development of assays for molecules which increase bone mineralization, and use of such molecules to increase bone mineral content, and in the treatment or prevention of a wide number of diseases.

TGF-beta Super-Family

The Transforming Growth Factor-beta (TGF-beta) super-family contains a variety of growth factors that share common sequence elements and structural motifs (at both the secondary and tertiary levels). This protein family is known to exert a wide spectrum of biological responses on a large variety of cell types. Many of them have important functions during the embryonal development in pattern formation and tissue specification; in adults they are involved, e.g., in wound healing and bone repair and bone remodeling, and in the modulation of the immune system. In addition to the three TGF-beta's, the super-family includes the Bone Morphogenic Proteins (BMPs), Activins, Inhibins, Growth and Differentiation Factors (GDFs), and Glial-Derived Neurotrophic Factors (GDNFs). Primary classification is established through general sequence features that bin a specific protein into a general sub-family. Additional stratification within the sub-family is possible due to stricter sequence

conservation between members of the smaller group. In certain instances, such as with BMP-5, BMP-6 and BMP-7, this can be as high as 75 percent amino acid homology between members of the smaller group. This level of identity enables a single representative sequence to illustrate the key biochemical elements of the sub-group that separates it from other members of the larger family.

TGF-beta signals by inducing the formation of hetero-oligomeric complexes of type I and type II receptors. The crystal structure of TGF-beta2 has been determined. The general fold of the TGF-beta2 monomer contains a stable, compact, cysteine knotlike structure formed by three disulfide bridges. Dimerization, stabilized by one disulfide bridge, is antiparallel.

TGF-beta family members initiate their cellular action by binding to receptors with intrinsic serine/threonine kinase activity. This receptor family consists of two subfamilies, denoted type I and type II receptors. Each member of the TGF-beta family binds to a characteristic combination of type I and type II receptors, both of which are needed for signaling. In the current model for TGF-beta receptor activation, TGF-beta first binds to the type II receptor (TbR-II), which occurs in the cell membrane in an oligomeric form with activated kinase. Thereafter, the type I receptor (TbR-I), which can not bind ligand in the absence of TbR-II, is recruited into the complex. TbR-II then phosphorylates TbR-I predominantly in a domain rich in glycine and serine residues (GS domain) in the juxtamembrane region, and thereby activates TbR-I.

Thus far seven type I receptors and five type II receptors have been identified.

Bone Morphogenic Proteins (BMPs) are Key Regulatory Proteins in Determining Bone Mineral Density in Humans

A major advance in the understanding of bone formation was the identification of the bone morphogenic proteins (BMPs), also known as osteogenic proteins (OPs), which regulate cartilage and bone differentiation *in vivo*. BMPs/OPs induce endochondral bone differentiation through a cascade of events which include formation of cartilage, hypertrophy and calcification of the cartilage, vascular invasion, differentiation of osteoblasts, and formation of bone. As described above, the BMPs/OPs (BMP 2-14, and osteogenic protein 1 and -2, OP-1 and OP-2) are members of the TGF-beta super-family. The striking evolutionary conservation between members the BMP/OP sub-family suggests that they are critical in the normal development and function of animals. Moreover, the presence of multiple forms of BMPs/OPs raises an important question about the biological relevance of this apparent redundancy. In addition to postfetal chondrogenesis and osteogenesis, the BMPs/OPs play multiple roles in skeletogenesis (including the development of craniofacial and dental tissues) and in embryonic development and organogenesis of parenchymatous organs, including the kidney. It is now understood that nature relies on common (and few) molecular mechanisms tailored to provide the emergence of specialized tissues and organs. The BMP/OP super-family is an elegant example of nature parsimony in programming multiple specialized functions deploying molecular isoforms with minor variation in amino acid motifs within highly conserved carboxy-terminal regions.

BMP Antagonism

The BMP and Activin sub-families are subject to significant post-translational regulation. An intricate extracellular

cellkline

control system exists, whereby a high affinity antagonist is synthesized and exported, and subsequently complexes selectively with BMPs or activins to disrupt their biological activity (W. C. Smith (1999) TIG 15(1) 3-6). A number of these natural antagonists have been identified, and based on sequence divergence to have evolved independently due to the lack of primary sequence conservation. There has been no structural work to date on this class of proteins. Studies of these antagonists has highlighted a distinct preference for interacting and neutralizing BMP-2 and BMP-4. Furthermore, the mechanism of inhibition seems to differ for the different antagonists (S. Iemura et al. (1998) *Proc Natl Acad Sci USA* 95 9337-9342).

Novel TGF-beta Binding-Proteins

1. Background re: TGF-beta Binding-proteins

As noted above, the present invention provides a novel class of TGF-beta binding-proteins that possess a nearly identical cysteine (disulfide) scaffold when compared to Human DAN, Human Gremlin, and Human Cerberus, and SCGF (U.S. Pat. No. 5,780,263) but almost no homology at the nucleotide level (for background information, see generally Hsu, D. R., Economides, A. N., Wang, X., Eimon, P. M., Harland, R. M., "The Xenopus Dorsalizing Factor Gremlin Identifies a Novel Family of Secreted Proteins that Antagonize BMP Activities," *Molecular Cell* 1:673-683, 1998).

One representative example of the novel class of TGF-beta binding-proteins is disclosed in Sequence ID Nos. 1, 5, 9, 11, 13, and 15. Representative members of this class of binding proteins should also be understood to include variants of the TGF-beta binding-protein (e.g., Sequence ID Nos. 5 and 7). As utilized herein, a "TGF-beta binding-protein variant gene" refers to nucleic acid molecules that encode a polypeptide having an amino acid sequence that is a modification of SEQ ID Nos: 2, 10, 12, 14 or 16. Such variants include naturally-occurring polymorphisms or allelic variants of TGF-beta binding-protein genes, as well as synthetic genes that contain conservative amino acid substitutions of these amino acid sequences. Additional variant forms of a TGF-beta binding-protein gene are nucleic acid molecules that contain insertions or deletions of the nucleotide sequence described herein. TGF-beta binding-protein variant genes can be identified by determining whether the genes hybridize with a nucleic acid molecule having the nucleotide sequence of SEQ ID Nos: 1, 5, 7, 9, 11, 13, or 15 under stringent conditions. In addition, TGF-beta binding-protein variant genes should encode a protein having a cysteine backbone.

As an alternative, TGF-beta binding-protein variant genes can be identified by sequence comparison. As used herein, two amino acid sequences have "100% amino acid sequence identity" if the amino acid residues of the two amino acid sequences are the same when aligned for maximal correspondence. Similarly, two nucleotide sequences have "100% nucleotide sequence identity" if the nucleotide residues of the two nucleotide sequences are the same when aligned for maximal correspondence. Sequence comparisons can be performed using standard software programs such as those included in the LASERGENE bioinformatics computing suite, which is produced by DNASTAR (Madison, Wis.). Other methods for comparing two nucleotide or amino acid sequences by determining optimal alignment are well-known to those of skill in the art (see, for example, Peruski and Peruski, *The Internet and the New Biology: Tools for Genomic and Molecular Research* (ASM Press, Inc. 1997), Wu et al. (eds.), "Information Superhighway and Computer Databases of Nucleic Acids and Proteins," in Methods in

Gene Biotechnology, pages 123-151 (CRC Press, Inc. 1997), and Bishop (ed.), Guide to Human Genome Computing, 2nd Edition (Academic Press, Inc. 1998)).

A variant TGF-beta binding-protein should have at least a 50% amino acid sequence identity to SEQ ID NOs: 2, 6, 10, 12, 14 or 16 and preferably, greater than 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95% identity. Alternatively, TGF-beta binding-protein variants can be identified by having at least a 70% nucleotide sequence identity to SEQ ID NOs: 1, 5, 9, 11, 13 or 15. Moreover, the present invention contemplates TGF-beta binding-protein gene variants having greater than 75%, 80%, 85%, 90%, or 95% identity to SEQ ID NO:1. Regardless of the particular method used to identify a TGF-beta binding-protein variant gene or variant TGF-beta binding-protein, a variant TGF-beta binding-protein or a polypeptide encoded by a variant TGF-beta binding-protein gene can be functionally characterized by, for example, its ability to bind to and/or inhibit the signaling of a selected member of the TGF-beta family of proteins, or by its ability to bind specifically to an anti-TGF-beta binding-protein antibody.

The present invention include functional fragments of TGF-beta binding-protein genes. Within the context of this invention, a "functional fragment" of a TGF-beta binding-protein gene refers to a nucleic acid molecule that encodes a portion of a TGF-beta binding-protein polypeptide which either (1) possesses the above-noted function activity, or (2) specifically binds with an anti-TGF-beta binding-protein antibody. For example, a functional fragment of a TGF-beta binding-protein gene described herein comprises a portion of the nucleotide sequence of SEQ ID Nos: 1, 5, 9, 11, 13, or 15.

2. Isolation of the TGF-beta Binding-protein Gene

DNA molecules encoding a binding-protein gene can be obtained by screening a human cDNA or genomic library using polynucleotide probes based upon, for example, SEQ ID NO:1.

For example, the first step in the preparation of a cDNA library is to isolate RNA using methods well-known to those of skill in the art. In general, RNA isolation techniques must provide a method for breaking cells, a means of inhibiting RNase-directed degradation of RNA, and a method of separating RNA from DNA, protein, and polysaccharide contaminants. For example, total RNA can be isolated by freezing tissue in liquid nitrogen, grinding the frozen tissue with a mortar and pestle to lyse the cells, extracting the ground tissue with a solution of phenol/chloroform to remove proteins, and separating RNA from the remaining impurities by selective precipitation with lithium chloride (see, for example, Ausubel et al. (eds.), *Short Protocols in Molecular Biology*, 3rd Edition, pages 4-1 to 4-6 (John Wiley & Sons 1995) ["Ausubel (1995)"]; Wu et al., *Methods in Gene Biotechnology*, pages 33-41 (CRC Press, Inc. 1997) ["Wu(1997)"]).

Alternatively, total RNA can be isolated by extracting ground tissue with guanidinium isothiocyanate, extracting with organic solvents, and separating RNA from contaminants using differential centrifugation (see, for example, Ausubel (1995) at pages 4-1 to 4-6, Wu(1997) at pages 33-41).

In order to construct a cDNA library, poly(A)⁺ RNA must be isolated from a total RNA preparation. Poly(A)⁺ RNA can be isolated from total RNA by using the standard technique of oligo(dT)-cellulose chromatography (see, for example, Ausubel (1995) at pages 4-11 to 4-12).

Double-stranded cDNA molecules are synthesized from poly(A)⁺ RNA using techniques well-known to those in the

next:
appear

sequences

sequences

beta

Peruski

breaking

Wu (1997)

4-6

art. (see, for example, Wu (1997) at pages 41-46). Moreover, commercially available kits can be used to synthesize double-stranded cDNA molecules. For example, such kits are available from Life Technologies, Inc. (Gaithersburg, Md.), CLONTECH Laboratories, Inc. (Palo Alto, Calif.), Promega Corporation (Madison, Wis.) and Stratagene Cloning Systems (La Jolla, Calif.).

The basic approach for obtaining TGF-beta binding-protein cDNA clones can be modified by constructing a subtracted cDNA library which is enriched in TGF-binding-protein-specific cDNA molecules. Techniques for constructing subtracted libraries are well-known to those of skill in the art (see, for example, Sargent, "Isolation of Differentially Expressed Genes," in *Meth. Enzymol.* 152:423, 1987, and Wu et al. (eds.), "Construction and Screening of Subtracted and Complete Expression cDNA Libraries," in *Methods in Gene Biotechnology*, pages 29-65 (CRC Press, Inc. 1997)).

Various cloning vectors are appropriate for the construction of a cDNA library. For example, a cDNA library can be prepared in a vector derived from bacteriophage, such as a λ gt10 vector (see, for example, Huynh et al., "Constructing and Screening cDNA Libraries in λ gt10 and λ gt11," in *DNA Cloning: A Practical Approach Vol. I*, Glover (ed.), page 49 (IRL Press, 1985); Wu (1997) at pages 47-52).

Alternatively, double stranded cDNA molecules can be inserted into a plasmid vector, such as a pBluescript vector (Stratagene Cloning Systems; La Jolla, Calif.), a LambdaGEM-4 (Promega Corp.; Madison, Wis.) or other commercially available vectors. Suitable cloning vectors also can be obtained from the American Type Culture Collection (Rockville, Md.).

In order to amplify the cloned cDNA molecules, the cDNA library is inserted into a prokaryotic host, using standard techniques. For example, a cDNA library can be introduced into competent *E. coli* DH5 cell, which can be obtained from Life Technologies, Inc. (Gaithersburg, Md.).

A human genomic DNA library can be prepared by means well-known in the art (see, for example, Ausubel (1995) at pages 5-1 to 5-6; Wu (1997) at pages 307-327). Genomic DNA can be isolated by lysing tissue with the detergent Sarkosyl, digesting the lysate with proteinase K, clearing insoluble debris from the lysate by centrifugation, precipitating nucleic acid from the lysate using isopropanol, and purifying resuspended DNA on a cesium chloride density gradient.

DNA fragments that are suitable for the production of a genomic library can be obtained by the random shearing of genomic DNA or by the partial digestion of genomic DNA with restriction endonucleases. Genomic DNA fragments can be inserted into a vector, such as a bacteriophage or cosmid vector, in accordance with conventional techniques, such as the use of restriction enzyme digestion to provide appropriate termini, the use of alkaline phosphatase treatment to avoid undesirable joining of DNA molecules, and ligation with appropriate ligases. Techniques for such manipulation are well-known in the art (see, for example, Ausubel (1995) at pages 5-1 to 5-6; Wu (1997) at pages 307-327).

Nucleic acid molecules that encode a TGF-beta binding-protein gene can also be obtained using the polymerase chain reaction (PCR) with oligonucleotide primers having nucleotide sequences that are based upon the nucleotide sequences of the human TGF-beta binding-protein gene, as described herein. General methods for screening libraries with PCR are provided by, for example, Yu et al., "Use of the Polymerase Chain Reaction to Screen Phase Libraries," in *Methods in Molecular Biology*, Vol. 15: *PCR Protocols: Current Methods and Applications*, White (ed.), pages

211-215 (Humana Press, Inc. 1993). Moreover, techniques for using PCR to isolate related genes are described by, for example, Preston, "Use of Degenerate Oligonucleotide Primers and the Polymerase Chain Reaction to Clone Gene Family Members," in *Methods in Molecular Biology*, Vol. 15: *PCR Protocols: Current Methods and Applications*, White (ed.), pages 317-337 (Humana Press, Inc. 1993).

Alternatively, human genomic libraries can be obtained from commercial sources such as Research Genetics (Huntsville, Ala.) and the American Type Culture Collection (Rockville, Md.).

A library containing cDNA or genomic clones can be screened with one or more polynucleotide probes based upon SEQ ID NO:1, using standard methods (see, for example, Ausubel (1995) at pages 6-1 to 6-11).

Anti-TGF-beta binding-protein antibodies, produced as described below, can also be used to isolate DNA sequences that encode TGF-beta binding-protein genes from cDNA libraries. For example, the antibodies can be used to screen λ gt11 expression libraries, or the antibodies can be used for immunoscreening following hybrid selection and translation (see, for example, Ausubel (1995) at pages 6-12 to 6-16, Margolis et al., "Screening λ expression libraries with antibody and protein probes," in *DNA Cloning 2: Expression Systems*, 2nd Edition, Glover et al. (eds.), pages 1-14 (Oxford University Press 1995)).

The sequence of a TGF-beta binding-protein cDNA or TGF-beta binding-protein genomic fragment can be determined using standard methods. Moreover, the identification of genomic fragments containing a TGF-beta binding-protein promoter or regulatory element can be achieved using well-established techniques, such as deletion analysis (see, generally, Ausubel (1995)).

As an alternative, a TGF-beta binding-protein gene can be obtained by synthesizing DNA molecules using mutually priming long oligonucleotides and the nucleotide sequences described herein (see, for example, Ausubel (1995) at pages 8-8 to 8-9). Established techniques using the polymerase chain reaction provide the ability to synthesize DNA molecules at least two kilobases in length (Adang et al., *Plant Molec. Biol.* 21:1131, 1993; Bambot et al., *PCR Methods and Applications* 2:266, 1993; Dillon et al., "Use of the Polymerase Chain Reaction for the Rapid Construction of Synthetic Genes," in *Method in Molecular Biology*, Vol. 15: *PCR Protocols: Current Methods and Application*, White (ed.), pages 263-268, (Humana Press, Inc. 1993); Holowachuk et al., *PCR Methods Appl.* 4:299, 1995).

3. Production of TGF-beta Binding-protein Genes

Nucleic acid molecules encoding variant TGF-beta binding-protein genes can be obtained by screening various cDNA or genomic libraries with polynucleotide probes having nucleotide sequence based upon SEQ ID NO:1, 5, 9, 11, 13, or 15, using procedures described above. TGF-beta binding-protein gene variants can also be constructed synthetically. For example, a nucleic acid molecule can be devised that encodes a polypeptide having a conservative amino acid change, compared with the amino acid sequence of SEQ ID NOs. 2, 6, 8, 10, 12, 14, or 16. That is, variants can be obtained that contain one or more amino acid substitutions of SEQ ID NOs: 2, 6, 8, 10, 12, 14 or 16, in which an alkyl amino acid is substituted for an alkyl amino acid in a TGF-beta binding-protein amino acid sequence, an aromatic amino acid is substituted for an aromatic amino acid in a TGF-beta binding-protein amino acid sequence, a sulfur-containing amino acid is substituted for a sulfur-containing amino acid in a TGF-beta binding-protein amino acid sequence, a hydroxy-containing amino acid is substituted

Mol

Rockville

alkaline

○

tuted for a hydroxy-containing amino acid in a TGF-beta binding-protein amino acid sequence, an acidic amino acid is substituted for an acidic amino acid in a TGF-beta binding-protein amino acid sequence, a basic amino acid is substituted for a basic amino acid in a TGF-beta binding-protein amino acid sequence, or a dibasic monocarboxylic amino acid is substituted for a dibasic monocarboxylic amino acid in a TGF-beta binding-protein amino acid sequence.

Among the common amino acids, for example, a "conservative amino acid substitution" is illustrated by a substitution among amino acids within each of the following groups: (1) glycine, alanine, valine, leucine, and isoleucine, (2) phenylalanine, tyrosine, and tryptophan, (3) serine and threonine, (4) aspartate and glutamate, (5) glutamine and asparagine, and (6) lysine, arginine and histidine. In making such substitutions, it is important to, where possible, maintain the cysteine backbone outlined in FIG. 1.

Conservative amino acid changes in a TGF-beta binding-protein gene can be introduced by changing nucleotides for the nucleotides recited in SEQ ID NO:1. Such "conservative amino acid" variants can be obtained, for example, by oligonucleotide-directed mutagenesis, linker-scanning mutagenesis, mutagenesis using the polymerase chain reactions, and the like (see Ausubel (1995) at pages 8-10 to 8-22; and McPherson (ed.), *Directed Mutagenesis: A Practical Approach* (IRL Press 1991)). The functional ability of such variants can be determined using a standard method, such as the assay described herein. Alternatively, a variant TGF-beta binding-protein polypeptide can be identified by the ability to specifically bind anti-TGF-beta binding-protein antibodies.

Routine deletion analyses of nucleic acid molecules can be performed to obtain "functional fragments" of a nucleic acid molecule that encodes a TGF-beta binding-protein polypeptide. As an illustration, DNA molecules having the nucleotide sequence of SEQ ID NO:1 can be digested with Bal31 nuclease to obtain a series of nested deletions. The fragments are then inserted into expression vectors in proper reading frame, and the expressed polypeptides are isolated and tested for activity, or for the ability to bind anti-TGF-beta binding-protein antibodies. One alternative to exonuclease digestion is to use oligonucleotide-directed mutagenesis to introduce deletions or stop codons to specify production of a desired fragment. Alternatively, particular fragments of a TGF-beta binding-protein gene can be synthesized using the polymerase chain reaction.

Standard techniques for functional analysis of proteins are described by, for example, Treuter et al., *Molec. Gen. Genet.* 240:113, 1993; Content et al., "Expression and preliminary deletion analysis of the 42 kDa 2-5A synthetase induced by human interferon," in *Biological Interferon Systems, Proceedings of ISIR-TNO Meeting on Interferon Systems*, Cantell (ed.), pages 65-72 (Nijhoff 1987); Herschman, "The EGF Receptor," in *Control of Animal Cell Proliferation*, Vol. 1, Boynton et al., (eds.) pages 169-199 (Academic Press 1985); Coumilleau et al. *J. Biol. Chem.* 270:29270, 1995; Fukunaga et al., *J. Biol. Chem.* 270:25291, 1995; Yamaguchi et al., *Biochem. Pharmacol.* 50:1295, 1995; and Meisel et al. *Plant Molec. Biol.* 30:1, 1996.

The present invention also contemplates functional fragments of a TGF-beta binding-protein gene that have conservative amino acid changes.

A TGF-beta binding-protein variant gene can be identified on the basis of structure by determining the level of identity with nucleotide and amino acid sequences of SEQ ID NOs: 1, 5, 9, 11, 13, or 15 and 2, 6, 10, 12, 14, or 16, as discussed

above. An alternative approach to identifying a variant gene on the basis of structure is to determine whether a nucleic acid molecule encoding a potential variant TGF-beta binding-protein gene can hybridize under stringent conditions to a nucleic acid molecule having the nucleotide sequence of SEQ ID Nos: 1, 5, 9, 11, 13, or 15, or a portion thereof of at least 15 or 20 nucleotides in length. As an illustration of stringent hybridization conditions, a nucleic acid molecule having a variant TGF-beta binding-protein sequence can bind with a fragment of a nucleic acid molecule having a sequence from SEQ ID NO:1 in a buffer containing, for example, 5xSSPE (1xSSPE=180 mM sodium chloride, 10 mM sodium phosphate, 1 mM EDTA (pH 7.7), 5xDenhardt's solution (100xDenhardt's=2% (w/v) bovine serum albumin, 2% (w/v) Ficoll, 2% (w/v) polyvinylpyrrolidone) and 0.5% SDS incubated overnight at 55-60° C. Post-hybridization washes at high stringency are typically performed in 0.5xSSC (1xSSC=150 mM sodium chloride, 15 mM trisodium citrate) or in 0.5xSSPE at 55-60° C.

Regardless of the particular nucleotide sequence of a variant TGF-beta binding-protein gene, the gene encodes a polypeptide that can be characterized by its functional activity, or by the ability to bind specifically to an anti-TGF-beta binding-protein antibody. More specifically, variant TGF-beta binding-protein genes encode polypeptides which exhibit at least 50%, and preferably, greater than 60, 70, 80 or 90%, of the activity of polypeptides encoded by the human TGF-beta binding-protein gene described herein.

4. Production of TGF-beta Binding-protein in Cultured Cells
To express a TGF-beta binding-protein gene, a nucleic acid molecule encoding the polypeptide must be operably linked to regulatory sequences that control transcriptional expression in an expression vector and then introduced into a host cell. In addition to transcriptional regulatory sequences, such as promoters and enhancers, expression vectors can include translational regulatory sequences and a marker gene which is suitable for selection of cells that carry the expression vector.

Expression vectors that are suitable for production of a foreign protein in eukaryotic cells typically contain (1) prokaryotic DNA elements coding for a bacterial replication origin and an antibiotic resistance marker to provide for the growth and selection of the expression vector in a bacterial host; (2) eukaryotic DNA elements that control initiation of transcription, such as a promoter; and (3) DNA elements that control the processing of transcripts, such as a transcription termination/polyadenylation sequence.

TGF-beta binding-proteins of the present invention are preferably expressed in mammalian cells. Examples of mammalian host cells include African green monkey kidney cells (Vero; ATCC CRI 1587), human embryonic kidney cells (293-HEK; ATCC CRL 1573), baby hamster kidney cells (BHK-21; ATCC CRL8544), canine kidney cells (MDCK; ATCC CCL 34), Chinese hamster ovary cells (CHO-K1; ATCC CCL61), rat pituitary cells (GH1; ATCC CCL82), HeLa S3 cells (ATCC CCL2.2), rat hepatoma cells (H-4-II-E; ATCC CRL 1548) SV40-transformed monkey kidney cells (COS-1; ATCC CRL 1650) and murine embryonic cells (NIH-3T3; ATCC CRL 1658).

For a mammalian host, the transcriptional and translational regulatory signals may be derived from viral sources, such as adenovirus, bovine papilloma virus, simian virus, or the like, in which the regulatory signals are associated with a particular gene which has a high level of expression. Suitable transcriptional and translational regulatory sequences also can be obtained from mammalian genes, such as actin, collagen, myosin, and metallothionein genes.

substituting

deletions

Transcriptional regulatory sequences include a promoter region sufficient to direct the initiation of RNA synthesis. Suitable eukaryotic promoters include the promoter of the mouse metallothionein gene [Hamer et al., *J. Molec. Appl. Genet.* 1:273, 1982], the TK promoter of Herpes virus [McKnight, *Cell* 31:55, 1992], the SV40 early promoter [Benoist et al., *Nature* 290:304, 1981], the Rous sarcoma virus promoter [Gorman et al., *Proc. Nat'l Acad. Sci. USA* 79:6777 1982], the cytomegalovirus promoter [Foecking et al., *Gene* 45:101, 1980], and the mouse mammary tumor virus promoter (see generally, Eicheverry, "Expression of Engineered Proteins, Mammalian Cell Culture," in *Protein Engineering: Principles and Practice*, Cleland et al. (eds.), pages 163-181 (John Wiley & Sons, Inc. 1996)).

Alternatively, a prokaryotic promoter, such as the bacteriophage T3 RNA polymerase promoter, can be used to control TGF-beta binding-protein gene expression in mammalian cells if the prokaryotic promoter is rated by a eukaryotic promoter (Zhou et al., *Mol. Cell. Biol.* 14:529, 1990; Kaufman et al., *Nucl. Acids Res.* 19:4485, 1991).

TGF-beta binding-protein genes may also be expressed in bacterial, yeast, insect or plant cells. Suitable promoters that can be used to express TGF-beta binding-protein polypeptides in a prokaryotic host are well-known to those of skill in the art and include promoters capable of recognizing the T4, T3, Sp6 and T7 polymerases, the P_R and P_L promoters of bacteriophage lambda, the trp, recA, heat shock, lacUV5, tac, lpp-lacSpr, phoA, and lacZ promoters of *E. coli*; promoters of *B. subtilis*, the promoters of the bacteriophages of *Bacillus*, *Streptomyces* promoters, the int promoter of bacteriophage lambda, the bla promoter of pBR322, and the CAT promoter of the chloramphenicol acetyl transferase gene. Prokaryotic promoters have been reviewed by Glick, *J. Ind. Microbiol.* 1:277, 1987; Watson et al., *Molecular Biology of the Gene*, 4th Ed. (Benjamin Cummins 1987), and by Ausubel et al. (1995).

Preferred prokaryotic hosts include *E. coli* and *Bacillus subtilis*. Suitable strains of *E. coli* include BL21(DE3), BL21(DE3)pLysS, BL21(DE3)pLysE, DH1, DH4I, DH5, DH5I, DH5IF, DH5IMCR, DH10B, DH10B/p3, DH11S, C600, HR101, JM101, JM105, JM109, JM110, K38, RR1, Y1088, Y1089, CSH18, ER1451, and ER1647 (see, for example, Brown (Ed.), *Molecular Biology Labfax* (Academic Press 1991)). Suitable strains of *Bacillus subtilis* include BR151, YB886, M1119, M1120, and B170 (see, for example, Hardy, "Bacillus Cloning Methods," in *DNA Cloning: A Practical Approach*, Glover (Ed.) (IRL Press 1985)).

Methods for expressing proteins in prokaryotic hosts are well-known to those of skill in the art (see, for example, Williams et al., "Expression of foreign proteins in *E. coli* using plasmid vectors and purification of specific polyclonal antibodies," in *DNA Cloning 2: Expression Systems, 2nd Edition*, Glover et al. (eds.), page 15 (Oxford University Press 1995); Ward et al., "Genetic Manipulation and Expression of Antibodies," in *Monoclonal Antibodies: Principles and Applications*, page 137 (Wiley-Liss, Inc. 1995); and Georgiou, "Expression of Proteins in Bacteria," in *Protein Engineering: Principles and Practice*, Cleland et al. (eds.), page 101 (John Wiley & Sons, Inc. 1996)).

The baculovirus system provides an efficient means to introduce cloned TGF-beta binding-protein genes into insect cells. Suitable expression vectors are based upon the *Autographa californica* multiple nuclear polyhedrosis virus (AcMNPV), and contain well-known promoters such as *Drosophila* heat shock protein (hsp) 70 promoter, *Autographa californica* nuclear polyhedrosis virus immediate-early gene promoter (ie-1) and the delayed early

39K promoter, baculovirus p10 promoter, and the *Drosophila* metallothionein promoter. Suitable insect host cells include cell lines derived from IPLB-Sf-21, a *Spodoptera frugiperda* pupal ovarian cell line, such as Sf9 (ATCC CRL 1711), Sf21AE, and Sf21 (Invitrogen Corporation; San Diego, Calif.), as well as *Drosophila* Schneider-2 cells. Established techniques for producing recombinant proteins in baculovirus systems are provided by Bailey et al., "Manipulation of Baculovirus Vectors," in *Methods in Molecular Biology, Volume 7: Gene Transfer and Expression Protocols*, Murray (ed.), pages 147-168 (The Humana Press, Inc. 1991), by Patel et al., "The baculovirus expression system," in *DNA Cloning 2: Expression Systems, 2nd Edition*, Glover et al. (eds.), pages 205-244 (Oxford University Press 1995), by Ausubel (1995) at pages 16-37 to 16-57, by Richardson (ed.), *Baculovirus Expression Protocols* (The Humana Press, Inc. 1995), and by Lucknow, "Insect Cell Expression Technology," in *Protein Engineering: Principles and Practice*, Cleland et al. (eds.), pages 183-218 (John Wiley & Sons, Inc. 1996).

Promoters for expression in yeast include promoters from GAL1 (galactose), PGK (phosphoglycerate kinase), ADH (alcohol dehydrogenase), AOX1 (alcohol oxidase), HIS4 (histidinol dehydrogenase), and the like. Many yeast cloning vectors have been designed and are readily available. Those vectors include Ylp-based vectors, such as Ylp5, YRp vectors, such as YRp17, YEpl3 and YCp vectors, such as YCp19. One skilled in the art will appreciate that there are a wide variety of suitable vectors for expression in yeast cells.

Expression vectors can also be introduced into plant protoplasts, intact plant tissues, or isolated plant cells. General methods of culturing plant tissues are provided, for example, by Miki et al., "Procedures for Introducing Foreign DNA into Plants," in *Methods in Plant Molecular Biology and Biotechnology*, Glick et al. (eds.), pages 67-88 (CRC Press, 1993).

An expression vector can be introduced into host cells using a variety of standard techniques including calcium phosphate transfection, liposome-mediated transfection, microprojectile-mediated delivery, electroporation, and the like. Preferably, the transfected cells are selected and propagated to provide recombinant host cells that comprise the expression vector stably integrated in the host cell genome. Techniques for introducing vectors into eukaryotic cells and techniques for selecting such stable transformants using a dominant selectable marker are described, for example, by Ausubel (1995) and by Murray (ed.), *Gene Transfer and Expression Protocols* (Humana Press 1991). Methods for introducing expression vectors into bacterial, yeast, insect, and plant cells are also provided by Ausubel (1995).

General methods for expressing and recovering foreign protein produced by a mammalian cell system is provided by, for example, Eicheverry, "Expression of Engineered Proteins in Mammalian Cell Culture," in *Protein Engineering: Principles and Practice*, Cleland et al. (eds.), pages 163 (Wiley-Liss, Inc. 1996). Standard techniques for recovering protein produced by a bacterial system is provided by, for example, Grisshammer et al., "Purification of over-produced proteins from *E. coli* cells," in *DNA Cloning 2: Expression Systems, 2nd Edition*, Glover et al. (eds.), pages 59-92 (Oxford University Press 1995). Established methods for isolating recombinant proteins from a baculovirus system are described by Richardson (ed.), *Baculovirus Expression Protocols* (The Humana Press, Inc., 1995).

More generally, TGF-beta binding-protein can be isolated by standard techniques, such as affinity chromatography,

region
metalothionein
insect in
regulated
no hypoxan
no hypoxan
HB101
proteins
heat

mediated delivery
cell

those
size exclusion chromatography, ion exchange chromatography, HPLC and the like. Additional variations in TGF-beta binding-protein isolation and purification can be devised by those of skill in the art. For example, anti-TGF-beta binding-protein antibodies, obtained as described below, can be used to isolate large quantities of protein by immunoaffinity purification.

5. Production of Antibodies to TGF-beta Binding-proteins.

Antibodies to TGF-beta binding-protein can be obtained, for example, using the product of an expression vector as an antigen. Particularly useful anti-TGF-beta binding-protein antibodies "bind specifically" with TGF-beta binding-protein of Sequence ID Nos. 2, 6, 10, 12, 14, or 16, but not to other TGF-beta binding-proteins such as Dan, Cerberus, SEGF, or Gremlin. Antibodies of the present invention (including fragments and derivatives thereof) may be a polyclonal or, especially a monoclonal antibody. The antibody may belong to any immunoglobulin class, and may be for example an IgG, for example IgG₁, IgG₂, IgG₃, IgG₄, IgE, IgM; or IgA antibody. It may be of animal, for example mammalian origin, and may be for example a murine, rat, human or other primate antibody. Where desired the antibody may be an internalising antibody.

polyclonal proteins
Polyclonal antibodies to recombinant TGF-beta binding-protein can be prepared using methods well-known to those of skill in the art (see, for example, Green et al., "Production of Polyclonal Antisera," in *Immunochemical Protocols* (Manson, ed.), pages 1-5 (Humana Press 1992); Williams et al., "Expression of foreign proteins in *E. coli* using plasmid vectors and purification of specific polyclonal antibodies," in *DNA Cloning 2: Expression Systems, 2nd Edition*, Glover et al. (eds.), page 15 (Oxford University Press 1995)). Although polyclonal antibodies are typically raised in animals such as rats, mice, rabbits, goats, or sheep, an anti-TGF-beta binding-protein antibody of the present invention may also be derived from a subhuman primate antibody. General techniques for raising diagnostically and therapeutically useful antibodies in baboons may be found, for example, in Goldenberg et al., international patent publication No. WO 91/11465 (1991), and in Losman et al., *Int. J. Cancer* 46:310, 1990.

The antibody should comprise at least a variable region domain. The variable region domain may be of any size or amino acid composition and will generally comprise at least one hypervariable amino acid sequence responsible for antigen binding embedded in a framework sequence. In general terms the variable (V) region domain may be any suitable arrangement of immunoglobulin heavy (V_H) and/or light (V_L) chain variable domains. Thus for example the V region domain may be monomeric and be a V_H or V_L domain where these are capable of independently binding antigen with acceptable affinity. Alternatively the V region domain may be dimeric and contain V_H-V_H, V_H-V_L, or V_L-V_L dimers in which the V_H and V_L chains are non-covalently associated (abbreviated hereinafter as F_v). Where desired, however, the chains may be covalently coupled either directly for example via a disulfide bond between the two variable domains, or through a linker, for example a peptide linker, to form a single chain domain (abbreviated hereinafter as scF_v).

The variable region domain may be any naturally occurring variable domain or an engineered version thereof. By engineered version is meant a variable region domain which has been created using recombinant DNA engineering techniques. Such engineered versions include those created for example from natural antibody variable regions by insertions, deletions or changes in or to the amino acid

sequences of the natural antibodies. Particular examples of this type include those engineered variable region domains containing at least one CDR and optionally one or more framework amino acids from one antibody and the remainder of the variable region domain from a second antibody.

The variable region domain may be covalently attached at a C-terminal amino acid to at least one other antibody domain or a fragment thereof. Thus, for example where a V_H domain is present in the variable region domain this may be linked to an immunoglobulin C_H1 domain or a fragment thereof. Similarly a V_L domain may be linked to a C_K domain or a fragment thereof. In this way for example the antibody may be a Fab fragment wherein the antigen binding domain contains associated V_H and V_L domains covalently linked at their C-termini to a CH1 and C_K domain respectively. The CH1 domain may be extended with further amino acids, for example to provide a hinge region domain as found in a Fab' fragment, or to provide further domains, such as antibody CH2 and CH3 domains.

Another form of an antibody fragment is a peptide coding for a single complementarity-determining region (CDR). CDR peptides ("minimal recognition units") can be obtained by constructing genes encoding the CDR of an antibody of interest. Such genes are prepared, for example, by using the polymerase chain reaction to synthesize the variable region from RNA of antibody-producing cells (see, for example, Larrick et al., *Methods: A Companion to Methods in Enzymology* 2:106, 1991; Courtenay-Luck, "Genetic Manipulation of Monoclonal Antibodies," in *Monoclonal Antibodies: Production, Engineering and Clinical Application*, Ritter et al. (eds.), page 166 (Cambridge University Press 1995); and Ward et al., "Genetic Manipulation and Expression of Antibodies," in *Monoclonal Antibodies: Principles and Applications*, Birch et al., (eds.), page 137 (Wiley-Liss, Inc. 1995)).

Antibodies for use in the invention may in general be monoclonal (prepared by conventional immunisation and cell fusion procedures) or in the case of fragments, derived therefrom using any suitable standard chemical e.g. reduction or enzymatic cleavage and/or digestion techniques, for example by treatment with pepsin.

More specifically, monoclonal anti-TGF-beta binding-protein antibodies can be generated utilizing a variety of techniques. Rodent monoclonal antibodies to specific antigens may be obtained by methods known to those skilled in the art (see, for example, Kohler et al., *Nature* 256:495, 1975; and Coligan et al. (eds.), *Current Protocols in Immunology*, 1:2.5.1-2.6.7 (John Wiley & Sons 1991) ["Coligan"]; Picklesley et al., "Production of monoclonal antibodies against proteins expressed in *E. coli*," in *DNA Cloning 2: Expression Systems, 2nd Edition*, Glover et al. (eds.), page 93 (Oxford University Press 1995)).

Briefly, monoclonal antibodies can be obtained by injecting mice with a composition comprising a TGF-beta binding-protein gene product, verifying the presence of antibody production by removing a serum sample, removing the spleen to obtain B-lymphocytes, fusing the B-lymphocytes with myeloma cells to produce hybridomas, cloning the hybridomas, selecting positive clones which produce antibodies to the antigen, culturing the clones that produce antibodies to the antigen, and isolating the antibodies from the hybridoma cultures.

In addition, an anti-TGF-beta binding-protein antibody of the present invention may be derived from a human monoclonal antibody. Human monoclonal antibodies are obtained from transgenic mice that have been engineered to produce specific human antibodies in response to antigenic chal-

missing word =

antibody

leage. In this technique, elements of the human heavy and light chain locus are introduced into strains of mice derived from embryonic stem cell lines that contain targeted disruptions of the endogenous heavy chain and light chain loci. The transgenic mice can synthesize human antibodies specific for human antigens, and the mice can be used to produce human antibody-secreting hybridomas. Methods for obtaining human antibodies from transgenic mice are described, for example, by Green et al., *Nature Genet.* 7:13, 1994; Lonberg et al., *Nature* 368:856, 1994; and Taylor et al., *Int. Immun.* 6:579, 1994.

Monoclonal antibodies can be isolated and purified from hybridoma cultures by a variety of well-established techniques. Such isolation techniques include affinity chromatography with Protein-A Sepharose, size-exclusion chromatography, and ion-exchange chromatography (see, for example, Coligan at pages 2.7.1-2.7.12 and pages 2.9.1-2.9.3; Baines et al., "Purification of Immunoglobulin G (IgG)," in *Methods in Molecular Biology*, Vol. 10, pages 79-104 (The Humana Press, Inc. 1992)).

For particular uses, it may be desirable to prepare fragments of anti-TGF-beta binding-protein antibodies. Such antibody fragments can be obtained, for example, by proteolytic hydrolysis of the antibody. Antibody fragments can be obtained by pepsin or papain digestion of whole antibodies by conventional methods. As an illustration, antibody fragments can be produced by enzymatic cleavage of antibodies with pepsin to provide a 5S fragment denoted F(ab')₂. This fragment can be further cleaved using a thiol reducing agent to produce 3.5S Fab' monovalent fragments. Optionally, the cleavage reaction can be performed using a blocking group for the sulfhydryl groups that result from cleavage of disulfide linkages. As an alternative, an enzymatic cleavage using pepsin produces two monovalent Fab fragments and an Fc fragment directly. These methods are described, for example, by Goldenberg, U.S. Pat. No. 4,331, 647, Nisonoff et al., *Arch Biochem. Biophys.* 89:230, 1960, Porter, *Biochem. J.* 73:119, 1959, Edelman et al., in *Methods in Enzymology* 1:422 (Academic Press 1967), and by Coligan at pages 2.8.1-2.8.10 and 2.10.-2.10.4.

Other methods of cleaving antibodies, such as separation of heavy chains to form monovalent light-heavy chain fragments, further cleavage of fragments, or other enzymatic, chemical or genetic techniques may also be used, so long as the fragments bind to the antigen that is recognized by the intact antibody.

Alternatively, the antibody may be a recombinant or engineered antibody obtained by the use of recombinant DNA techniques involving the manipulation and re-expression of DNA encoding antibody variable and/or constant regions. Such DNA is known and/or is readily available from DNA libraries including for example phage-antibody libraries (see Chiswell, D J and McCafferty, *J. Tibtech* 10 80-84 (1992)) or where desired can be synthesised. Standard molecular biology and/or chemistry procedures may be used to sequence and manipulate the DNA, for example, to introduce codons to create cysteine residues, to modify, add or delete other amino acids or domains as desired.

From here, one or more replicable expression vectors containing the DNA may be prepared and used to transform an appropriate cell line, e.g. a non-producing myeloma cell line, such as a mouse NSO line or a bacterial, e.g. *E. coli* line in which production of the antibody will occur. In order to obtain efficient transcription and translation, the DNA sequence in each vector should include appropriate regulatory sequences, particularly a promoter and leader sequence operably linked to the variable domain sequence. Particular

methods for producing antibodies in this way are generally well known and routinely used. For example, basic molecular biology procedures are described by Maniatis et al (Molecular Cloning, Cold Spring Harbor Laboratory, New York, 1989); DNA sequencing can be performed as described in Sanger et al (PNAS 74, 5463, (1977)) and the Amersham International plc sequencing handbook; and site directed mutagenesis can be carried out according to the method of Kramer et al (Nucl. Acids Res. 12, 9441, (1984)) and the Anglian-Biotechnology Ltd handbook. Additionally, there are numerous publications, detailing techniques suitable for the preparation of antibodies by manipulation of DNA, creation of expression vectors and transformation of appropriate cells, for example as reviewed by Mountain A and Adair, J R in Biotechnology and Genetic Engineering Reviews (ed. Tombs, M P, 10, Chapter 1, 1992, Intercept, Andover, UK) and in International Patent Specification No. WO 91/09967.

Where desired, the antibody according to the invention may have one or more effector or reporter molecules attached to it and the invention extends to such modified proteins. The effector or reporter molecules may be attached to the antibody through any available amino acid side-chain, terminal amino acid or, where present carbohydrate functional group located in the antibody, always provided of course that this does not adversely affect the binding properties and eventual usefulness of the molecule. Particular functional groups include, for example any free amino, imino, thiol, hydroxyl, carboxyl or aldehyde group. Attachment of the antibody and the effector and/or reporter molecule(s) may be achieved via such groups and an appropriate functional group in the effector or reporter molecules. The linkage may be direct or indirect, through spacing or bridging groups.

Effector molecules include, for example, antineoplastic agents, toxins (such as enzymatically active toxins of bacterial or plant origin and fragments thereof e.g. ricin and fragments thereof) biologically active proteins, for example enzymes, nucleic acids and fragments thereof, e.g. DNA, RNA and fragments thereof, naturally occurring and synthetic polymers e.g. polysaccharides and polyalkylene polymers such as poly(ethylene glycol) and derivatives thereof, radionuclides, particularly radioiodide, and chelated metals. Suitable reporter groups include chelated metals, fluorescent compounds or compounds which may be detected by NMR or ESR spectroscopy.

Particular antineoplastic agents include cytotoxic and cytostatic agents, for example alkylating agents, such as nitrogen mustards (e.g. chlorambucil, melphalan, mechlorethamine, cyclophosphamide, or uracil mustard) and derivatives thereof, triethylenephosphoramide, triethylenethiophosphoramide, busulphan, or cisplatin; antimetabolites, such as methotrexate, fluorouracil, floxuridine, cytarabine, mercaptopurine, thioguanine, fluoracetic acid or fluorocitric acid, antibiotics, such as bleomycins (e.g. bleomycin sulphate), doxorubicin, daunorubicin, mitomycins (e.g. mitomycin C), actinomycins (e.g. dactinomycin) plicamycin, calicheamicin and derivatives thereof, or esperamicin and derivatives thereof; mitotic inhibitors, such as etoposide, vincristine or vinblastine and derivatives thereof; alkaloids, such as ellipticine; polyols such as taxicin-I or taxicin-II; hormones, such as androgens (e.g. dromostanolone or testolactone), progestins (e.g. megestrol acetate or medroxyprogesterone acetate), estrogens (e.g. dimethylstilbestrol diphosphate, polyestradiol phosphate or estramustine phosphate) or antiestrogens (e.g. tamoxifen); anthraquinones, such as mitoxantrone, ureas, such as

delete comma

are preparation

Methods

cyclophospham

delete comma

Calicheamicin

insert comma

hydroxyurea; hydrazines, such as procarbazine; or imidazoles, such as dacarbazine.

Particularly useful effector groups are calicheamicin and derivatives thereof (see for example South African Patent Specifications Nos. 85/8794, 88/8127 and 90/2839).

Chelated metals include chelates of di- or tripositive metals having a coordination number from 2 to 8 inclusive. Particular examples of such metals include technetium (Tc), rhenium (Re), cobalt (Co), copper (Cu), gold (Au), silver (Ag), lead (Pb), bismuth (Bi), indium (In), gallium (Ga), yttrium (Y), terbium (Tb), gadolinium (Gd), and scandium (Sc). In general the metal is preferably a radionuclide. Particular radionuclides include ^{99m}Tc , ^{188}Re , ^{186}Re , ^{58}Co , ^{60}Co , ^{67}Cu , ^{195}Au , ^{199}Au , ^{110}Ag , ^{203}Pb , ^{206}Bi , ^{207}Bi , ^{111}In , ^{67}Ga , ^{68}Ga , ^{88}Y , ^{90}Y , ^{160}Tb , ^{153}Gd and ^{47}Sc .

The chelated metal may be for example one of the above types of metal chelated with any suitable polydentate chelating agent, for example acyclic or cyclic polyamines, polyethers, (e.g. crown ethers and derivatives thereof); polyamides; porphyrins; and carbocyclic derivatives.

In general, the type of chelating agent will depend on the metal in use. One particularly useful group of chelating agents in conjugates according to the invention, however, are acyclic and cyclic polyamines, especially polyamino-carboxylic acids, for example diethylenetriamine-pentaacetic acid and derivatives thereof, and macrocyclic amines, e.g. cyclic tri-aza and tetra-aza derivatives (for example as described in International Patent Specification No. WO 92/22583); and polyamides, especially desferrioxamine and derivatives thereof.

Thus for example when it is desired to use a thiol group in the antibody as the point of attachment this may be achieved through reaction with a thiol reactive group present in the effector or reporter molecule. Examples of such groups include an α -halocarboxylic acid or ester, e.g. iodoacetamide, an imide, e.g. maleimide, a vinyl sulphone, or a disulfide. These and other suitable linking procedures are generally and more particularly described in International Patent Specifications Nos. WO 93/06231, WO 92/22583, WO 90/091195 and WO 89/01476.

Assays for Selecting Molecules Which Increase Bone Density

As discussed above, the present invention provides methods for selecting and/or isolating compounds which are capable of increasing bone density. For example, within one aspect of the present invention methods are provided for determining whether a selected molecule is capable of increasing bone mineral content, comprising the steps of (a) mixing a selected molecule with TGF-beta binding protein and a selected member of the TGF-beta family of proteins, (b) determining whether the selected molecule stimulates signaling by the TGF-beta family of proteins, or inhibits the binding of the TGF-beta binding protein to the TGF-beta family of proteins. Within certain embodiments, the molecule enhances the ability of TGF-beta to function as a positive regulator of mesenchymal cell differentiation.

Within other aspects of the invention, methods are provided for determining whether a selected molecule is capable of increasing bone mineral content, comprising the steps of (a) exposing a selected molecule to cells which express TGF-beta binding-protein from and (b) determining whether the expression (or activity) of TGF-beta binding-protein from said exposed cells decreases, and therefrom determining whether the compound is capable of increasing bone mineral content. Within one embodiment, the cells are selected from the group consisting of the spontaneously

transformed or untransformed normal human bone from bone biopsies and rat parietal bone osteoblasts. Such methods may be accomplished in a wide variety of assay formats including, for example, Countercurrent Immuno-Electrophoresis (CIEP), Radioimmunoassays, Radioimmunoprecipitations, Enzyme-Linked Immunosorbent Assays (ELISA), Dot Blot assays, Inhibition or Competition assays, and sandwich assays (see U.S. Pat. Nos. 4,376,110 and 4,486,530; see also *Antibodies: A Laboratory Manual*, supra).

Representative embodiments of such assays are provided below in Examples 5 and 6. Briefly, a family member of the TGF-beta super-family or a TGF-beta binding protein is first bound to a solid phase, followed by addition of a candidate molecule. The labeled family member of the TGF-beta super-family or a TGF-beta binding protein is then added to the assay, the solid phase washed, and the quantity of bound or labeled TGF-beta super-family member or TGF-beta binding protein on the solid support determined. Molecules which are suitable for use in increasing bone mineral content as described herein are those molecules which decrease the binding of TGF-beta binding protein to a member or members of the TGF-beta super-family in a statistically significant manner. Obviously, assays suitable for use within the present invention should not be limited to the embodiments described within Examples 2 and 3. In particular, numerous parameters may be altered, such as by binding TGF-beta to a solid phase, or by elimination of a solid phase entirely.

Within other aspects of the invention, methods are provided for determining whether a selected molecule is capable of increasing bone mineral content, comprising the steps of (a) exposing a selected molecule to cells which express TGF-beta and (b) determining whether the activity of TGF-beta from said exposed cells is altered, and therefrom determining whether the compound is capable of increasing bone mineral content. Similar to the above described methods, a wide variety of methods may be utilized to assess the changes of TGF-beta binding-protein expression due to a selected test compound.

For example, within one aspect of the present invention methods are provided for determining whether a selected molecule is capable of increasing bone mineral content, comprising the steps of (a) mixing a selected molecule with TGF-beta binding-protein and a selected member of the TGF-beta family of proteins, (b) determining whether the selected molecule up-regulates the signaling of the TGF-beta family of proteins, or inhibits the binding of the TGF-beta binding-protein to the TGF-beta family of proteins. Within certain embodiments, the molecule enhances the ability of TGF-beta to function as a positive regulator of mesenchymal cell differentiation.

Similar to the above described methods, a wide variety of methods may be utilized to assess stimulation of TGF-beta due to a selected test compound. One such representative method is provided below in Example 6 (see also Durham et al., *Endo.* 136:1374-1380).

Within yet other aspects of the present invention, methods are provided for determining whether a selected molecule is capable of increasing bone mineral content, comprising the step of determining whether a selected molecule inhibits the binding of TGF-beta binding-protein to bone, or an analogue thereof. As utilized herein, it should be understood that bone or analogues thereof refers to hydroxyapatite, or a surface composed of a powdered form of bone, crushed bone or intact bone. Similar to the above described methods, a wide variety of methods may be utilized to assess the inhibition of

derivatives

may be in italics

macrocyclic

mesenchymal

(1380)

are

TGF-beta binding-protein localization to bone matrix. One such representative method is provided below in Example 7.

It should be noted that while the methods recited herein may refer to the analysis of an individual test molecule, that the present invention should not be so limited. In particular, the selected molecule may be contained within a mixture of compounds. Hence, the recited methods may further comprise the step of isolating a molecule which inhibits the binding of TGF-beta binding-protein to a TGF-beta family member.

Candidate Molecules

A wide variety of molecules may be assayed for their ability to inhibit the binding of TGF-beta binding-protein to a TGF-beta family member. Representative examples which are discussed in more detail below include organic molecules, proteins or peptides, and nucleic acid molecules. Although it should be evident from the discussion below that the candidate molecules described herein may be utilized in the assays described herein, it should also be readily apparent that such molecules can also be utilized in a variety of diagnostic and therapeutic settings.

1. Organic Molecules

Numerous organic molecules may be assayed for their ability to inhibit the binding of TGF-beta binding-protein to a TGF-beta family member.

For example, within one embodiment of the invention suitable organic molecules may be selected from either a chemical library, wherein chemicals are assayed individually, or from combinatorial chemical libraries where multiple compounds are assayed at once, then deconvoluted to determine and isolate the most active compounds.

Representative examples of such combinatorial chemical libraries include those described by Agrafiotis et al., "System and method of automatically generating chemical compounds with desired properties," U.S. Pat. No. 5,463,564; Armstrong, R. W., "Synthesis of combinatorial arrays of organic compounds through the use of multiple component combinatorial arrays syntheses," WO 95/2566; Baldwin, J. J. et al., "Sulfonamide derivatives and their use," WO 95/24186; Baldwin, J. J. et al., "Combinatorial dihydrobenzopyran library," WO 95/30642; Brenner, S., "New kit for preparing combinatorial libraries," WO 95/16918; Chenery, B. et al., "Preparation of library of resin-bound aromatic carbocyclic compounds," WO 95/16712; Ellman J. A., "Solid phase and combinatorial synthesis of benzodiazepine compounds on a solid support," U.S. Pat. No. 5,288,514; Felder, E. et al., "Novel combinatorial compound libraries," WO 95/16209; Lerner, R. et al., "Encoded combinatorial chemical libraries," WO 93/20242; Pavia, M. R. et al., "A method for preparing and selecting pharmaceutically useful non-peptide compounds from a structurally diverse universal library," WO 95/04277; Summerton, J. E. and D. D. Weller, "Morpholino-subunit combinatorial library and method," U.S. Pat. No. 5,506,337; Holmes, C., "Methods for the Solid Phase Synthesis of Thiazolidinones, Metathiazanones, and Derivatives thereof," WO 96/00148; Phillips, G. B. and G. C. P. Wei, "Solid-phase Synthesis of Benzimidazoles," *Tet. Letters* 37:4887-90, 1996; Ruhland, B. et al., "Solid-supported Combinatorial Synthesis of Structurally Diverse β -Lactams," *J. Amer. Chem. Soc.* 111:253-4, 1996; Look, G. C. et al., "The Identification of Cyclooxygenase-1 Inhibitors from 4-Thiazolidinone Combinatorial Libraries," *Bioorg and Med. Chem. Letters* 6:707-12, 1996.

2. Proteins and Peptides

A wide range of proteins and peptides may likewise be utilized as candidate molecules for inhibitors of the binding of TGF-beta binding-protein to a TGF-beta family member.

a. Combinatorial Peptide Libraries

Peptide molecules which are putative inhibitors of the binding of TGF-beta binding-protein to a TGF-beta family member may be obtained through the screening of combinatorial peptide libraries. Such libraries may either be prepared by one of skill in the art (see e.g., U.S. Pat. Nos. 4,528,266 and 4,359,535, and Patent Cooperation Treaty Publication Nos. WO 92/15679, WO 92/15677, WO 90/07862, WO 90/02809, or purchased from commercially available sources (e.g., New England Biolabs Ph.D.TM Phage Display Peptide Library Kit).

b. Antibodies

Antibodies which inhibit the binding of TGF-beta binding-protein to a TGF-beta family member may readily be prepared given the disclosure provided herein. Within the context of the present invention, antibodies are understood to include monoclonal antibodies, polyclonal antibodies, anti-idiotypic antibodies, antibody fragments (e.g., Fab, and F(ab')₂, F₁, variable regions, or complementarity determining regions). As discussed above, antibodies are understood to be specific against TGF-beta binding-protein, or against a specific TGF-beta family member, if they bind with a K_d of greater than or equal to 10⁻⁷M, preferably greater than or equal to 10⁻⁸M, and do not bind to other TGF-beta binding-proteins, or bind with a K_d of less than or equal to 10⁻⁶M. Furthermore, antibodies of the present invention should block or inhibit the binding of TGF-beta binding-protein to a TGF-beta family member.

The affinity of a monoclonal antibody or binding partner, as well as inhibition of binding can be readily determined by one of ordinary skill in the art (see Scatchard, *Ann. N.Y. Acad. Sci.* 51:660-67 1949).

Briefly, polyclonal antibodies may be readily generated by one of ordinary skill in the art from a variety of warm-blooded animal such as horses, cows, various fowl, rabbits, mice, or rats. Typically, the TGF-beta binding-protein or unique peptide thereof of 13-20 amino acids (preferably conjugated to keyhole limpet hemocyanin by cross-linking with glutaraldehyde) is utilized to immunize the animal through intraperitoneal, intramuscular, intraocular, or subcutaneous injections, along with an adjuvant such as Freund's complete or incomplete adjuvant. Following several booster immunizations, samples of serum are collected and tested for reactivity to the protein or peptide. Particularly preferred polyclonal antisera will give a signal on one of these assays that is at least three times greater than background. Once the titer of the animal has reached a plateau in terms of its reactivity to the protein, larger quantities of antisera may be readily obtained either by weekly bleedings, or by exsanguinating the animal.

Monoclonal antibodies may also be readily generated using conventional techniques (see U.S. Pat. Nos. RE 32,011, 4,902,614, 4,543,439, and 4,411,993 which are incorporated herein by reference; see also *Monoclonal Antibodies, Hybridomas: A New Dimension in Biological Analyses*, Plenum Press, Kennett, McKeam, and Bechtol (eds.), 1980, and *Antibodies: A Laboratory Manual*, Harlow and Lane (eds.), Cold Spring Harbor Laboratory Press, 1988, which are also incorporated herein by reference).

Briefly, within one embodiment a subject animal such as a rat or mouse is immunized with TGF-beta binding-protein or portion thereof as described above. The protein may be admixed with an adjuvant such as Freund's complete or incomplete adjuvant in order to increase the resultant immune response. Between one and three weeks after the initial immunization the animal may be reimmunized with another booster immunization, and tested for reactivity to the

array
Preparation
Benzimidazole

prepared

polyclonal

immunization

protein utilizing assays described above. Once the animal has reached a plateau in its reactivity to the injected protein, it is sacrificed and organs which contain large numbers of B cells such as the spleen and lymph nodes are harvested.

Cells which are obtained from the immunized animal may be immortalized by infection with a virus such as the Epstein-Barr virus (EBV) (see Glasky and Reading, *Hybridoma* 8(4):377-389, 1989). Alternatively, within a preferred embodiment, the harvested spleen and/or lymph node cell suspensions are fused with a suitable myeloma cell in order to create a "hybridoma" which secretes monoclonal antibody. Suitable myeloma lines include, for example, NS-1 (ATCC No. TIB 18), and P3x63-Ag 8.653 (ATCC No. CRL 1580).

Following the fusion, the cells may be placed into culture plates containing a suitable medium, such as RPMI 1640, or DMEM (Dulbecco's Modified Eagles Medium) (JRH Biosciences, Lenexa, Kans.), as well as additional ingredients, such as fetal bovine serum (FBS, i.e., from Hyclone, Logan, Utah, or JRH Biosciences). Additionally, the medium should contain a reagent which selectively allows for the growth of fused spleen and myeloma cells such as HAT (hypoxanthine, aminopterin, and thymidine) (Sigma Chemical Co., St Louis, Miss.). After about seven days, the resulting fused cells or hybridomas may be screened in order to determine the presence of antibodies which are reactive against TGF-beta binding-protein (depending on the antigen used), and which block or inhibit the binding of TGF-beta binding-protein to a TGF-beta family member.

A wide variety of assays may be utilized to determine the presence of antibodies which are reactive against the proteins of the present invention, including for example counter-current immuno-electrophoresis, radioimmunoassays, radioimmunoprecipitations, enzyme-linked immunosorbent assays (ELISA), dot blot assays, western blots, immunoprecipitation, inhibition or competition assays, and sandwich assays (see U.S. Pat. Nos. 4,376,110 and 4,486,530; see also *Antibodies: A Laboratory Manual*, Harlow and Lane (eds.), Cold Spring Harbor Laboratory Press, 1988). Following several clonal dilutions and reassays, a hybridoma producing antibodies reactive against the desired protein may be isolated.

Other techniques may also be utilized to construct monoclonal antibodies (see William D. Huse et al., "Generation of a Large Combinational Library of the Immunoglobulin Repertoire in Phage-Lambda," *Science* 246:1275-1281, December 1989; see also L. Sastry et al., "Cloning of the Immunological Repertoire in *Escherichia coli* for Generation of Monoclonal Catalytic Antibodies: Construction of a Heavy Chain Variable Region-Specific cDNA Library," *Proc. Natl. Acad. Sci. USA* 86:5728-5732, August 1989; see also Michelle Altling-Mees et al., "Monoclonal Antibody Expression Libraries: A Rapid Alternative to Hybridomas," *Strategies in Molecular Biology* 3:1-9, January 1990). These references describe a commercial system available from Stratagene (La Jolla, Calif.) which enables the production of antibodies through recombinant techniques. Briefly, mRNA is isolated from a B cell population, and utilized to create heavy and light chain immunoglobulin cDNA expression libraries in the λ ImmunoZap(H) and λ ImmunoZap(L) vectors. These vectors may be screened individually or co-expressed to form Fab fragments or antibodies (see Huse et al., supra; see also Sastry et al., supra). Positive plaques may subsequently be converted to a non-lytic plasmid which allows high level expression of monoclonal antibody Feats from *E. coli*.

Similarly, portions or fragments, such as Fab and Fv fragments, of antibodies may also be constructed utilizing

conventional enzymatic digestion or recombinant DNA techniques to incorporate the variable regions of a gene which encodes a specifically binding antibody. Within one embodiment, the genes which encode the variable region from a hybridoma producing a monoclonal antibody of interest are amplified using nucleotide primers for the variable region. These primers may be synthesized by one of ordinary skill in the art, or may be purchased from commercially available sources. (Stratagene (La Jolla, Calif.) sells primers for mouse and human variable regions including, among others primers for $V_{H\alpha}$, $V_{H\beta}$, $V_{H\gamma}$, $V_{H\delta}$, $C_{H\alpha}$, V_L and C_L regions. These primers may be utilized to amplify heavy or light chain variable regions, which may then be inserted into vectors such as ImmunoZAP™ H or ImmunoZAP™ L (Stratagene), respectively. These vectors may then be introduced into *E. coli*, yeast, or mammalian-based systems for expression. Utilizing these techniques, large amounts of a single-chain protein containing a fusion of the V_H and V_L domains may be produced (see Bird et al., *Science* 242:423-426, 1988). In addition, such techniques may be utilized to change a "murine" antibody to a "human" antibody, without altering the binding specificity of the antibody.

Once suitable antibodies have been obtained, they may be isolated or purified by many techniques well known to those of ordinary skill in the art (see *Antibodies: A Laboratory Manual*, Harlow and Lane (eds.), Cold Spring Harbor Laboratory Press, 1988). Suitable techniques include peptide or protein affinity columns, HPLC or RP-HPLC, purification on protein A or protein G columns, or any combination of these techniques.

c. Mutant TGF-beta Binding-proteins

As described herein and below in the Examples (e.g., Examples 8 and 9), altered versions of TGF-beta binding-protein which compete with native TGF-beta binding-protein's ability to block the activity of a particular TGF-beta family member should lead to increased bone density. Thus, mutants of TGF-beta binding-protein which bind to the TGF-beta family member but do not inhibit the function of the TGF-beta family member would meet the criteria. The mutant versions must effectively compete with the endogenous inhibitory functions of TGF-beta binding-protein.

d. Production of Proteins

Although various genes (or portions thereof) have been provided herein, it should be understood that within the context of the present invention, reference to one or more of these genes includes derivatives of the genes that are substantially similar to the genes (and, where appropriate, the proteins (including peptides and polypeptides) that are encoded by the genes and their derivatives). As used herein, a nucleotide sequence is deemed to be "substantially similar" if: (a) the nucleotide sequence is derived from the coding region of the above-described genes and includes, for example, portions of the sequence or allelic variations of the sequences discussed above, or alternatively, encodes a molecule which inhibits the binding of TGF-beta binding-protein to a member of the TGF-beta family, (b) the nucleotide sequence is capable of hybridization to nucleotide sequences of the present invention under moderate, high or very high stringency (see Sambrook et al., *Molecular Cloning: A Laboratory Manual*, 2nd ed., Cold Spring Harbor Laboratory Press, New York, 1989); or (c) the DNA sequences are degenerate as a result of the genetic code to the DNA sequences defined in (a) or (b). Further, the nucleic acid molecule disclosed herein includes both complementary and non-complementary sequences, provided the sequences otherwise meet the criteria set forth herein.

add comma

secreted

MO

binding

Construct

cdi

Nutl.

Acad.

Stratagene

insert

Spring

fragments

31

Within the context of the present invention, high stringency means standard hybridization conditions (e.g., 5xSSPE, 0.5% SDS at 65° C., or the equivalent).

The structure of the proteins encoded by the nucleic acid molecules described herein may be predicted from the primary translation products using the hydrophobicity plot function of, for example, P/C Gene or Intelligentsias Suite (Intelligentsias, Mountain View, Calif.), or according to the methods described by Kyte and Doolittle (*J. Mol. Biol.* 157:105-132, 1982).

Proteins of the present invention may be prepared in the form of acidic or basic salts, or in neutral form. In addition, individual amino acid residues may be modified by oxidation or reduction. Furthermore, various substitutions, deletions, or additions may be made to the amino acid or nucleic acid sequences, the net effect of which is to retain or further enhance or decrease the biological activity of the mutant or wild-type protein. Moreover, due to degeneracy in the genetic code, for example, there may be considerable variation in nucleotide sequences encoding the same amino acid sequence.

Other derivatives of the proteins disclosed herein include conjugates of the proteins along with other proteins or polypeptides. This may be accomplished, for example, by the synthesis of N-terminal or C-terminal fusion proteins which may be added to facilitate purification or identification of proteins (see U.S. Pat. No. 4,851,341, see also, Hopp et al., *BioTechnology* 6:1204, 1988.) Alternatively, fusion proteins such as Flag/TGF-beta binding protein, be constructed in order to assist in the identification, expression, and analysis of the protein.

Proteins of the present invention may be constructed using a wide variety of techniques described herein. Further, mutations may be introduced at particular loci by synthesizing oligonucleotides containing a mutant sequence, flanked by restriction sites enabling ligation to fragments of the native sequence. Following ligation, the resulting reconstructed sequence encodes a derivative having the desired amino acid insertion, substitution, or deletion.

Alternately, oligonucleotide-directed site-specific (or segment specific) mutagenesis procedures may be employed to provide an altered gene having particular codons altered according to the substitution, deletion, or insertion required. Exemplary methods of making the alterations set forth above are disclosed by Walder et al. (*Gene* 42:133, 1986); Bauer et al. (*Gene* 37:73, 1985); Craik (*BioTechniques*, January 1985, 12-19); Smith et al. (*Genetic Engineering; Principles and Methods*, Plenum Press, 1981); and Sambrook et al. (supra). Deletion or truncation derivatives of proteins (e.g., a soluble extracellular portion) may also be constructed by utilizing convenient restriction endonuclease sites adjacent to the desired deletion. Subsequent to restriction, overhangs may be filled in, and the DNA religated. Exemplary methods of making the alterations set forth above are disclosed by Sambrook et al. (*Molecular Cloning: A Laboratory Manual*, 2d Ed., Cold Spring Harbor Laboratory Press, 1989).

Mutations which are made in the nucleic acid molecules of the present invention preferably preserve the reading frame of the coding sequences. Furthermore, the mutations will preferably not create complementary regions that could hybridize to produce secondary mRNA structures, such as loops or hairpins, that would adversely affect translation of the mRNA. Although a mutation site may be predetermined, it is not necessary that the nature of the mutation per se be predetermined. For example, in order to select for optimum characteristics of mutants at a given site, random mutagen-

32

esis may be conducted at the target codon and the expressed mutants screened for indicative biological activity. Alternatively, mutations may be introduced at particular loci by synthesizing oligonucleotides containing a mutant sequence, flanked by restriction sites enabling ligation to fragments of the native sequence. Following ligation, the resulting reconstructed sequence encodes a derivative having the desired amino acid insertion, substitution, or deletion.

Nucleic acid molecules which encode proteins of the present invention may also be constructed utilizing techniques of PCR mutagenesis, chemical mutagenesis (Drinkwater and Klinedinst, *PNAS* 83:3402-3406, 1986), by forced nucleotide misinformation (e.g., Liao and Wise *Gene* 88:107-111, 1990), or by use of randomly mutagenized oligonucleotides (Horwitz et al., *Genome* 3:112-117, 1989).

The present invention also provides for the manipulation and expression of the above described genes by culturing host cells containing a vector capable of expressing the above-described genes. Such vectors or vector constructs include either synthetic or cDNA-derived nucleic acid molecules encoding the desired protein, which are operably linked to suitable transcriptional or translational regulatory elements. Suitable regulatory elements may be derived from a variety of sources, including bacterial, fungal, viral, mammalian, insect, or plant genes. Selection of appropriate regulatory elements is dependent on the host cell chosen and may be readily accomplished by one of ordinary skill in the art. Examples of regulatory elements include: a transcriptional promoter and enhancer or RNA polymerase binding sequence, a transcriptional terminator, and a ribosomal binding sequence, including a translation initiation signal.

Nucleic acid molecules that encode any of the proteins described above may be readily expressed by a wide variety of prokaryotic and eukaryotic host cells, including bacterial, mammalian, yeast or other fungi, viral, insect, or plant cells. Methods for transforming or transfecting such cells to express foreign DNA are well known in the art (see, e.g., Itakura et al., U.S. Pat. No. 4,704,362; Hinnen et al., *Proc. Natl. Acad. Sci. USA* 75:1929-1933, 1978; Murray et al., U.S. Pat. No. 4,801,542; Upshall et al., U.S. Pat. Nos. 4,935,349; Hagen et al., U.S. Pat. No. 4,784,950; Axel et al., U.S. Pat. No. 4,399,216; Goeddel et al., U.S. Pat. No. 4,766,075; and Sambrook et al. *Molecular Cloning: A Laboratory Manual*, 2nd ed., Cold Spring Harbor Laboratory Press, 1989; for plant cells see Czako and Marton, *Plant Physiol.* 104:1067-1071, 1994; and Paszkowski et al., *Biotech* 24:387-392, 1992).

Bacterial host cells suitable for out the present invention include *E. coli*, *B. subtilis*, *Salmonella typhimurium*, and various species within the genera *Pseudomonas*, *Streptomyces*, and *Staphylococcus*, as well as many other bacterial species well known to one of ordinary skill in the art. Representative examples of bacterial host cells include DH5α (Stratagene, LaJolla, Calif.).

Bacterial expression vectors preferably comprise a promoter which functions in the host cell, one or more selectable phenotypic markers, and a bacterial-origin of replication. Representative promoters include the β-lactamase (penicillinase) and lactose promoter system (see Chang et al., *Nature* 275:615, 1978), the T7 RNA polymerase promoter (Studier et al., *Meth. Enzymol.* 185:60-89, 1990), the lambda promoter (Elvin et al., *Gene* 87:123-126, 1990), the trp promoter (Nichols and Yanofsky, *Meth. in Enzymology* 101:155, 1983) and the lac promoter (Russell et al., *Gene* 20:231, 1982). Representative selectable markers include various antibiotic resistance markers such as the kanamycin

may

add " "

create

capable

bacterial

add comma

"trp" in italics

transforming

phosphotransferase

or ampicillin resistance genes. Many plasmids suitable for transforming host cells are well known in the art, including among others, pBR322 (see Bolivar et al., *Gene* 2:95, 1977), the pUC plasmids pUC18, pUC19, pUC118, pUC119 (see Messing, *Meth. in Enzymology* 101:20-77, 1983 and Vieira and Messing, *Gene* 19:259-268, 1982), and pNH8A, pNH16a, pNH18a, and Bluescript M13 (Stratagene, La Jolla, Calif.).

Yeast and fungi host cells suitable for carrying out the present invention include, among others, *Saccharomyces pombe*, *Saccharomyces cerevisiae*, the genera *Pichia* or *Kluyveromyces* and various species of the genus *Aspergillus* (McKnight et al., U.S. Pat. No. 4,935,349). Suitable expression vectors for yeast and fungi include, among others, YCp50 (ATCC No. 37419) for yeast, and the amdS cloning vector pV3 (Trumbull, *Bio/Technology* 7:169, 1989), YRp7 (Struhl et al., *Proc. Natl. Acad. Sci. USA* 76:1035-1039, 1978), YEpl3 (Broach et al., *Gene* 8:121-133, 1979), pJDB249 and pJDB219 (Beggs, *Nature* 275:104-108, 1978) and derivatives thereof.

Preferred promoters for use in yeast include promoters from yeast glycolytic genes (Hitzeman et al., *J. Biol. Chem.* 255:12073-12080, 1980; Alber and Kawasaki, *J. Mol. Appl. Genet.* 1:419-434, 1982) or alcohol dehydrogenase genes (Young et al., in *Genetic Engineering of Microorganisms for Chemicals*, Hollaender et al. (eds.), p. 355, Plenum, New York, 1982; Ammerer, *Meth. Enzymol.* 101:192-201, 1983). Examples of useful promoters for fungi vectors include those derived from *Aspergillus nidulans* glycolytic genes, such as the adh3 promoter (McKnight et al., *EMBO J.* 4:2093-2099, 1985). The expression units may also include a transcriptional terminator. An example of a suitable terminator is the adh3 terminator (McKnight et al., *ibid.*, 1985).

As with bacterial vectors, the yeast vectors will generally include a selectable marker, which may be one of any number of genes that exhibit a dominant phenotype for which a phenotypic assay exists to enable transformants to be selected. Preferred selectable markers are those that complement host cell auxotrophy, provide antibiotic resistance or enable a cell to utilize specific carbon sources, and include *leu2* (Broach et al., *ibid.*), *ura3* (Bosch et al., *Gene* 8:17, 1979), or *his3* (Struhl et al., *ibid.*). Another suitable selectable marker is the *cap* gene, which confers chloramphenicol resistance on yeast cells.

Techniques for transforming fungi are well known in the literature, and have been described, for instance, by Beggs (*ibid.*), Hinnen et al. (*Proc. Natl. Acad. Sci. USA* 75:1929-1933, 1978), Yelton et al. (*Proc. Natl. Acad. Sci. USA* 81:1740-1747, 1984), and Russell (*Nature* 301:167-169, 1983). The genotype of the host cell may contain a genetic defect that is complemented by the selectable marker present on the expression vector. Choice of a particular host and selectable marker is well within the level of ordinary skill in the art.

Protocols for the transformation of yeast are also well known to those of ordinary skill in the art. For example, transformation may be readily accomplished either by preparation of spheroplasts of yeast with DNA (see Hinnen et al., *PNAS USA* 75:1929, 1978) or by treatment with alkaline salts such as LiCl (see Itoh et al., *J. Bacteriology* 153:163, 1983). Transformation of fungi may also be carried out using polyethylene glycol as described by Cullen et al. (*Bio/Technology* 5:369, 1987).

Viral vectors include those which comprise a promoter that directs the expression of an isolated nucleic acid molecule that encodes a desired protein as described above. A wide variety of promoters may be used within the context of

the present invention, including for example, promoters such as MoMLV LTR, RSV LTR, Friend MuLV LTR, adenoviral promoter (Ohno et al., *Science* 265:781-784, 1994), neomycin phosphotransferase promoter/enhancer, late parvovirus promoter (Koenig et al., *Hum. Gene Therap.* 5:457-463, 1994), Herpes TK promoter, SV40 promoter, metallothionein IIa gene enhancer/promoter, cytomegalovirus immediate early promoter, and the cytomegalovirus immediate late promoter. Within particularly preferred embodiments of the invention, the promoter is a tissue-specific promoter (see e.g., WO 91/02805; EP 0,415,731; and WO 90/07936). Representative examples of suitable tissue specific promoters include neural specific enolase promoter, platelet derived growth factor beta promoter, bone morphogenic protein promoter, human alpha-chimaerin promoter, synapsin I promoter and synapsin II promoter. In addition to the above-noted promoters, other viral-specific promoters (e.g., retroviral promoters (including those noted above, as well as others such as HIV promoters), hepatitis, herpes (e.g., EBV), and bacterial, fungal or parasitic (e.g., malarial) -specific promoters may be utilized in order to target a specific cell or tissue which is infected with a virus, bacteria, fungus or parasite.

Mammalian cells suitable for carrying out the present invention include, among others COS, CHO, SaOS, osteosarcomas, KS483, MG-63, primary osteoblasts, and human or murine bone marrow stroma. Mammalian expression vectors for use in carrying out the present invention will include a promoter capable of directing the transcription of a cloned gene or cDNA. Preferred promoters include viral promoters and cellular promoters. Bone specific promoters include the bone sialo-protein and the promoter for osteocalcin. Viral promoters include the cytomegalovirus immediate early promoter (Boshart et al., *Cell* 41:521-530, 1985), cytomegalovirus immediate late promoter, SV40 promoter (Subramani et al., *Mol. Cell Biol.* 1:854-864, 1981), MMTV LTR, RSV LTR, metallothionein-1, adenovirus Ela. Cellular promoters include the mouse metallothionein-1 promoter (Palmiter et al., U.S. Pat. No. 4,579,821), a mouse V_{κ} promoter (Bergman et al., *Proc. Natl. Acad. Sci. USA* 81:7041-7045, 1983; Grant et al., *Nucl. Acids Res.* 15:5496, 1987) and a mouse V_{H} promoter (Loh et al., *Cell* 33:85-93, 1983). The choice of promoter will depend, at least in part, upon the level of expression desired or the recipient cell line to be transfected.

Such expression vectors may also contain a set of RNA splice sites located downstream from the promoter and upstream from the DNA sequence encoding the peptide or protein of interest. Prefer RNA splice sites may be obtained from adenovirus and/or immunoglobulin genes. Also contained in the expression vectors is a polyadenylation signal located downstream of the coding sequence of interest. Suitable polyadenylation signals include the early or late polyadenylation signals from SV40 (Kaufman and Sharp, *ibid.*), the polyadenylation signal from the Adenovirus 5 E1B region and the human growth hormone gene terminator (DeNoto et al., *Nucl. Acids Res.* 9:3719-3730, 1981). The expression vectors may include a noncoding viral leader sequence, such as the Adenovirus 2 tripartite leader, located between the promoter and the RNA splice sites. Preferred vectors may also include enhancer sequences, such as the SV40 enhancer. Expression vectors may also include sequences encoding the adenovirus VA RNAs. Suitable expression vectors can be obtained from commercial sources (e.g., Stratagene, La Jolla, Calif.).

Vector constructs comprising cloned DNA sequences can be introduced into cultured mammalian cells by, for

Kluyveromyces

Turnbull

should be in italics

Sources should be in italics

should be in italics

Natl.

utilized

Botstein

virus

osteocalcin

Boshart

Biol

polyadenylation

transfection

US 6,803,453 B1

35

36

example, calcium phosphate-mediated transfecting (Wigler et al., *Cell* 14:725, 1978; Corsaro and Pearson, *Somatic Cell Genetics* 7:603, 1981; Gaharm and Van der Eb, *Virology* 52:456, 1973), electroporation (Neumann et al., *EMBO J.* 1:841-845, 1982), or DEAE-dextran mediated transfecting (Ausubel et al. (eds.), *Current Protocols in Molecular Biology*, John Wiley and Sons, Inc., New York, 1987). To identify cells that have stably integrated the cloned DNA, a selectable marker is generally introduced into the cells along with the gene or cDNA of interest. Preferred selectable markers for use in cultured mammalian cells include genes that confer resistance to drugs, such as neomycin, hygromycin, and methotrexate. The selectable marker may be an amplifiable selectable marker. Preferred amplifiable selectable markers are the DHFR gene and the neomycin resistance gene. Selectable markers are reviewed by Thilly (*Mammalian Cell Technology*, Butterworth Publishers, Stoneham, Mass., which is incorporated herein by reference).

Mammalian cells containing a suitable vector are allowed to grow for a period of time, typically 1-2 days, to begin expressing the DNA sequence(s) of interest. Drug selection is then applied to select for growth of cells that are expressing the selectable marker in a stable fashion. For cells that have been transfected with an amplifiable, selectable marker the drug concentration may be increased in a stepwise manner to select for increased copy number of the cloned sequences, thereby increasing expression levels. Cells expressing the introduced sequences are selected and screened for production of the protein of interest in the desired form or at the desired level. Cells that satisfy these criteria can then be cloned and scaled up for production.

Protocols for the transfecting of mammalian cells are well known to those of ordinary skill in the art. Representative methods include calcium phosphate mediated transfecting, electroporation, lipofection, retroviral, adenoviral and protoplast fusion-mediated transfecting (see Sambrook et al., supra). Naked vector constructs can also be taken up by muscular cells or other suitable cells subsequent to injection into the muscle of a mammal (or other animals).

Numerous insect host cells known in the art can also be useful within the present invention, in light of the subject specification. For example, the use of baculoviruses as vectors for expressing heterologous DNA sequences in insect cells has been reviewed by Atkinson et al. (*Pestic. Sci.* 28:215-224, 1990).

Numerous plant host cells known in the art can also be useful within the present invention, in light of the subject specification. For example, the use of *Agrobacterium rhizogenes* as vectors for expressing genes in plant cells has been reviewed by Simkar et al. (*J. Biosci. (Bangalore)* 11:47-58, 1987).

Within related aspects of the present invention, proteins of the present invention may be expressed in a transgenic animal whose germ cells and somatic cells contain a gene which encodes the desired protein and which is operably linked to a promoter effective for the expression of the gene. Alternatively, in a similar manner transgenic animals may be prepared that lack the desired gene (e.g., "knock-out" mice). Such transgenic may be prepared in a variety of non-human animals, including mice, rats, rabbits, sheep, dogs, goats and pigs (see Hammer et al., *Nature* 315:681-683, 1985; Palmer et al., *Science* 222:809-814, 1983; Brinster et al., *Proc. Natl. Acad. Sci. USA* 82:4438-442, 1985; Palmiter and Brinster, *Cell* 41:343-345, 1985, and U.S. Pat. Nos. 5,175,383 5,087,571, 4,736,866 5,387,742, 5,347,075, 5,221,778, and 5,175,384). Briefly, an expression vector, including a

nucleic acid molecule to be expressed together with appropriately positioned expression control sequences, is introduced into pronuclei of fertilized eggs, for example, by microinjection. Integration of the injected DNA is detected by blot analysis of DNA from tissue samples. It is preferred that the introduced DNA be incorporated into the germ line of the animal so that it is passed on to the animal's progeny. Tissue-specific expression may be achieved through the use of a tissue-specific promoter, or through the use of an inducible promoter, such as the metallothionein gene promoter (Palmiter et al., 1983, *ibid*), which allows regulated expression of the transgene.

Proteins can be isolated by, among other methods, culturing suitable host and vector systems to produce the recombinant translation products of the present invention. Supernatants from such cell lines, or protein inclusions or whole cells where the protein is not excreted into the supernatants, can then be treated by a variety of purification procedures in order to isolate the desired proteins. For example, the supernatants may be first concentrated using commercially available protein concentration filters, such as an Amicon or Millipore Pellicon ultrafiltration unit. Following concentration, the concentrate may be applied to a suitable purification matrix such as, for example, an anti-protein antibody bound to a suitable support. Alternatively, anion or cation exchange resins may be employed in order to purify the protein. As a further alternative, one or more reverse-phase high performance liquid chromatography (RP-HPLC) steps may be employed to further purify the protein. Other methods of isolating the proteins of the present invention are well known in the skill of the art.

A protein is deemed to be "isolated" within the context of the present invention if no other (undesired) protein is detected pursuant to SDS-PAGE analysis followed by Coomassie blue staining. Within other embodiments, the desired protein can be isolated such that no other (undesired) protein is detected pursuant to SDS-PAGE analysis followed by silver staining.

3. Nucleic Acid Molecules

Within other aspects of the invention, nucleic acid molecules are provided which are capable of inhibiting TGF-beta binding-protein binding to a member of the TGF-beta family. For example, within one embodiment antisense oligonucleotide molecules are provided which specifically inhibit expression of TGF-beta binding-protein nucleic acid sequences (see generally, Hirashima et al. in *Molecular Biology of RNA: New Perspectives* (M. Inouye and B. S. Dudock, eds.; 1987 Academic Press, San Diego, p. 401); *Oligonucleotides: Antisense Inhibitors of Gene Expression* (J. S. Cohen, ed., 1989 MacMillan Press, London); Stein and Cheng, *Science* 261:1004-1012, 1993; WO 95/10607; U.S. Pat. No. 5,359,051; WO 92/06693; and EP-A2-612844). Briefly, such molecules are constructed such that they are complementary to, and able to form Watson-Crick base pairs with, a region of transcribed TGF-beta binding-protein mRNA sequence. The resultant double-stranded nucleic acid interferes with subsequent processing of the mRNA, thereby preventing protein synthesis (see Example 10).

Within other aspects of the invention, ribozymes are provided which are capable of inhibiting the TGF-beta binding-protein binding to a member of the TGF-beta family. As used herein, "ribozymes" are intended to include RNA molecules that contain anti-sense sequences for specific recognition, and an RNA-cleaving enzymatic activity. The catalytic strand cleaves a specific site in a target RNA at greater than stoichiometric concentration. A wide variety of ribozymes may be utilized within the context of the present

inducible

Supernatant

Oligonucleotide

strand

mammalian
ordinary
transfection

Agrobacterium

transgenics

transgenic Palm transgenic

~~Haseloff~~ et al

US 6,803,453 B1

37

invention, including for example, the hammerhead ribozyme (for example, as described by Forster and Symons, *Cell* 48:211-220, 1987; Haseloff and Gerlach, *Nature* 328:596-600, 1988; Walbot and Bruening, *Nature* 334:196, 1988; Haseloff and Gerlach, *Nature* 334:585, 1988); the hairpin ribozyme (for example, as described by Haseloff et al., U.S. Pat. No. 5,254,678, issued Oct. 19, 1993 and Hempel et al., European Patent Publication No. 0 360,257, published Mar. 26, 1990); and *Tetrahymena ribosomal* RNA-based ribozymes (see Cech et al., U.S. Pat. No. 4,987,071). Ribozymes of the present invention typically consist of RNA, but may also be composed of DNA, nucleic acid analogs (e.g., phosphorothioates), or chimerics thereof (e.g., DNA/RNA/RNA).

4. Labels

The gene product or any of the candidate molecules described above and below, may be labeled with a variety of compounds, including for example, fluorescent molecules, toxins, and radionuclides. Representative examples of fluorescent molecules include fluorescein, Phycobilli proteins, such as phycoerythrin, rhodamine, Texas red and luciferase. Representative examples of toxins include ricin, abrin diphtheria toxin, cholera toxin, gelonin, pokeweed antiviral protein, tritin, Shigella toxin, and Pseudomonas exotoxin A. Representative examples of radionuclides include Cu-64, Ga-67, Ga-68, Zr-89, Ru-97, Tc-99m, Rh-105, Pd-109, Iu-111, I-123, I-125, I-131, Re-186, Re-188, Au-198, Au-199, Pb-203, At-211, Pb-212 and Bi-212. In addition, the antibodies described above may also be labeled or conjugated to one partner of a ligand binding pair. Representative examples include avidin-biotin, and riboflavin-riboflavin binding protein.

Methods for conjugating or labeling the molecules described herein with the representative labels set forth above may be readily accomplished by one of ordinary skill in the art (see Trichothecene Antibody Conjugate, U.S. Pat. No. 4,744,981; Antibody Conjugate, U.S. Pat. No. 5,106,951; Fluorogenic Materials and Labeling Techniques, U.S. Pat. No. 4,018,884; Metal Radionuclide Labeled Proteins for Diagnosis and Therapy, U.S. Pat. No. 4,897,255; and Metal Radionuclide Chelating Compounds for Improved Chelation Kinetics, U.S. Pat. No. 4,988,496; see also Inman *Methods In Enzymology*, Vol. 34, *Affinity Techniques, Enzyme Purification*: Part B, Jakoby and Wilchek (eds.), Academic Press, New York, p. 30, 1974; see also Wilchek and Bayer, "The Avidin-Biotin Complex in Bioanalytical Applications," *Anal. Biochem.* 171:1-32, 1988).

Pharmaceutical Compositions

As noted above, the present invention also provides a variety of pharmaceutical compositions, comprising one of the above-described molecules which inhibits the TGF-beta binding-protein binding to a member of the TGF-beta family along with a pharmaceutically or physiologically acceptable carrier, excipients or diluents. Generally, such carriers should be nontoxic to recipients at the dosages and concentrations employed. Ordinarily, the preparation of such compositions entails combining the therapeutic agent with buffers, antioxidants such as ascorbic acid, low molecular weight (less than about 10 residues) polypeptides, proteins, amino acids, carbohydrates including glucose, sucrose or dextrans, chelating agents such as EDTA, glutathione and other stabilizers and excipients. Neutral buffered saline or saline mixed with nonspecific serum albumin are exemplary appropriate diluents.

In addition, the pharmaceutical compositions of the present invention may be prepared for administration by a

38

variety of different routes. In addition, pharmaceutical compositions of the present invention may be placed within containers, along with packaging material which provides instructions regarding the use of such pharmaceutical compositions. Generally, such instructions will include a tangible expression describing the reagent concentration, as well as within certain embodiments, relative amounts of excipient ingredients or diluents (e.g., water, saline or PBS) which may be necessary to reconstitute the pharmaceutical composition.

Methods of Treatment

The present invention also provides methods for increasing the mineral content and mineral density of bone. Briefly, numerous conditions result in the loss of bone mineral content, including for example, disease, genetic predisposition, accidents which result in the lack of use of bone (e.g., due to fracture), therapeutics which effect bone resorption, or which kill bone forming cells and normal aging. Through use of the molecules described herein which inhibit the TGF-beta binding-protein binding to a TGF-beta family member such conditions may be treated or prevented. As utilized herein, it should be understood that bone mineral content has been increased, if bone mineral content has been increased in a statistically significant manner (e.g., greater than one-half standard deviation), at a selected site.

A wide variety of conditions which result in loss of bone mineral content may be treated with the molecules described herein. Patients with such conditions may be identified through clinical diagnosis utilizing well known techniques (see, e.g., Harrison's Principles of Internal Medicine, McGraw-Hill, Inc.). Representative examples of diseases that may be treated include dysplasias, wherein there is abnormal growth or development of bone. Representative examples of such conditions include achondroplasia, cleidocranial dysostosis, enchondromatosis, fibrous dysplasia, Gaucher's, hypophosphatemic rickets, Marfan's, multiple hereditary exotoses, neurofibromatosis, osteogenesis imperfecta, osteopetrosis, osteopoikilosis, sclerotic lesions, fractures, periodontal disease, pseudoarthrosis and pyogenic osteomyelitis.

Other conditions which may be treated or prevented include a wide variety of causes of osteopenia (i.e., a condition that causes greater than one standard deviation of bone mineral content or density below peak skeletal mineral content at youth). Representative examples of such conditions include anemic states, conditions caused by steroids, conditions caused by heparin, bone marrow disorders, scurvy, malnutrition, calcium deficiency, idiopathic osteoporosis, congenital osteopenia or osteoporosis, alcoholism, chronic liver disease, senility, postmenopausal state, oligomenorrhea, amenorrhea, pregnancy, diabetes mellitus, hyperthyroidism, Cushing's disease, acromegaly, hypogonadism, immobilization or disuse, reflex sympathetic dystrophy syndrome, transient regional osteoporosis and osteomalacia.

Within one aspect of the present invention, bone mineral content or density may be increased by administering to a warm-blooded animal a therapeutically effective amount of a molecule which inhibits the TGF-beta binding-protein binding to a TGF-beta family member. Examples of warm-blooded animals that may be treated include both vertebrates and mammals, including for example horses, cows, pigs, sheep, dogs, cats, rats and mice. Representative examples of therapeutic molecules include ribozymes, ribozyme genes, antisense oligonucleotides and antibodies (e.g., humanized antibodies).

should not be in italics

phycoerythrin

preparation

Briefly

treated

treated
no hyphen

diagnosis

osteogenesis
hereditary

by
Calcium
osteopenia

add "."

Within other aspects of the present invention, methods are provided for increasing bone density, comprising the step of introducing into cells which home to bone a vector which directs the expression of a molecule which inhibits the TGF-beta binding-protein binding to a member of the TGF-beta family, and administering the vector containing cells to a warm-blooded animal. Briefly, cells which home to bone may be obtained directly from the bone of patents (e.g. cells obtained from the bone marrow such as CD34+, osteoblasts, osteocytes, and the like), from periphery blood, or from cultures.

A vector which direct the expression of a molecule that inhibits the TGF-beta binding-protein binding to a member of the TGF-beta family is introduced into the cells. Representative examples of suitable vectors include viral vectors such as herpes viral vectors (e.g., U.S. Pat. No. 5,288,641), adenoviral vectors (e.g., WO 94/26914, WO 93/9191; Kolls et al., *PNAS* 91(1):215-219, 1994; Kass-Eisler et al., *PNAS* 90(24):11498-502, 1993; Guzman et al., *Circulation* 88(6):2838-48, 1993; Guzman et al., *Cir. Res.* 73(6):1202-1207, 1993; Zabner et al., *Cell* 75(2):207-216, 1993; Li et al., *Hum Gene Ther.* 4(4):403-409, 1993; Caillaud et al., *Eur. J. Neurosci.* 5(10):1287-1291, 1993; Vincent et al., *Nat. Genet.* 5(2):130-134, 1993; Jaffe et al., *Nat. Genet.* 1(5):372-378, 1992; and Levrero et al., *Gene* 101(2):195-202, 1991), adeno-associated viral vectors (WO 95/13365; Flotte et al., *PNAS* 90(22):10613-10617, 1993), baculovirus vectors, parvovirus vectors (Koering et al., *Hum. Gene Therap.* 5:457-463, 1994), pox virus vectors (Panicali and Paoletti, *PNAS* 79:4927-4931, 1982; and Ozaki et al., *Biochem. Biophys. Res. Comm.* 193(2):653460, 1993), and retroviruses (e.g., EP 0,415,731; WO 90/07936; WO 91/0285, WO 94/03622; WO 93/25698; WO 93/25234; U.S. Pat. No. 5,219,740; WO 93/11230; WO 93/10218). Viral vectors may likewise be constructed which contain a mixture of different elements (e.g., promoters, envelope sequences and the like) from different viruses, or non-viral sources. Within various embodiments, either the viral vector itself, or a viral particle which contains the viral vector may be utilized in the methods and compositions described below.

Within other embodiments of the invention, nucleic acid molecules which encode a molecule which inhibits the TGF-beta binding-protein binding to a member of the TGF-beta family themselves may be administered by a variety of techniques, including, for example, administration of asialoglycomucoid (ASOR) conjugated with poly-L-lysine DNA complexes (Crimino et al., *PNAS* 92:122-92126 1993), DNA linked to killed adenovirus (Curiel et al., *Hum. Gene Ther.* 3(2):147-154, 1992), cytofectin-mediated introduction (DMRIE-DOPE, Vical, Calif.), direct DNA injection (Acandi et al., *Nature* 352:815-818, 1991); DNA ligand (Wu et al., *J. of Biol. Chem.* 264:16985-16987, 1989); lipofection (Felgner et al., *Proc. Natl. Acad. Sci. USA* 84:7413-7417, 1999); liposomes (Picketing et al., *Circ.* 89(1):13-21, 1994; and Wang et al., *PNAS* 84:7851-7855, 1987); microprojectile bombardment (Williams et al., *PNAS* 88:2726-2730, 1991); and direct delivery of nucleic acids which encode the protein itself either alone (Vile and Hart, *Cancer Res.* 53:3860-3864, 1993), or utilizing PEG-nucleic acid complexes.

Representative examples of molecules which may be expressed by the vectors of present invention include ribozymes and antisense molecules, each of which are discussed in more detail above.

Determination of increased bone mineral content may be determined directly through the use of X-rays (e.g., Dual Energy X-ray Absorptometry or "DEXA"), or by inference through bone turnover markers (osteoblast specific alkaline phosphatase, osteocalcin, type 1 procollagen C' propeptide

(PICP), and total alkaline phosphatase; see Comier, C., *Curr. Opin. in Rheu.* 7:243, 1995), or markers of bone resorption (pyridinoline, deoxypyridinoline, N-telopeptide, urinary hydroxyproline, plasma tartrate-resistant acid phosphatase and galactosyl hydroxylysine; see Comier, supra). The amount of bone mass may also be calculated from body weights, or utilizing other methods (see Guiness-Hey, *Metab. Bone Dis. and Rel. Res.* 5:177-181, 1984).

As will be evident to one of skill in the art, the amount and frequency of administration will depend, of course, on such factors as the nature and severity of the indication being treated, the desired response, the condition of the patient, and so forth. Typically, the compositions may be administered by a variety of techniques, as noted above.

The following examples are offered by way of illustration, and not by way of limitation.

EXAMPLES

Example 1

Sclerosteosis Maps to the Long Arm of Human Chromosome 17

Genetic mapping of the defect responsible for sclerosteosis in humans, located the gene responsible for this disorder to the region of human chromosome 17 that encodes a novel TGF-beta binding-protein family member. In sclerosteosis skeletal bone displays a substantial increase in mineral density relative to that of unaffected individuals. Bone in the head displays overgrowth as well. Sclerosteosis patients are generally healthy although they may exhibit variable degrees of syndactyly at birth and variable degrees of cranial compression and nerve compression in the skull.

Linkage analysis of the gene defect associated with sclerosteosis was conducted by applying the homozygosity mapping method to DNA samples collected from 24 South African Afrikaner families in which the disease occurred. (Sheffield et al., 1994, *Human Molecular Genetics* 3:1331-1335. "Identification of a Bardet-Biedt syndrome locus on chromosome 3 and evaluation of an efficient approach to homozygosity mapping"). The Afrikaner population of South Africa is genetically homogeneous; the population is descended from a small number of founders who colonized the area several centuries ago, and it has been isolated by geographic and social barriers since the founding. Sclerosteosis is rare everywhere in the world outside the Afrikaner community, which suggests that a mutation in the gene was present in the founding population and has since increased in numbers along with the increase in the population. The use of homozygosity mapping is based on the assumption that DNA mapping markers adjacent to a recessive mutation are likely to be homozygous in affected individuals from consanguineous families and isolated populations.

A set of 371 microsatellite markers (Research Genetics, Set 6) from the autosomal chromosomes was selected to type pools of DNA from sclerosteosis patient samples. The DNA samples for this analysis came from 29 sclerosteosis patients in 24 families, 59 unaffected family members and a set of unrelated control individuals from the same population. The pools consisted of 4-6 individuals, either affected individuals, affected individuals from consanguineous families, parents and unaffected siblings, or unrelated controls. In the pools of unrelated individuals and in most of the pools with affected individuals or family members analysis of the markers showed several allele sizes for each marker.

One marker, D17S1299, showed an indication of homozygosity: one band in several of the pools of affected individuals.

All 24 sclerosteosis families were typed with a total of 19 markers in the region of D17S1299 (at 17q12-q21). Affected individuals from every family were shown to be homozygous in this region, and 25 of the 29 individuals were homozygous for a core haplotype; they each had the same alleles between D17S1787 and D17S930. The other four individuals had one chromosome which matched this haplotype and a second which did not. In sum, the data compellingly suggested that this 3 megabase region contained the sclerosteosis mutation. Sequence analysis of most of the exons in this 3 megabase region identified a nonsense mutation in the novel TGF-beta binding-protein coding sequence (C>T mutation at position 117 of Sequence ID No. 1 results in a stop codon). This mutation was shown to be unique to sclerosteosis patients and carriers of Afrikaner descent. The identity of the gene was further confirmed by identifying a mutation in its intron (A>T mutation at position +3 of the intron) which results in improper mRNA processing in a single, unrelated patient with diagnosed sclerosteosis.

Example 2

Tissue-specificity of TGF-beta Binding-protein Gene Expression by RT-PCR:

A. Human Beer Gene Expression by RT-PCR:

Fist-strand cDNA was prepared from the following total RNA samples using a commercially available kit ("Superscript Preamplification System for First-Strand cDNA Synthesis", Life Technologies, Rockville, Md.): human brain, human liver, human spleen, human thymus, human placenta, human skeletal muscle, human thyroid, human pituitary, human osteoblast (NHOst from Clonetics Corp., San Diego, Calif.), human osteosarcoma cell line (Saos-2, ATCC# HTB-85), human bone, human bone marrow, human cartilage, vervet monkey bone, saccharomyces cerevisiae, and human peripheral blood monocytes. All RNA samples were purchased from a commercial source (Clontech, Palo Alto, Calif.), except the following which were prepared in-house; human osteoblasts, human osteosarcoma cell line, human bone, human cartilage and vervet monkey bone. These in-house RNA samples were prepared using a commercially available kit ("TRI Reagent", Molecular Research Center, Inc., Cincinnati, Ohio).

PCR was performed on these samples, and additionally on a human genomic sample as a control. The sense Beer oligonucleotide primer had the sequence 5'-CCGGAGCTGGAGAACAACAAG-3' (SEQ ID NO:19). The antisense Beer oligonucleotide primer had the sequence 5'-GCACTGGCCGGAGCACACC-3' (SEQ ID NO:20). In addition, PCR was performed using primers for the human beta-actin gene, as a control. The sense beta-actin oligonucleotide primer had the sequence 5'-AGGCCAACCAGGAGGAGATGACC-3' (SEQ ID NO:21). The antisense beta-actin oligonucleotide primer had the sequence 5'-GAAGTCCAGGGCAGCTAGCA-3' (SEQ ID NO:22). PCR was performed using standard conditions in 25 ul reactions, with an annealing temperature of 61 degrees Celsius. Thirty-two cycles of PCR were performed with the Beer primers and twenty-four cycles were performed with the beta-actin primers.

Following amplification, 12 ul from each reaction were analyzed by agarose gel electrophoresis and ethidium bromide staining. See FIG. 2A.

B. RNA In-situ Hybridization of Mouse Embryo Sections:

The full length mouse Beer cDNA (Sequence ID No. 11) was cloned into the pCR2.1 vector (Invitrogen, Carlsbad, Calif.) in the antisense and sense direction using the manufacturer's protocol. ³⁵S-alpha-GTP-labeled cRNA sense and antisense transcripts were synthesized using in-vitro transcription reagents supplied by Ambion, Inc (Austin, Tex.) In-situ hybridization was performed according to the protocols of Lyons et al. (J. Cell Biol. 111:2427-2436, 1990).

The mouse Beer cRNA probe detected a specific message expressed in the neural tube, limb buds, blood vessels and ossifying cartilages of developing mouse embryos. Panel A in FIG. 3 shows expression in the apical ectodermal ridge (aer) of the limb (1) bud, blood vessels (bv) and the neural tube (nt). Panel B shows expression in the 4th ventricle of the brain (4). Panel C shows expression in the mandible (ma) cervical vertebrae (cv), occipital bone (oc), palate (pa) and a blood vessel (bv). Panel D shows expression in the ribs (r) and a heart valve (va). Panel A is a transverse section of 10.5 dpc embryo. Panel B is a sagittal section of 12.5 dpc embryo and panels C and D are sagittal sections of 15.5 dpc embryos. ba=branchial arch, b=heart, te=telencephalon (forebrain), b=brain, f=frontonasal mass, g=gut, h=heart, j=jaw, li=liver, lu=lung, ot=otic vesicle, ao=, sc=spinal cord, skm=skeletal muscle, ns=nasal sinus, th=thymus, to=tongue, fl=forelimb, di=diaphragm

Example 3

Expression and Purification of Recombinant Beer Protein

A. Expression in COS-1 Cells;

The DNA sequence encoding the full length human Beer protein was amplified using the following PCR oligonucleotide primers: The 5' oligonucleotide primer had the sequence 5'-AAGCTTGGTACCATGCAGC TCCCAC-3' (SEQ ID NO:23) and contained a HindIII restriction enzyme site (in bold) followed by 19 nucleotides of the Beer gene starting 6 base pairs prior to the presumed amino terminal start codon (ATG). The 3' oligonucleotide primer had the sequence 5'-AAGTTCTA CTGTGTCATCGTCGTCCTGTAGTCGTAGGCGTT CTC CAGCT-3' (SEQ ID NO:24) and contained a HindIII restriction enzyme site (in bold) followed by a reverse complement stop codon (CTA) followed by the reverse complement of the FLAG epitope (underlined, Sigma-Aldrich Co., St. Louis, Mo.) flanked by the reverse complement of nucleotides coding for the carboxy terminal 5 amino acids of the Beer. The PCR product was TA cloned ("Original TA Cloning Kit", Invitrogen, Carlsbad, Calif.) and individual clones were screened by DNA sequencing. A sequence-verified clone was then digested by HindIII and purified on a 1.5% agarose gel using a commercially available reagents ("QIAquick Gel Extraction Kit", Qiagen Inc., Valencia, Calif.). This fragment was then ligated to High digested phosphatase-treated pcDNA3.1 (Invitrogen, Carlsbad, Calif.) plasmid with T4 DNA ligase. DH10B E. coli were transformed and plated on LB, 100 µg/ml ampicillin plates. Colonies bearing the desired recombinant in the proper orientation were identified by a PCR-based screen, using a 5' primer corresponding to the T7 promoter/priming site in pcDNA3.1 and a 3' primer with the sequence 5'-GCACTGGCCGGAGCACACC-3' (SEQ ID NO:25) that corresponds to the reverse complement of internal BEER sequence. The sequence of the cloned fragment was confirmed by DNA sequencing.

COS-1 cells (ATCC# CRL-1650) were used for transfecting 50 µg of the expression plasmid pcDNA-Beer-Flag was transfected using a commercially available kit following

Standard 100

additionally
antisense
Seq ID
PCR
annealing

transcription
in italics
heart
insert
thymus

Purification
should be
in bold
epitope
amino
Hind III
all
delete

Transfection Kit
transfection
St.
protocols supplied by the manufacturer ("DEAE-Dextran Transfecting Kit") Sigma Chemical Co., St. Louis, Mo.). The final media following transfection was DMEM (Life Technologies, Rockville, Md.) containing 0.1% Fetal Bovine Serum. After 4 days in culture, the media was removed. Expression of recombinant BEER was analyzed by SDS-PAGE and Western Blot using anti-FLAG M2 monoclonal antibody (Sigma-Aldrich Co., St. Louis, Mo.). Purification of recombinant BEER protein was performed using an anti-FLAG M2 affinity column ("Mammalian Transient Expression System", Sigma-Aldrich Co., St. Louis, Mo.). The column profile was analyzed via SDS-PAGE and Western Blot using anti-FLAG M2 monoclonal antibody. B. Expression in SF9 Insect Cells:

The human Beer gene sequence was amplified using PCR with standard conditions and the following primers:

Sense primer: 5'-GTCGTCGGATCCATGGGGTGG CAGGCGTTCAAGAAATGAT-3' (SEQ ID NO:26)

Antisense primer: 5'-GTCGTCGAAGCTTCTACTTG TCATCGTCCTTGTAGTCGTA GCGTTTCACGCTCGGC-3' (SEQ ID NO:27)

epitope
BamHI
invitrogen
transfection
The resulting cDNA contained the coding region of Beer with two modifications. The N-terminal secretion signal was removed and a FLAG epitome tag (Sigma) was fused in frame to the C-terminal end of the insert. BamI and HindIII cloning sites were added and the gene was subcloned into pMelBac vector (Invitrogen) for transfer into a baculoviral expression vector using standard methods.

Recombinant baculoviruses expressing Beer protein were made using the Bac-N-Blue transfecting kit (Invitrogen) and purified according to the manufacturers instructions.

invitrogen
transfection
SF9 cells (Invitrogen) were maintained in TNM-FH media (Invitrogen) containing 10% fetal calf serum. For protein expression, SF9 cultures in spinner flasks were infected at an MOI of greater than 10. Samples of the media and cells were taken daily for five days, and Beer expression monitored by western blot using an anti-FLAG M2 monoclonal antibody (Sigma) or an anti-Beer rabbit polyclonal antiserum.

NaCl
elution
After five days the baculovirus-infected SF9 cells were harvested by centrifugation and cell associated protein was extracted from the cell pellet using a high salt extraction buffer (1.5 M NaCl, 50 mM Tris pH 7.5). The extract (20 ml per 300 ml culture) was clarified by centrifugation, dialyzed three times against four liters of Tris buffered saline (150 mM NaCl, 50 mM Tris pH 7.5), and clarified by centrifugation again. This high salt fraction was applied to Hitrap Heparin (Pharmacia; 5 ml bed volume), washed extensively with HEPES buffered saline (25 mM HEPES 7.5, 150 mM NaCl) and bound proteins were eluted with a gradient from 150 mM NaCl to 1200 mM NaCl. Beer elution was observed at approximately 800 mM NaCl. Beer containing fractions were supplemented to 10% glycerol and 1 mM DTT and frozen at -80 degrees C.

Example 4

Preparation and Testing of Polyclonal Antibodies to Beer, Gremlin, and Dan

A. Preparation of Antigen:

Human
The DNA sequences of Human Beer, Human Gremlin, and Human Dan were amplified using PCR methods with the following oligonucleotide primers:

H. Beer

Sense: 5'-GACTTGGATCCAGGGGTGGGCGTTC-3' (SEQ ID NO:28) Antisense 5'-AGCAAAAGCTT CAGTAGGCGTTCTCCAG-3' (SEQ ID NO:29)

H. Gremlin

Sense:

5'-GACTTGGATCCGAAGGGAAGAAAGAAAGGG-3' (SEQ ID NO:30) Antisense: 5'-AGCA TAAGCTTAATCCAAATCGATGGA-3' (SEQ ID NO:31)

H. Dan

Sense: 5'-ACTACGAGCTCGGCCCCACCACCCATC AACAAG-3' (SEQ ID NO:32) Antisense: 5'-ACTTAGAAGCTTTCAGTCCTCAGCCCCCT CTCC-3' (SEQ ID NO:33)

In each case the listed primers amplified the entire coding region minus the secretion signal sequence. These include restriction sites for subcloning into the bacterial expression vector pQE-30 (Qiagen Inc., Valencia, Calif.) at sites BamHI/HindIII for Beer and Gremlin, and sites SacI/HindIII for Dan. pQE30 contains a coding sequence for a 6xHis tag at the 5' end of the cloning region. The completed constructs were transformed into *E. coli* strain M-15/pRep (Qiagen Inc) and individual clones verified by sequencing. Protein expression in M-15/pRep and purification (6xHis affinity tag binding to Ni-NTA coupled to Sepharose) were performed as described by the manufacturer (Qiagen, The QIAexpressionist).

strain
solubilization
The *E. coli*-derived Beer protein was recovered in significant quantity using solubilization in 6M guanidine and dialyzed to 2-4M to prevent precipitation during storage. Gremlin and Dan protein were recovered in higher quantity with solubilization in 6M guanidine and a post purification guanidine concentration of 0.5M.

B. Production and Testing of Polyclonal Antibodies:

Polyclonal antibodies to each of the three antigens were produced in rabbit and in chicken hosts using standard protocols (R & R Antibody, Stanwood, Wash.; standard protocol for rabbit immunization and antisera recovery; Short Protocols in Molecular Biology, 2nd edition, 1992 11.37-11.41; Contributors Helen M. Cooper and Yvonne Paterson; chicken antisera was generated with Strac Biosolutions, Ramona, Calif.) Peterson

Contributors
generated
Rabbit antisera and chicken egg IgY fraction were screened for activity via Western blot. Each of the three antigens was separated by PAGE and transferred to 0.45 um nitrocellulose (Novex, San Diego, Calif.). The membrane was cut into strips with each strip containing approximately 75 ng of antigen. The strips were blocked in 3% Blotting Grade Block (Bio-Rad Laboratories, Hercules, Calif.) and washed 2 times in 1xTris buffer saline (TBS)/0.02% TWEEN buffer. The primary antibody (preimmunization bleeds, rabbit antisera or chick egg IgY in dilutions ranging from 1:100 to 1:10,000 in blocking buffer) was incubated with the strips for one hour with gentle rocking. A second series of three washes 1xTBS/0.02%TWEEN was followed by an one hour incubation with the secondary antibody (peroxidase conjugated donkey anti-rabbit, Amersham Life Science, Piscataway, N.J.; or peroxidase conjugated donkey anti-chicken, Jackson ImmunoResearch, West Grove, Pa.). A final cycle of 3x washes of 1xTBS/0.02%TWEEN was performed and the strips were developed with Lumi-Light Western Blotting Substrate (Roche Molecular Biochemicals, Mannheim, Germany).

C. Antibody Cross-reactivity Test:

Following the protocol described in the previous section, nitrocellulose strips of Beer, Gremlin or Dan were incubated with dilutions (1:5000 and 1:10,000) of their respective rabbit antisera or chicken egg IgY as well as to antisera or chicken egg IgY (dilutions 1:1000 and 1:5000) made to the remaining two antigens. The increased levels of nonmatch

ID CTAG AGG IgY

45

ing antibodies was performed to detect low affinity binding by those antibodies that may be seen only at increased concentration. The protocol and duration of development is the same for all three binding events using the protocol described above. There was no antigen cross-reactivity observed for any of the antigens tested.

Example 5

Interaction of Beer With TGF-beta Super-family Proteins

The interaction of Beer with proteins from different phylogenetic arms of the TGF- β superfamily were studied using immunoprecipitation methods. Purified TGF β -1, TGF β -2, TGF β -3, BMP-4, BMP-5, BMP-6 and GDNF were obtained from commercial sources (R&D systems; Minneapolis, Minn.). A representative protocol is as follows. Partially purified Beer was dialyzed into HEPES buffered saline (25 mM HEPES 7.5, 150 mM NaCl). Immunoprecipitations were done in 300 μ l of IP buffer (150 mM NaCl, 25 mM Tris pH 7.5, 1 mM EDTA, 1.4 mM β -mercaptoethanol, 0.5% Triton \times 100, and 10% glycerol). 30 ng recombinant human BMP-5 protein (R&D systems) was applied to 15 μ l of FLAG affinity matrix (Sigma; St Louis Mo.). In the presence and absence of 500 ng FLAG epitope-tagged Beer. The proteins were incubated for 4 hours @ 4 $^{\circ}$ C. and then the affinity matrix-associated proteins were washed 5 times in IP buffer (1 ml per wash). The bound proteins were eluted from the affinity matrix in 60 microliters of 1 \times SDS PAGE sample buffer. The proteins were resolved by SDS PAGE and Beer associated BMP-5 was detected by western blot using anti-BMP-5 antiserum (Research Diagnostics, Inc) (see FIG. 5). BEER Ligand Binding Assay:

FLAG-Beer protein (20 ng) is added to 100 μ l PBS/0.2% BSA and adsorbed into each well of 96 well microtiter plate previously coated with anti-FLAG monoclonal antibody (Sigma; St Louis Mo.) and blocked with 10% BSA in PBS. This is conducted at room temperature for 60 minutes. This protein solution is removed and the wells are washed to remove unbound protein. BMP-5 is added to each well in concentrations ranging from 10 pM to 500 nM in PBS/0.2% BSA and incubated for 2 hours at room temperature. The binding solution is removed and the plate washed with three times with 200 μ l volumes of PBS/0.2% BSA. BMP-5 levels are then detected using BMP-5 anti-serum via ELISA (F. M. Ausubel et al (1998) Current Protocols in Mol Biol. Vol 2 11.2.1-11.2.22). Specific binding is calculated by subtracting non-specific binding from total binding and analyzed by the LIGAND program (Munson and Podbard, Anal. Biochem, 107, p220-239, (1980)).

In a variation of this method, Beer is engineered and expressed as a human Fc fusion protein. Likewise the ligand BMP is engineered and expressed as mouse Fc fusion. These proteins are incubated together and the assay conducted as described by Mellor et al using homogeneous time resolved fluorescence detection (G. W. Mellor et al., J of Biomol Screening, 3(2) 91-99, 1998).

Example 6

Screening Assay for Inhibition of TGF-beta Binding-protein Binding to TGF-beta Family Members

The assay described above is replicated with two exceptions. First, BMP concentration is held fixed at the Kd determined previously. Second, a collection of antagonist candidates is added at a fixed concentration (20 μ M in the

46

case of the small organic molecule collections and 1 μ M in antibody studies). These candidate molecules (antagonists) of TGF-beta binding-protein binding include organic compounds derived from commercial or internal collections representing diverse chemical structures. These compounds are prepared as stock solutions in DMSO and are added to assay wells at \leq 1% of final volume under the standard assay conditions. These are incubated for 2 hours at room temperature with the BMP and Beer, the solution removed and the bound BMP is quantitated as described. Agents that inhibit 40% of the BMP binding observed in the absence of compound or antibody are considered antagonists of this interaction. These are further evaluated as potential inhibitors based on titration studies to determine their inhibition constants and their influence on TGF-beta binding-protein binding affinity. Comparable specificity control assays may also be conducted to establish the selectivity profile for the identified antagonist through studies using assays dependent on the BMP ligand action (e.g. BMP/BMP receptor competition study).

Example 7

Inhibition of TGF-beta Binding-protein Localization to Bone Matrix

Evaluation of inhibition of localization to bone matrix (hydroxyapatite) is conducted using modifications to the method of Nicolas (Nicolas, J Cell Tissue Int 57:206, 1995). Briefly, 125 I-labelled TGF-beta binding-protein is prepared as described by Nicolas (supra). Hydroxyapatite is added to each well of a 96-well microtiter plate equipped with a polypropylene filtration membrane (Polyfiltroninc, Weymouth Mass.). TGF-beta binding-protein is added to 0.2% albumin in PBS buffer. The wells containing matrix are washed 3 times with this buffer. Adsorbed TGF-beta binding-protein is eluted using 0.3M NaOH and quantitated.

Inhibitor identification is conducted via inactivation of TGF-beta binding-protein with test molecules and applying the mixture to the matrix as described above. The matrix is washed 3 times with 0.2% albumin in PBS buffer. Adsorbed TGF-beta binding-protein is eluted using 0.3 M NaOH and quantified. Agents that inhibit 40% of the TGF-beta binding-protein binding observed in the absence of compound or antibody are considered bone localization inhibitors. These inhibitors are further characterized through dose response studies to determine their inhibition constants and their influence on TGF-beta binding-protein binding affinity.

Example 8

Construction of TGF-beta Binding-protein Mutant A Mutagenesis:

A full-length TGF-beta binding-protein cDNA in pBlue-script SK serves as a template for mutagenesis. Briefly, appropriate primers (see the discussion provided above) are utilized to generate the DNA fragment by polymerase chain reaction using Vent DNA polymerase (New England Biolabs, Beverly, Mass.). The polymerase chain reaction is run for 23 cycles in buffers provided by the manufacturer using a 57 $^{\circ}$ C. annealing temperature. The product is then exposed to two restriction enzymes and after isolation using agarose gel electrophoresis, ligated back into pRBP4-503 from which the matching sequence has been removed by enzymatic digestion. Integrity of the mutant is verified by DNA sequencing.

B. Mammalian Cell Expression and Isolation of Mutant TGF-beta Binding-protein:

The mutant TGF-beta binding-protein cDNAs are transferred into the pcDNA3.1 mammalian expression vector described in EXAMPLE 3. After verifying the sequence, the resultant construct are transfected into COS-1 cells, and secreted protein is purified as described in EXAMPLE 3.

Example 9

Animal Models-I

Generation of Transgenic Mice Overexpressing the Beer Gene

The ~200 kilobase (kb) BAC clone 15G5, isolated from the CITB mouse genomic DNA library (distributed by Research Genetics, Huntsville, Ala.) was used to determine the complete sequence of the mouse Beer gene and its 5' and 3' flanking regions. A 41 kb Sall fragment, containing the entire gene body, plus ~17 kb of 5' flanking and ~20 kb of 3' flanking sequence was sub-cloned into the BamHI site of the SuperCos1 cosmid vector (Stratagene, La Jolla, Calif.) and propagated in the *E. coli* strain DH10B. From this cosmid construct, a 35 kb MluI-AvIII restriction fragment (Sequence No. 6), including the entire mouse Beer gene, as well as 17 kb and 14 kb of 5' and 3' flanking sequence, respectively, was then gel purified, using conventional means, and used for microinjection of mouse zygotes (DNX Transgress, U.S. Pat. No. 4,873,191). Founder animals in which the cloned DNA fragment was integrated randomly into the genome were obtained at a frequency of 5-30% of live-born pups. The presence of the transgene was ascertained by performing Southern blot analysis of genomic DNA extracted from a small amount of mouse tissue, such as the tip of a tail. DNA was extracted using the following protocol: tissue was digested overnight at 55° C. in a lysis buffer containing 200 mM NaCl, 100 mM Tris pH8.5, 5 mM EDTA, 0.2% SDS and 0.5 mg/ml Proteinase K. The following day, the DNA was extracted once with phenol/chloroform (50:50), once with chloroform/isoamylalcohol (24:1) and precipitated with ethanol. Upon resuspension in TE (10 mM Tris pH7.5, 1 mM EDTA) 8-10 ug of each DNA sample were digested with a restriction endonuclease, such as EcoRI, subjected to gel electrophoresis and transferred to a charged nylon membrane, such as HyBondN+ (Amersham Arlington Heights, Ill.). The resulting filter was then hybridized with a radioactively labelled fragment of DNA deriving from the mouse Beer gene locus, and able to recognize both a fragment from the endogenous gene locus and a fragment of a different size deriving from the transgene. Founder animals were bred to normal non-transgenic mice to generate sufficient numbers of transgenic and non-transgenic progeny in which to determine the effects of Beer gene overexpression. For these studies, animals at various ages (for example, 1 day, 3 weeks, 6 weeks, 4 months) are subjected to a number of different assays designed to ascertain gross skeletal formation, bone mineral density, bone mineral content, osteoclast and osteoblast activity, extent of endochondral ossification, cartilage formation, etc. The transcriptional activity from the transgene may be determined by extracting RNA from various tissues, and using an RT-PCR assay which takes advantage of single nucleotide polymorphisms between the mouse strain from which the transgene is derived (129Sv/J) and the strain of mice used for DNA microinjection [(C57BL/6J x SJL/J)F2].

Animal Models-II

Disruption of the Mouse Beer Gene by Homologous Recombination

Homologous recombination in embryonic stem (ES) cells can be used to inactivate the endogenous mouse Beer gene

and subsequently generate animals carrying the loss-of-function mutation. A reporter gene, such as the *E. coli* β -galactosidase gene, was engineered into the targeting vector so that its expression is controlled by the endogenous Beer gene's promoter and translational initiation signal. In this way, the spatial and temporal patterns of Beer gene expression can be determined in animals carrying a targeted allele.

The targeting vector was constructed by first cloning the drug-selectable phosphoglycerate kinase (PGK) promoter driven neomycin-resistance gene (neo) cassette from pGT-N29 (New England Biolabs, Beverly, Mass.) into the cloning vector pSP72 (Promega-Madison, Wis.). PCR was used to flank the PGKneo cassette with bacteriophage P1 loxP sites, which are recognition sites for the P1 Cre recombinase (Hoess et al., PNAS USA, 79:3398, 1982). This allows subsequent removal of the neo-resistance marker in targeted ES cells or ES cell-derived animals (U.S. Pat. No. 4,959, 317). The PCR primers were comprised of the 34 nucleotide (ntd) loxP sequence, 15-25 ntd complementary to the 5' and 3' ends of the PGKneo cassette, as well as restriction enzyme recognition sites (BamHI in the sense primer and EcoRI in the anti-sense primer) for cloning into pSP72. The sequence of the sense primer was 5'-AATCTGGATCCATACTTCGTATAGCATACATTATACGAAGTTATCTGCAGGATTCGAGGGGCCCT-3' (SEQ ID NO:34); sequence of the anti-sense primer was 5'-AATCTGAATTCCACC GTGTTAATTAAATAACTTCGTATAATGTATGCTATACGAAGTTATAGATCTAGAGTCAGCTTCTGA-3' (SEQ ID NO:35).

The next step was to clone a 3.6 kb XhoI-HindIII fragment, containing the *E. coli* β -galactosidase gene and SV40 polyadenylation signal from pSV β (Clontech, Palo Alto, Calif.) into the pSP72-PGKneo plasmid. The "short arm" of homology from the mouse Beer gene locus was generated by amplifying a 2.4 kb fragment from the BAC clone 15G5. The 3' end of the fragment coincided with the translational initiation site of the Beer gene, and the anti-sense primer used in the PCR also included 30 ntd complementary to the 5' end of the β -galactosidase gene so that its coding region could be fused to the Beer initiation site in-frame. The approach taken for introducing the "short arm" into the pSP72- β gal-PGKneo plasmid was to linearize the plasmid at a site upstream of the β -gal gene and then to co-transform this fragment with the "short arm" PCR product and to select for plasmids in which the PCR product was integrated by homologous recombination. The sense primer for the "short arm" amplification included 30 ntd complementary to the pSP72 vector to allow for this recombination event. The sequence of the sense primer was 5'-ATAGGTGACACT ATAGAACTCGAGCAGCTGAAGCTTAAACCACATGGTGGCTCACAACCAT-3' (SEQ ID NO:36) and the sequence of the anti-sense primer was 5'-AACGACGGCCAGTGAATCCGTAATCATGGTCATGCTGCCAGGTGGAGGAGGGCA-3' (SEQ ID NO:37).

The "long arm" from the Beer gene locus was generated by amplifying a 6.1 kb fragment from BAC clone 15G5 with primers which also introduce the rare-cutting restriction enzyme sites SgrAI, FseI, AscI and PacI. Specifically, the sequence of the sense primer was 5'-ATTACCACCGGTGACACCCGCTTCCTGACAG-3' (SEQ ID NO:38); the sequence of the anti-sense primer was 5'-ATTACTTAATTAAACATCGCGCGCCCATATGGCCGGCCCCCTAATTGCGGCCATCGTAATT-3' (SEQ ID NO:39). The resulting PCR product was cloned into the TA vector (Invitrogen, Carlsbad, Calif.) as an intermediate step.

should be in *Italis*

Madison

the sidase

should be ATT (not A)

ATG (not ATC)

delete "T" (not "I")

Transgenic

should be in *Italis*

Transgenic

should be in *Italis*

HyBondN+ transgenic

should be in *Italis*

osteoclast

transgenic strain

transgene

extracting

Sequence ID no. 1: human BEER cDNA (complete coding region plus 5' and 3' UTRs)

AGAGCCTGTGCTACTGGAAGGTGGCGTGCCTCTCTCGCTGGTACCATGCAGCTCCCACTGGCCCTGTGTCCTCGTCTGC

CTGCTGGTACACACAGCCTTCCGTGTAGTGGAGGGCCAGGGGTGGCAGGCGTTCAAGAATGATGCCACGAAATCATCCCC

CGAGCTCGGAGAGTACCCCGAGCCTCCACCGAGAGCTGGAGAACAACAAGACCATGAACCGGGCGGAGAACGGAGGGCGGC

CTCCCCACACCCCTTTGAGACCAAAGACGTGTCCGAGTACAGCTGCCGCGAGCTGCACCTTACC CGCTACGTGACCGAT

GGGCGCTGCGCGCAGCGCCAAGCCGGTCAACCAGCTGTGTGTGCTCCGGCCAGTGC GGCCCGGGCGCGCTGTGTC AACGC

CATCGGCCCGGGCAAGTGTGGCGACCTAGTGGGCCGACTTCCGCTGCATCCCCGACCCTACCGCGCGCAGCGCGTGC

AGTGTCTGTGTCCCGTGGTGTAGGCGCCGCGCGCGCAAGGTGCGCTGTGTGGCCTGTGCAAGTGAAGCGCCTCACC

CGCTTCCACAACCAGTCGGAGCTCAAGGACTTCGGGACCAGGCGCTCGGCCGCAAGAAGGGCGGAAGCCGCGGCCCG

CGCCCGGAGCGCCAAAGCCAACAGGCCGAGCTGGAGAAGCCCTACTAGAGCCCGCCCGCGCCCTCCCCACCGGGCGGC

GCCCCGGCCCTGAACCCGCGCCCCACATTTCTGTCTCTGCGCGTGGTTTGATTGTTTATATTTCATGTGTAATGCTCG

AACCCAGGGCAGGGGGCTGAGACCTTCAGGCCCTGAGGAATCCGGGCGCGCGCAAGGCCCCCTCAGCCCGCCAGCTG

AGGGGTCCACAGGGGCAGGGGAGGGAATTGAGAGTCAAGACACTGAGCCACGCGAGCCCGCCTCTGGGGCGCCCTACTC

TTGCTGGTCCCACTTCAGAGGAGGCAGAAATGGAAGCATTTTCACCGCCCTGGGGTTTTAAGGGAGCGGTGTGGGAGTGG

GAAAGTCCAGGGACTGGTTAAGAAAGTTGGATAAGATTCCCCCTGCACTCTGCTGCCCATCAGAAAGCCTGAGGCGTGC

CCAGAGCACAAGACTGGGGGCAACTGTAGATGTGGTTTCTAGTCTGGCTCTGCCACTAACTTGTGTGTAACCTTGAAAC

TACACAATTCCTCTCGGGACCTCAATTTCCACTTTGTAAAAATGAGGGTGGAGGTGGGAATAGGATCTCGAGGAGACTAT

TGGCATATGATTCGAAGGACTCCAGTGCTTTTGAAATGGGCAGAGGTGAGAGAGAGAGAGAGAAAGAGAGAGAATGAATG

CAGTTGCATTGATTCACTGTCGAAGGTCACTTCCAGAATTCAGAGTTGTGATGCTCTCTTCGACAGCCAAAGATGAAAAA

CAACAGAAAAAAGAAAGTAAAGAGTCTATTATGGCTGACATATTTACGGCTGACAAACTCTGGGAAGAGCTATGCTG

-continued

CTTCCAGCCTGGCTTCCCGGATGTTTGGCTACCTCCACCCCTCCATCTCAAAGAAATAACATCATCCATTGGGGTAGA
AAAGGAGAGGGTCCGAGGGTGGTGGGAGGATAGAAATCACATCCGCCCAACTTCCCAAAGAGCAGCATCCCTCCCCG
ACCCATAGCCATGTTTAAAGTCACCTCCGAAGAGAAGTGAAGGTCAAGGACACTGGCCTTGCAGGCCGAGGGAGC
AGCCATCACAACCTCAGACACGACATCCCTTTGAGACACCGCCTTCTGCCACCACCTCAGGACACATTTCTGCCT
AGAAAACAGCTTCTTACTGCTCTTACATGTGATGGCATATCTTACACTAAAAGAAATATTATTGGGGGAAAACTACAAGT
GCTGTACATATGCTGAGAACTGCAGAGCATAATAGCTGCCACCCAAAAATCTTTTGAATAATCTTCCAGACAACCTC
TTACTTTCTGTGTAGTTTAAATTGTTAAAAAAGTTTAAACAGAAGCACATGACATATGAAAGCCTGCAGGACT
GGTCGTTTTTTTGGCAATTCTCCACGTGGGACTTGTCCACAAGATGAAAGTAGTGGTTTTAAAGAGTTAAGTTACAT
ATTTATTTCTCACTTAAGTTATTTATGCAAAAGTTTCTTGTAGAGAATGACAATGTTAATATTGCTTTATGAATAA
CAGTCTGTTCTTCCAGAGTCCAGAGACATTGTTAATAAGACAATGAATCATGACCGAAAG

Sequence ID No. 2: Human BEER protein (complete sequence)

MLPLALCLVLLVHTAPRVVEGGWQAPKNDATETIPELGEYPEPPPELENNKTMNRAEMGGRPHHPFETKDVSEYSC
RELHPTRYVTDGPCRSAPVTELVCSGGQCGPARLLPNAIGRGKWRPSPGDFRCIPDRYRAQVRQLLCPGGEAPRARKVR
LVASCKRLTRFHNQSELKDFGTEAARPQKGRKPRPRARSAKANQAELENAY

Sequence ID No. 3: Human Beer cDNA containing Sclerosteosis nonsense mutation

AGAGCCTGTGCTACTGGAAGGTGGCGTCCCTCTCTGGCTGGTACCATGCAGCTCCCACTGGCCCTGTGTCTCGTCTGC
CTGCTGGTACACACAGCCTTCCTGTAGTGGAGGGCTAGGGGTGGCAGGCGTTCAAGAATGATGCCACGGAAATCATCC
CGAGCTCGGAGAGTACCCCGAGCTCCACCGAGCTGGAGAACAAGACCATGAACCGGGCGGAGAACGGAGGGCGGC
CTCCCCACACCCCTTTGAGACCAAGACGTGTCCGAGTACAGCTGCCGCGAGCTGCACTTACCCGCTACGTGACCGAT
GGCCCGTGGCGCAGCGCAAGCCGGTCAACCGAGCTGGTGTGCTCCGGCCAGTGGCGCCGCGCGCTGCTGCCAACGC
CATCGGCGCGGCAAGTGGTGGCGACCTAGTGGGCGCGACTTCCTGCTGCATCCCGACCGCTACCGCGCGCAGCGGTGC
AGCTGCTGTGCTCCGGTGGTGAAGCGCCGCGCGCGCAAGGTGCGCCTGGTGGCTCGTGCAGTGAAGCGCCTCACC
CGCTTCCACAACAGTGGAGCTCAAGGACTTCGGGACCGAGGCGCTCGGCGCAGAAGGGCCGAAGCCGCGGCCCG
CGCCCGGAGCGCAAGGCCAACAGCCGAGCTGGAGAAGCGCTACTAGAGCCCGCCGCGCCCTCCCCACCGCGGGC
GCCCCGGCCCTGAACCCGCGCCCAATTTCTGCTCTGCGCGTGGTTGATTGTTTATATTTTCAATTGTAAATGCCGTC
AACCAGGGCAGGGGGCTGAGACCTTCAGGCCCTGAGGAATCCCGGCGCGCGCAAGGCCCCCTCAGCCCGCCAGCTG
AGGGGTCCACGGGGCAGGGGAGGGAATTGAGAGTCACAGACACTGAGCCACGCGACCCCGCTCTGGGGCCGCTACCT
TTGCTGGTCCCACTTCAGAGGAGGAGCAAAATGAAGCAATTTTACCGCCCTGGGGTTTAAAGGAGCGGTGTGGAGTGG
GAAAGTCCAGGACTGGTTAAGAAAGTTGGATAAGATTCCCTTGCACCTCGCTGCCATCAGAAAGCTGAGGCGTGC
CCAGAGCACAAGACTGGGGCAACTGTAGATGTGGTTTCTAGTCTGGCTCTGCCACTAATCTGTGTGTAACCTTGAAC
TACACAATTCCTTCGGGACCTCAATTTCCACTTTGTAAATGAGGGTGGAGGTGGGAATAGGATCTCGAGGAGACTAT
TGGCATATGATTCCAAGGACTCCAGTGCCTTTTGAATGGGCAGAGGTGAGAGAGAGAGAGAAAGAGAGAAATGAATG
CAGTTGCATTGATTAGTCCAAAGGTCACTTCCAGAAATTCAGAGTTGTGATGCTCTCTCTGACAGCCAAAGATGAAAA
CAACAGAAAAAAGTAAAGAGTCTATTATGGCTGACATATTACGGCTGACAACTCTGGAAGAAGCTATGCTG
CTTCCAGCCTGGCTTCCCGGATGTTTGGCTACCTCCACCCCTCCATCTCAAAGAAATAACATCATCCATTGGGGTAGA
AAAGGAGAGGGTCCGAGGGTGGTGGGAGGATAGAAATCACATCCGCCCAACTTCCCAAAGAGCAGCATCCCTCCCCG
ACCCATAGCCATGTTTAAAGTCACCTCCGAAGAGAAGTGAAGGTCAAGGACACTGGCCTTGCAGGCCGAGGGAGC
AGCCATCACAACCTCAGACACGACATCCCTTTGAGACACCGCCTTCTGCCACCACCTCAGGACACATTTCTGCCT
AGAAAACAGCTTCTTACTGCTCTTACATGTGATGGCATATCTTACACTAAAAGAAATATTATTGGGGGAAAACTACAAGT
GCTGTACATATGCTGAGAACTGCAGAGCATAATAGCTGCCACCCAAAAATCTTTTGAATAATCTTCCAGACAACCTC
TTACTTTCTGTGTAGTTTAAATTGTTAAAAAAGTTTAAACAGAAGCACATGACATATGAAAGCCTGCAGGACT

-continued

GGTCGTTTTTTGGCAATTCTTCCACGTGGGACTTGTCACAAGAATGAAAGTAGTGGTTTTTAAAGAGTTAAGTTACAT
ATTATTTTCTCACTTAAGTTATTTATGCAAAAGTTTTCTTGTAGAGAATGACAATGTTAATATTGCTTTATGAATTAA
CAGTCTGTTCTTCCAGAGTCCAGAGACATTGTTAATAAAGACAATGAATCATGACCGAAAG

Sequence ID No. 4: Truncated Human Beer protein from Sclerosteosis
MQLPLALCLVCLLVHTAPRVVEG*

Sequence ID No. 5: Human BEER cDNA encoding protein variant (V10X)
AGAGCCTGTGCTACTGGAAGGTGGCTGCCCTCCTCTGGCTGGTACCATGCAGCTCCACTGGCCCTGTGTCTCATCTGC
CTGCTGGTACACACAGCCTTCCGTGTAGTGGAGGGCCAGGGGTGGCAGGCGTTCAAGAATGATGCCACGGAAATCATCCG
CGAGCTCGGAGAGTACCCCGAGCTCCACCGGAGCTGGAGAACACAAGACCATGAACCGGGCGGAGAACGGAGGGCGGC
CTCCCCACACCCCTTTGAGACCAAAGACGTGTCCGAGTACAGCTGCCGCGAGCTGCATTACCCGCTACGTGACCGAT
GGGCGGTGCCGACGCGCAAGCCGGTACCGAGCTGGTGTGCTCCGGCCAGTGGCGCCGGCGCGCTGTGCTGCCAACGC
CATCGGCCCGCGCAAGTGGTGGCGACCTAGTGGGCGGACTTCCGCTGCATCCCGACCGCTACCGCGCGCAGCGCGTGC
AGCTGCTGTGTCCTGGTGGTGGCGCCGCGCGCGCAAGGTGCCCTGGTGGCTCTGTGCAAGTGAAGCGCCTCACC
CGCTTCCACAACCACTCGGAGCTCAAGGACTTCGGGACCGAGGCGCTCGGCCGCGAAGGGCCGGAAGCCGCGGCCCG
CGCCCCGAGCGCCAAAGCCAAACAGCGGAGCTGGAGAAGCCTACTAGAGCCCGCCGCGCCCTCCCCACCGCGGGC
GCCCCGGCCCTGAACCCGCGCCCACTTTCTGTCTCTGCGCGTGGTTTGATTGTTTATATTTCATTGTAATGCTGTC
AACCAGGGCAGGGGGCTGAGACCTTCCAGGCCCTGAGGAATCCCGGGCGCCGGCAAGGCCCCCTCAGCCCGCCAGCTG
AGGGGTCCACGGGGCAGGGGAGGGAATGAGAGTCAAGACACTGAGCCACGCGAGCCCGCCTCTGGGGCGGCCTACCT
TTGCTGGTCCCACCTTCAGAGGAGCGAGAAATGAAGCATTTTCACGCCCTGGGGTTTTAAGGGAGCGGTGTGGAGTGG
GAAAGTCCAGGACTGGTTAAGAAAGTTGGATAAGATTCCTTGCACCTCGCTGCCCATCAGAAAGCCTGAGGCGTGC
CCAGAGCACAAGACTGGGGCAACTGTAGATGTGGTTTCTAGTCCCTGGCTCTGCCACTAATTGCTGTGTAACCTTGAAC
TACACAATTCCTTCCGGGACCTCAATTTCACCTTTGTAATGAGGGTGGAGGTGGGAATAGGATCTCGAGGAGACTAT
TGGCATATGATTCAAGGACTCCAGTGCCCTTTGTAATGGGAGAGGTGAGAGAGAGAGAGAAAGAGAGAGAAATGAATG
CAGTTGCATTGATTAGTGCAGGTCACCTCCAGAATTCAGAGTTGTGATGCTCTTCTTGACAGCCAAAGATGAAAAA
CAAACAGAAAAAAGTAAAGAGTCTATTTATGGCTGACATATTTACGGCTGACAACTCCTGGAAGAAGCTATGCTG
CTTCCAGCCTGGCTTCCCGGATGTTGGCTACCTCCACCCCTCCATCTCAAGAAATAACATCATTCATTGGGGTAGA
AAAGGAGAGGGTCCGAGGGTGGTGGGAGGATAGAAATCACATCCGCCCAACTTCCAAAGAGCAGCATCCCTCCCCG
ACCCATAGCCATGTTTTAAAGTACCTTCCGAAGAGAAGTGAAGGTTCAAGGACACTGGCCTTGAGGCCGAGGGAGC
AGCCATCACAACCTCACAGACCAGCACATCCCTTTTGAGACACCGCTTCTGCCACCACTCACGGACACATTTCTGCCT
AGAAAACAGCTTCTTACTGCTCTTACATGTGATGGCATATCTTACACTAAAAGAAATATTATTTGGGGGAAAACTACAAGT
GCTGTACATATGCTGAGAAATGCAGAGCATAATAGCTGCCACCCAAAAATCTTTTGAATAATCATTTCCAGACAACCTC
TTACTTTCTGTGTAGTTTTTAATTGTTAAAAAAGTTTTAAACAGAAGCACATGACATATGAAGCCTGCAGGACT
GGTCGTTTTTTGGCAATTCTTCCACGTGGGACTTGTCACAAGAATGAAAGTAGTGGTTTTTAAAGAGTTAAGTTACAT
ATTATTTTCTCACTTAAGTTATTTATGCAAAAGTTTTCTTGTAGAGAATGACAATGTTAATATTGCTTTATGAATTAA
CAGTCTGTTCTTCCAGAGTCCAGAGACATTGTTAATAAAGACAATGAATCATGACCGAAAG

Sequence ID No. 6: Human BEER protein variant (V10I)
MQLPLALCLICLLVHTAPRVVEGQWQAPKNDATITRELGEYPEPPELENNKTMNRAENGRRPHHPFETKDVSEYSC
RELHFTRYVTDGPCRSAPVTELVCSGQCGPARLLPNAIGRGKWRPSGPDFRCIPDRYARQVQLLCPGGEAPRARKVR
LVASCKCKRLTRPHNQSELDGTEAARPQGRKPRPRARSAXAQAELENAY

Sequence ID No. 7: Human Beer cDNA encoding protein variant (P38R)
AGAGCCTGTGCTACTGGAAGGTGGCTGCCCTCCTCTGGCTGGTACCATGCAGCTCCACTGGCCCTGTGTCTCGTCTGC
CTGCTGGTACACACAGCCTTCCGTGTAGTGGAGGGCCAGGGGTGGCAGGCGTTCAAGAATGATGCCACGGAAATCATCCG

-continued

CGAGCTCGGAGAGTACCCCGAGCTCCACCGAGCTGGAGAACAAAGACCATGAACCGGGCGGAGAACGGAGGGCGG
 CTCGCCACCACTTTTGTAGACAAAGACGTGTCCGAGTACAGTGCCTGCGGAGCTGCACTTCACCGCTACGTGACCGAT
 GGGCCGTGCCGAGCGCAAAGCCGGTCAACGAGCTGGTGTGCTCCGGCCAGTGCAGCCCGGCGCGCTGTGCCAACGC
 CATCGGCCGCGCAAGTGGTGGCGACCTAGTGGGCCGACTTCCGCTGCATCCCGACCGCTACCGCGCGAGCGGGTGC
 AGTGCTGTGTCCGGTGGTGAAGCGCCGCGCGCGCAAGGTGCGCTGGTGGCTTCGTGCAAGTGCAAGCGCCTACCC
 CGCTTCACAAACAGTCGGAGCTCAAGGACTTCGGGACCGAGGCGCTCGGCCGAGAAGGGCCGAAGCCGCGGCCCG
 CGCCCGGAGCGCAAAGCCAAACAGGCGAGGTGGAGAACGCCTACTAGAGCCCGCGCGCCCTCCCAACCGCGGGC
 GCCCCGGCCCTGAACCCGCGCCACATTTCTGTCTCTGCGCGTGGTTTGTATTGTTTATATTTTCATTGTAAATGCCGTC
 AACCCAGGCGAGGGGGTGAAGCTTTCAGGCCCTGAGGAATCCCGGGCGCGGCAAGGCCCCCTCAGCCCGCCAGCTG
 AGGGGTCCACGGGGCAGGGGAGGGAAATTGAGAGTCACAGACACTGAGCCACGCGAGCCCGCTCTGGGGCGCCTACCT
 TTGCTGGTCCCACTTCAGAGGAGGCAAGAAATGAAGCAATTTACCGCCCTGGGGTTTAAAGGAGCGGTGTGGAGTGG
 GAAAGTCCAGGACTGGTTAAGAAAGTTGGATAAGATTCCCCCTTGACCTCGCTGCCATCAGAAAGCCTGAGGCGTGC
 CCAGAGCACAAGACTGGGGGCAACTGTAGATGTGGTTTCTAGTCTGGCTTGCCTACTAATTTGCTGTAACTTTGAAC
 TACACAATTTCTTCCGGGACCTCAATTTCCACTTTGTAAATGAGGGTGGAGGTGGGAATAGGATCTCGAGGAGACTAT
 TGGCATATGATTCCAAGGACTCCAGTGCCTTTTGAATGGGCGAGGTGAGAGAGAGAGAGAAAGAGAGAAATGAATG
 CAGTTGCATTGATTCAAGTCCAAAGTCACTTCCAGAAATCAGAGTTGTGATGCTCTCTCTGACAGCCAAAGATGAAAAA
 CAAACAGAAAAAAGTAAAGAGTCTATTTATGGCTGACATATTTACGGCTGACAACTCCTGGAAGAAGCTATGCTG
 CTTCACAGCTGGCTTCCCGGATGTTTGGCTACCTCCACCCCTCCATCTCAAAGAAATAACATCATCCATTGGGGTAGA
 AAAGGAGAGGGTCCGAGGGTGGTGGGAGGGATAGAAATCAGATCCGCCCACTTCCCAAAGAGCAGCATCCCTCCCCG
 ACCCATAGCCATGTTTTAAAGTCACTTCCGAAGAGAAGTGAAAGGTTCAAGGACACTGGCTTGCAGGCCGAGGGAGC
 AGCCATCACAACCTCAGAGACAGCACATCCCTTTTGAAGACCGCTTCTGCCCACTCAGGACACATTTCTGCCT
 AGAAAAACAGCTTCTTACTGCTCTTACATGTGATGGCATATCTTACACTAAAAAATATTATTTGGGGAAAACTACAAGT
 GCTGTACATATGCTGAGAACTGCAGAGCATAATAGTGCACCCAAAAATCTTTTGAATAATTTCCAGACAACCTC
 TTACTTTCTGTAGTGTTTTAAATGTTAAAAAAGTTTAAACAGAGCACATGACATATGAAAGCCTGCAGGACT
 GGTCTTTTTTTGGCAATTTCTCCAGTGGGACTTGTCCACAAGAATGAAAGTAGTGGTTTTTAAAGAGTTAAGTTACAT
 ATTTATTTCTCACTTAAAGTTATTTATGCAAAAGTTTCTTTGTAGAGAATGACAATGTTAATATGCTTTATGAATTAA
 CAGTCTGTCTTCCAGAGTCCAGAGACATTTCTAATAAGACAATGAATCATGACCGAAAG

Sequence ID No. 8: Human Beer protein variant (P38R)
 MQLPLALCLVCLLVHTAFRVVEGQWQAFKNDATETIIPELGEYFEPPELENNKTMNRAENGRRPPHPPETKDVSEYSC
 RELHFTRYVTDGPCRSAPVTELVSCGQCPARLLPNAIGRGKWRPSGPDFRCIPDRYRAQRVQLLCPGGEAPRARKVR
 LVASCKKRLTRFNNQSELKDPGTEAARPQKGRKPRPRAESAKANQAELENAY

Sequence ID No. 9: Vervet BEER cDNA (complete coding region)
 ATGCAGCTCCCACTGGCCCTGTGCTTGTCTGCCTGCTGGTACACGACGCTTCCGTGTAGTGAGGGCCAGGGGTGGCA
 GGCTTCAAGAAATGATGCCACGGAATCATCCCCGAGCTCGGAGAGTACCCGAGCCTCCACCGAGCTGGAGAACACA
 AGACCATGAACGGGCGGAGAAATGGAGGGCGGCTCCCAACACCCCTTTGAGACCAAGACGTGTCGAGTACAGCTGC
 CGAGAGCTGCACTTCAACCGCTACGTGACCGATGGGCGGTGCCGAGCGCCAAGCCAGTCACCGAGTTGGTGTGCTCGG
 CCAGTGGGGCCCGCAGCCTGCTGCCAACGCCATCGGCCGCGGCAAGTGGTGGCGCCGAGTGGGGCCGACTTCCGCT
 GCATCCCGACCGCTACCGCGCGCAGCGTGTGCAGCTGCTGTGTCCTGGTGGTGCAGCGCGCGCGCAAGGTGCGC
 CTGTGGGCTCTGTCAAGTGCAAGCGCCTCACCGCTTCCACAACAGTCGGAGCTCAAGGACTTCGGTCCGAGGCGCG
 TCGGCCGCAAGGGCCGAAGCCGCGGCCCGCGCCCGGGGGCCAAAGCAATCAGGCCGAGCTGGAGAACGCCTACT

AG

-continued

Sequence ID No. 17: MluI - AviII restriction fragment used to make mouse Beer transgene

CGCGTTTGGTGAGCAGCAATATTGCGCTTCGATGAGCCTTGGCGTTGAGATTGATACCTCTGCTGCACAAAAGGCAATC

GACCGAGCTGGACCAGCGCATTCGTGACACGCTCCTTCGAACCTATTGCGAATGGAGTGTCAATTCATCAAGGACNGCC

TGATCGCAAATGGTGCTATCCACGAGCGGCAATCGAAAACCTCAGCCGGTGACCAATATCTACAACATCAGCCTTGGT

ATCCTGCGTGATGAGCCAGCGCAGAACAAGGTAACCGTCAGTGCAGATAAGTTCAAAGTTAAACCTGGTGTGATACAA

CATTGAAACGTTGATCGAAAACGCGCTGAAAACGCTGCTGAATGTGCGGCGCTGGATGTCACAAAGCAAATGGCAGCAG

ACAAGAAAGCGATGGATGAACCTGGCTTCTATGTCCGACGGCCATCATGATGGAATGTTTAAAGGTGGTGTATCTGG

CAGCAGTGCCGTCGATAGTATGCAATTGATAATTATATCATTTGCGGGTCTCTTCCGGCGATCCGCTTGTACGGGGC

GGCGACCTCGCGGGTTTTCTGCTATTTATGAAAATTTTCCGGTTTAAGGCGTTTCCGTCTCTCTCGTCATAACTTAATGT

TTTTATTTAAAAATACCTCTGAAAAGAAAGGAAACGACAGGTGCTGAAAGCAGCTTTTGGCCCTCTGTCGTTTCTTTC

TCTGTTTTTGTCCGTGGAATGAACAATGGAAGTCAACAAAAGCAGAGCTTATCGATGATAAGCGGTCAAACATGAGAAT

TCGCGGCCGCATAATACGACTCACTATAGGGATCGACGCTACTCCCGCGCATGAAGCGGAGGAGCTGGACTCCGCATG

CCCAGAGCGCCCCCAACCCCCAAAGTGCTGACCTCAGCCTCTACCAGCTCTGGCTTGGGCTTGGCGGGGTCAAGGC

TACCACGTTCTCTTAACAGGTGGCTGGGCTGTCTCTTGGCCGCGCTCATGTGACAGCTGCCTAGTTCTGCAGTGAGGTC

ACCGTGGAATGTCTGCTTCTGTTGCCATGGCAACGGGATGACGTTACAATCTGGGTGTGGAGCTTTCTGTCCGTGTCA

GGAAATCCAATACCTAAAATACCTAGAAGAGGAAGTAGCTGAGCCAAGGCTTTCTGGCTTCTCCAGATAAAGTTTG

ACTTAGATGGAATAAATAAATGATAAGACCCGAGCCATCTGAAAATTCCTCTAATTGCACCACTAGGAAATGTGTA

TATTATTAGCTCGTATGTGTTCTTATTTTAAAAAGAAAACCTTAGTCATGTTATTAATAAGAAATTTCTCAGCAGTGGA

GAGAACCAATATTAAACACCAAGATAAAAGTTGGCATGATCCACATTGCAGGAAGATCCACGTTGGGTTTTTCATGAATGTG

AAGACCCATTTATTAAAGTCCTAAGCTCTGTTTTTGCACACTAGGAAGCGATGGCCGGGATGGCTGAGGGGCTGTAAGG

ATCTTTCAATGTCTTACATGTGTGTTTCTGTCTGCACCTAGGACCTGCTGCCTAGCCTGCAGCAGAGCCAGAGGGGTT

TCACATGATTAGTCTCAGACACTTGGGGCAGGTTGCATGTACTGCATCGCTTATTTCCATACGGAGCACCTACTATGTG

TCAAACACCATATGGTGTCTACTCTTCAGAACGGTGGTGGTCATCATGGTGCATTGCTGACGGTTGGATTGGTGGTAGA

GAGCTGAGATATATGGACGCACTCTTCAGCATTCTGTCAACGTGGCTGTGCATTCTTGCTCCTGAGCAAGTGGCTAAACA

GACTCACAGGGTCAGCCTCCAGCTCAGTCGCTGCATAGTCTTAGGGAACCTCTCCAGTCTCCCTACCTCAACTATCCA

AGAAGCCAGGGGGCTTGGCGGTCTCAGGAGCCTGCTTGTGTTGGGGACAGGTTGTTGAGTTTATCTGCAGTAGGTTGCCT

AGGCATAGTGTGAGGCTGATGGCTGCCTTGGAGAACACATCCTTTGCCCTCTATGCAAATCTGACCTTGACATGGGGC

GCTGCTCAGCTGGGAGGATCAATGCATACCTAAAGCCAAGCCTAAAGCTTCTTCGTCCACCTGAAACTCCTGGACCAAG

GGGCTTCCGGCAGATCTCTCAGGCCAGTGAGGAGTCTGTGTGAGCTGCACTTCCAATCTCAGGGCGTGAGAGGCAGA

GGGAGGTGGGGCAGAGCCTTGACGCTCTTCTCTCCATCTGGACAGCGCTCTGGCTCAGCAGCCATATGAGCACAGGC

ACATCCCCACCCACCCACCTTTCTGTCTCTGAGAAATTTAGGCTCTGTTACGGGGGGGGGGGGGGGGGGCAGTCC

TATCCTCTCTTAGGTAGACAGGACTCTGCAGGAGACACTGCTTTGTAAGATACTGCAGTTTAAATTTGGATGTTGTGAGG

GGAAAGCGAAGGGCCTCTTTGACCATTCAGTCAAGGTACCTTCTAACTCCATCGTATGGGGGGCTACTCTAGTGTCTAG

ACATTGCAGAGAGCCTCAGAACTGTAGTTACCAAGTGTGGTAGGATTGATCCTTCAGGGAGCCTGACATGTGACAGTTCCA

TTCTTCAACCACTCACCAGAACATTTATTCAGTACCTACCCCGTAACAGGCACCGTAGCAGGTACTGAGGGACGGACCACT

CAAAGAACTGACAGACCAGGCTTTGGAATATAAACACCAAAGCATCAGGCTCTGCCAACAGAACTCTTTAACTACTCA

GGCCCTTTAACTACTCAGGACCCCCACCCCAAGCAGTTGGCACTGCTATCCACATTTTACAGAGAGGAAAACTA

GGCACAGGACGATATAAGTGGCTTGTCTAAGCTTGTCTGCATGGTAAATGGCAGGGCTGGATTGAGACCCAGACATTCCA

ACTCTAGGGTCTATTTTTCTTTTTCTCGTTGTTGGAATCTGGGTGTTACTGGGTAAACTCAGGCTAGCCTCACACTCAT

ATCCTTCTCCATGGCTTACGAGTGTAGGATTCCAGGTGTGTGCTACCATGTCTGACTCCCTGTAGCTTGTCTATACCA

-continued

TAAACATTGAAGAAGACAGTAGATGCCAATTTTAAGCCCCACATGCACATGGACAAGTGTGCGTTTGAACACACATAT
GCACCTACTGTGAACCGGCATGCACACTCGGGCTTATCACACACATAATTGAAAGAGAGAGTGAGAGAGGAGAGTGCAC
ATTAGAGTTACAGGAAAGTGTGAGTGAGCACACCCATGCACACAGACATGTGTCCAGGGAGTAGGAAAGGAGCCTGGG
TTTGTTGATAAGAGGGAGCCATCATGTGTTCTAAGGAGGGCGTGTGAAGGAGGCGTTGTGTGGGCTGGGACTGGAGCAT
GGTTGTAACAGCATGCTCCCTGTGGGAAACAGGAGGGTGGCCACCCTGCAGAGGGTCCCACTGTCCAGCGGGATCAGT
AAAAGCCCCTGCTGAGAATTTAGGTAATAGCCAGAGAGAGAAAGGTAGGAAAGTGGGGGACTCCCATCTCTGATGTAG
GAGGATCTGGGCAAGTAGAGGTGCGTTTGAGGTAGAAAGAGGGGTGCAGAGGAGATGCTCTTAATCTGGGTCAGCAGTT
TCTTTTCAAATAATGCCGTGTGAGGAGGTGTAGGTGGCCATTCACTCACTCAGCAGAGGGATGATGATGCCCGGTGGA
TGCTGGAATAGGCCGAGCATCAACCTTGGCTCTGGAAGAACTCCATCTTTCAGAGGAGAGTGGATCTGTGTATGGCCAG
CGGGGTACAGGTGCTTGGGGCCCCGGGGACTCCTAGCACTGGGTGATGTTTATCGAGTGTCTTGTGTGCCAGGCAC
TGGCCTGGGGCTTTGTTTCTGCTCTGTTTTGTTTCTGTTTTTGGACAGACTCTTGTATGTATCCGTGTCAATCTTGG
AATCTCACTGCATAGCCAGGCTGCCGAGAGAGGGAGGGCAATAGGCCTTGTAAGCAAGCCACACTTCAGAGACTAGAC
TCCACCCTGCGAATGATGACAGGTACAGAGCTGAGTTCCGGAAGATTTTTTTCCAGCTGCCAGGTGGAGTGTGGAGTGGC
AGCTAGCGGCAAGGGTAGAGGGCGAGCTCCCTGTGCAGGAGAAATGCAAGCAAGAGATGGCAAGCCAGTGAAGTAAAGCAT
TCTGTGTGGGAGCAGGTGGATGAAGAGAGAGGCTGGGCTTTCGCCTCTGGGGGGGGGTGAGGGGTGGGGATGAGGTGA
GAGGAGGGCAGCTCCCTGCAGTGTGATGAGATTTTTCTTGACAGTGACCTTTGGCCTCTCCCTCCCACTTCCCTTCTT
TCCTTCTTCCACCATTGCTTTCCCTTGTCTTGAGAAATCTGAGTTTCCACTTCACTGGTGATGCAGACGGAACAGAA
AGCCGT
GTATGTGTGTGATGGGAATGGCTCATAGTCTGCAGGAAGGTGGGCAGGAAGGAATAAGCTGTAGGCTGAGGCAGTGTGG
GATGCAGGGAGAGAGGAGAGGGGATACCAGAGAAGGAAATTAAGGGAGCTACAAGAGGCAATTGTTGGGGTGTGTGTG
TGTGTGTGTTTATATTTGTATTTGAAATACATTCTTTTAAAAATACTTATCCATTTATTATTTTATGTGCACGT
GTGTGTGCTGCATGAGTTTATGTGTGCACAGTGTGTGCGGGAACCTTGGAGGCCACAAGGGGCATCTGATCCCCTGG
AACTGGAGTTGGAGGAGGTGTGAGTCCCTGACATGTTTGTGGGAACGAAACCCGGTCTTATGAAGAGCAGGAAGT
GCAGTTATCTGCTGAGCCATCTCTCCAGTCTGAAATCCATTCTCTTAAATACAGTGGCAGAGACATGATGGGATTTA
CGTATGGATTTAATGTGGCGGTCAATTAAGTTCCGGCACAGGCAAGCACCTGTAAAGCCATCACCACAACCGCAACAGTGA
ATGTGACCATCACCCCCATGTTCTTATGTCCCCGTGCCCTCCATCTCCATTCTCAAGCACCTCTTGCTCTGCCTCTG
TCGCTGGAGAACAGTGTGCATCTGCACACTCTTATGTGAGTGAAGTACACAGCCTGCACCCCTTCTGGTCTGAGTATT
TGGGTTCTGACTCTGCTATCACACTACTGTACTGCATTCTCTCGCTCTCTTTTTTAAACATATTTTATTGTTTGT
GTGTATGCACATGTGCCACATGTGTACAGATACTATGGAGGCCAGAAGAGGCCATGGCCGTCCCTGGAGCTGGAGTTACA
GGCAGCGTGTGAGCTGCTGGTGTGGGTGCTGGGAACAACTTGAATCTAAAGCAAGCACTTTAACTGCTGAGGCAGC
TCTCAGTACCCTTCTTCAATTTCCGCTGGGTTCATTGTATGGACACATGTAGCTAGAATATCTTGTCTTATCTAATTA
TGTACATTGTTTGTGCTAAGAGAGAGTAATGCTCTATAGCCTGAGCTGGCCTCAACCTTGCCATCTCTGCTCAGCC
TCCTCCTCTGAGTGTAGGATGACAGGCGAGTGGTAACCTACATGGTTTCTGTTTGTCTAAGACTGAAGGATAACAT
TCATACAGAGAAGGTCTGGGTCAAAAGTGTGAGTTCACGAAATGGCAACCCGTGATCAAGAAACAAATCAGGGG
CTGGAGAGATGGCACTGACTGCTCTCCAGAGTCCGGAGTTCAATTCCAGCAACCATGTTGGCTCACAGCCATCTA
TAACGAGATCTGACGCCCTCTTCTGGTGTGTCTGAAGACAGCTACAGTGTACTCACATAAAATAAATAAATCTTTAAAC
ACACACACACACAATTACCACCCAGAAAGCCCACTCCATGTTCCCTCCACGTCTCTGCTACAGTACTCCAGGTT
ACCACTGTTTACGGCTTCTAACAACCTGGTTTACTTGGGCTCTTTTCTGCTCTGTGGAGCCACACATTTGTGTGCTCAT
ACACGTTCTTTCTAGTAAAGTTGCATATTACTCTGCGTTTTTACATGTATTTATTTATGTAGTGTGTGTGTGTGGG
CCATGCATGGCAGTGTGTGGGATGTGAGATATTGTGAACAGGGGACAGTTCTTTCTTCAATCATGTGGGTTCAG

-continued

AGGTTGAAGCTCAGGTCATCATGTGTGGCAGCAAAATGCCTTTACCCACTGAGACATCTCCATATCTTTTTTTTCCCTG
AGGTGGGGGCTGTTCATAGCCCAAACTGGCTTTGCACTTGCAAGTGACTCCCTGCTCCACCTCTTAGAGTA
TTGGAATTACGATGTGTACTACCACACCTGACTGGATCATTAAATCTTTGATGGGGCGGGGAAGCGCACATGCTGCAGG
TGAAGGGATGACTGGACTGGACATGAGCGTGAAGCCAGAGAACAGCTTCAGTCTAATGCTCTCCCAACTGAGCTATTTC
GGTTTGCCAGAGAACTTACAGAACTTCTCAGTGCCATGTGGATTGCGGGTTGAGTTCAACTCATCAGCTTGACAT
TGGCTCCTCTACCCACTGAGCCTTCTACTACTCTCTACCTAGATCATTAAATCTTTTTTAAAAAGACTTATTAGGGGGC
TGGAGAGATGGCTCAGCCGTTAAGAGCACCGAATGCCCTTCCAGAGGTCCTGAGTTCAATTCACAGCATGCCATTGCTGG
GCAGTAGGGGGCGCAGGTGTTCAACGTGAGTAGCTGTGTCAGTTTTCGCGGTGGAGAACCTCTTGACACCCTGCTGTC
CCTGGTCATTCTGGTGGGTGCATGTTGATATGCTTGTGTATGGAAGACTTTGACTGTTACAGTGAAAGTTGGGCTTCCA
CAGTTACCACGCTCTCCCTGTTTCTTGAGGCGGGTGTCTGTCATTGCGCGAGGGCTACAGCCGCTCCCAACGCTA
GTTATCGCTACCTCATGATGCGGCAGAAGATGGACGTCAAGTCTGCTCTGAGTACTGTGAGGCAGAATCGTGAGATCGG
CCCCAACGATGGCTTCTTGCCCAACTCTGCCAGCTCAATGACAGACTAGCCAAGGAGGGCAAGGTGAAACTCTAGGGTG
CCACAGCCTCTTTTGAGAGGCTGACTGGGAGGGCCCTGGCAGCCATGTTTAGGAAACACAGTATACCCACTCCCTGC
ACCACCAGACAGTGGCCACATCTGTCCTCCTGCTCGGGGGCCACTCCACCCTTAGGGAGCACATGAAGAAGCTC
CCTAAGAACTTCTGCTCCTTAGCATCTCTTCTGTAATTATGCTCTCTCCCTGAGGTGAGGTTGAGGTTTATGTCCTG
TCTGTGGCATAGATACATCTCAGTGACCCAGGTGGGAGGGCTATCAGGGTGCATGGCCCGGACACGGGCACCTTCAT
GACCCCTCCCCACCTGGGTTCTCTGTGTGTCAGAACCCAGAGCCTGTAAGGAACTATGCAAAACACAGGCCCTG
ACCTCCCCATGTCTGTTCCTGCTCAGAGCCGACACGCCCTGCTGAGGCAGAGCAATGACATTAAGTTCTGAAGCAG
AGTGAGATAGATTAGTGACTAGATTCCAAAAAGAGGAAAAAAGGCTGCATTTTAAATATTCTCTTAGAATTAA
AGATACTACATAGGGGCCCTTGGGTAAGCAATCCATTTTCCAGAGGCTATCTTGATTCTTTGGAATGTTTAAAGTG
GCCTTGCCAGAGAGCTTACGATCTATATCTGCTGCTTCAGAGCCTTCCCTGAGGATGGCTCTGTCTCTTGCTTGTAGA
AGAGCGATGCCTTGGGAGGGTTTCCCTTTTCAGAAATACAGGGTGTAAGTCCAGCCTATTACAAACAAACAAACAAA
CAAACAAACAAAGGACCTCCATTTGGAGAAATGCAAGGATTTATCTGAATTATAGTGTGGTGAGTTCAAGTCATCAC
GCCAAGTGCTTGCCATCTGGTTGCTATTCTAAGAATAATTAGGAGGAGGAACCTAGCCAATTGCAGCTCATGTCCGTGG
GTGTGTGCACGGGTGCATATGTTGGAAGGGGTGCTGTCCCTTGGGGACAGAAGGAAATGAAAGGCCCTCTGCTCAC
CCTGGCCATTTACGGGAGGCTCTGCTGGTTCCACGGTGTCTGTGCAGGATCTGAAACTGACTCGCTGGACAGAAACGAG
ACTTGGCGGCACCATGAGAATGGAGAGAGAGAGCAAGAAAGAAACAGCCTTTAAAGAACTTTCTAAGGGTGGTTTT
TGAACCTCGCTGGACCTTGATGTGTGCACATTTGCCAGAGATTGAACATAATCCTCTTGGGACTTCACGTTCTCATTAT
TTGTATGTCTCCGGGTACAGCAGAGCCGTACGCCACCCAGCACCAGCAGCATAGGCGTCTATAAAAGCCCATTT
TATGAGAACAGAGCTGTTTGTAGTACCCGTGTATAGAGAGAGTTGTTGCTGGGGACCCGGATCCAGCAGCCTGGT
TGCTGCTGTAGGATGCTTACAGGAGTTTGACAGAAACCTTCTTGGAGGAAAGAAATATCAGGGATTTTGTGTA
ATATTTCAAATTCAGCTTTAAGTGTAAGACTCAGCAGTGTTCATGGTTAAGGTAAGGAACATGCCTTTTCCAGAGCTGCT
GCAAGAGGCAGGAGAAGCAGACCTGTCTTAGGATGTCACTCCAGGGTAAAGACCTCTGATCAGCAGGAGCAGAGCTG
TGCAGCCTGGATGGTCAATTGCTCCCTATTCTGTGTGACACAGCAACCTGGTGCATAGGGCTGGTCACTCTTTTTTT
TTTTTTTTTTTTTTTTTTGGCCAGAAATGAAGTGACCATAGCCAAGTTGTGTACCTCAGTCTTTAGTTTCCAAGCGGT
CTCTTGCTCAATACAATGTGCATTTCAAAATAACACTGTAGAGTTGACAGAACTGGTTCATGTGTTATGAGAGAGGAAAA
GAGAGGAAAGAAACAAACAAACAAACACCAACAAACAAACATCTGGGCTAGCCAGGCATGATTGCAATGTCTACAG
GCCAGTTTCATGAGAGGCAGAGACAGGAAGACCGCCGAAAGGTCAAGGATAGCATGGTCTACGTATCGAGACTCCAGCCA
GGGCTACGGTCCCAAGATCTAGGTTTGGATTTTGGGCTTTGGTTTTTGGAGACAGGTTTCTCTGTGTAGCCCTGGCTG
TCCTGGAACCTGCTCTGTAGACCAGGCTGGCTCAAACTTAGAGATCTGCCTGACTCTGCCTTTGAGGGCTGGGACGAAT

-continued

GCCACCACCTGCCCACTAAGATTCCATTAAAAAAGTTCAAGATAATTAAGAGTTGCCAGCTCGTTAAAGCTAA
GTAGAAGCAGTCTCAGGCCGTGCTGTGAGGCTGTCTTGGCTTGACCTGAAATCTGCCCCAACAGTGTCCAAGTGCA
CATGACTTTGAGCCATCTCCAGAGAAGGAAGTGAATAATGTGGCTCCCCAGTCGATTGGGACACAGTCTCTCTTTGTCTA
GGTAACACATGGTGACACATAGCATGAACTCTCACTCTGAGGGTGGGTTTCCTCCCCCTGCCTCTCTGGGTGGTC
ACCCCATAGGACAGCCACAGGACAGTCACTAGCACCTACTGGAACCTCTTTGTGGGAACATGAAGAAAGAGCCTTTGGG
AGATTCTCGGCTTTCCATTAGGGCTGAAAGTACAACGGTCTTGGTTGGCTTTGCCCTCGTGTATATAAACTAGCTACTA
TTCTTCAGGTAAAAATACCGATGTTGTGGAAGCAACCCCGTGGCTGCCGTGAGTAGGGGTGGGTTGGAATCCTG
GATAGTGTCTATCCATGGAAGTGGTGAATAGGAATTAAGGGTGTTCCTCCCCCCCCAACCTCTTCCTCAGACCCAG
CCACTTTCTATGACTTTATAAACATCCAGGTAAAAATTACAAACATAAAAAATGGTTTCTCTCTCAATCTCTTAAAGTCTG
CCTGCCCTTTCCAGGGTAGGTCTGTCTTAGCTGTCTATTGTCTTGAGAGCACAGACTAACACTTACCAAATGAGGG
AACTCTTGGCCCATACTAAGGCTCTTCGGGCTCCAGCACTCTTAAGTTATTTAAGAATTCACATTGGCCTTTAGCAC
ACCCGCCACCCCAAGTGGGTGTGGAATATGCCATGGCCAGCAGGGGGCACTGTTGAGCGGGTGCTTTCCACCCTTAAG
TTGCTTATAGTATTTAAGATGCTAAATGTTTAAATCAAGAGAAGCACTGATCTTATAATACGAGGATAAGAGATTTTCTC
ACAGGAAATGTCTTTTTCATAATCTTTTACAGGCTTTGTCTGATCGTAGCATAGAGAGAATAGCTGGATATTTAACT
TGTATTCCATTTTCTCTGCCAGCGTTAGGTTAACTCCGTAAAAAGTGATTCACTGGACCGAAGAGGCTCAGAGGGCAGG
GGATGGTGGGTGAGGCAGAGCACTGTCACTGCCAGGCATGGGAGGTCCTGCCATCCGGGAGGAAAAGGAAAGTTTAGC
CTCTAGTCTACCAACAGTGTAAACGCACCTTAAAGTTGTAACCAAAATAAATGTCTTACATTACAAAGACGCTGTGTTTG
TGTTTCTCTTTGTGTGTTTGGGCTTTTATGTGTGCTTTATAACTGCTGTGGTGGTGTGTTGTTAGTTTGGAGTAGGA
TCTCAGGCTGGCCTTGAACCTCTGATCGCTGCCCTGCCCTGCCCTGCCCTGCTCCTGCCCTCCAAGTGTAGGACT
AAAAGCACATGCCACCACACAGTACAGCATTTTCTAACATTTAAAAATAATCACCTAGGGGCTGGAGAGAGGGTTCCA
GCTAAGAGTGACACACTGCTCTTGGGTAGGACCTGAGTTTAGTTCCAGAACCTATACTGGGTGGCTCCAGGTCCAGAGGA
TCCAGGACCTCTGGCCTCCATGGGCATCTGCTCTTAGCACATACCCACATACAGATACACATAAAAAATAAAATGAAGC
CTTTAAAAACCTCTTAAACCTAGCCCTTGGAGGTACGACTCTGGAAGCTGGCATACTGTGTAAGTCCATCTCATGGTG
TTCTGGCTAACGTAAGACTTACAGAGACAGAAAAGAACTCAGGGTGTGCTGGGGTGGGATGGAGGAAGAGGGATGAGT
AGGGGGAGCAGGGGAACTTGGGCAGTGAATAATCTTGCAGGACACTAGAGGAGGATAAATACCAGTCAATGCACCCAC
TACTGGACAACCTCAGGGAATATGCTGGGTGAAAAGAGAAGGCCCCAGGTATTGGCTGCATTGGCTGCATTGCGTAAC
ATTTTTTAAATTTGAAAAGAAAAGATGTAAATCAAGGTTAGATGAGTGGTTGCTGTGAGCTGAGAGCTGGGGTGAAGTA
GACATGTGGACAACCTCATCAAAAAGCAGAGAAAGAACGGGCTGTGGTGACAGCTACCTCTAATCTCCACCTCCGGGAG
GTGATCAAGGTTAGCCCTCAGCTAGCTGTGGTGATGAGACCTGTTTCAAAAACCTTTAATAAAGAAATATGAAAAA
GACATCAGGGCAGATCTTGGGGCCAAAGCGGACAGGCGAGTCTCGTGGTAAGTCTGTGTAAGCGGATGCATGAGCA
CGTGCCGACAGCATCATGAGAGAGCCCTAGGTAAGTAAGGATGGATGTGAGTGTGCGGCTCGGCGCACTGCACGTCTT
GGCTGTGGTGCTGAGCTGGCATCTTTGGTGAGCTGTGGAGGGAAATGGGTAGGGAGATCAAAAATCCCTCCGAATTAT
TTCAAGAACTGTCTATTACAATTATCTCAAAATATTAATAAAAAAGAAATTAAAAAACAAAAACCTATCCAGGTGTG
GTGGTGTGCACCTATAGCCACGGGCACCTGGAAGCTGGAGCAAGAGGATGGCGAGTTGAAGGTATCTGGGGCTGTACA
GCAAGACCGTGTGCTCCCAAAACAAACAGCAAAACCAATTATGTACACAAGAGTGTATTAGTGAGCGGCCCTCGCT
GAGAGCATGGGGTGGGGTGGGGTGGGGGACAGAAATATCTAACTGCAGTCAATAGGGATCCACTGAGACCTCGGGG
TTGACTGCAGCTTAACCTTGGGAAATGATAAGGGTTTGTGTGTGAGTAAAGCATCGATTACTGACTTAACCTCAATGA
AGAAAAAGAAAAAGAAAAACAACAAAGCCAAACCAAGGGGTGGTGAGATGGCTCAGTGGGTAAGAGCACCCGACTGC
TCTCCGAAGGTCCAGAGTTCAAAATCCAGCAACACATGGTGGCTCACAACCATCTGTAAACAGATATGATGCCCTCTT
CTGGTGTGCTGAAGACAGCTACAGTGACTTACATATAATAAATAAATCTTAAAAAAGAAAAAAGCCAAA

-continued

CAGGTGGCACCACCTTGCCCTGAGCTTGACCTGACCCAGCTTTGCCTCATTCTGAGGACAGCAGAACTGTGGAGGC
AGAGCCAGCACAGAGAGATGCCTGGGGTGGGGTGGGGTATCACGCACGGAAGTAGCAGCAATGAATGGGGTGGGGTGG
CAGCTGGAGGGACACTCCAGAGAAATGACCTTGCTGGTCACCATTTGTGTGGGAGGAGCTCATTTCAGCTTGCCAC
CACATGCTGTCCCTCCTGTCTCCTAGCCAGTAAGGGATGTGGAGGAAAGGGCCACCCCAAAGGAGCATGCAATGCAGTCA
CGTTTTTGCAGAGGAAGTGCTTGACCTAAGGGCACTATTCTTGGAAAGCCCCAAAGCTAGTCCTTCCTGGGCAACAGG
CCTCCCCACATACCACCTCTGACGGGTGAGTAAATTAAGCCAGCCACAGAAGGGTGGCAAGGCCACACCTCCCCCT
GTTGTGCCCCCCCCCCCCGCGTGAAGTGATCTCGCCCTCTGCCCTCTGGCTTTGGTACTGGGATTTTTTTTTCCTT
TTATGTGATATTGATCTGACACCATGGAACTTTGGAGGTAGACAGGACCCACACATGGATTAGTTAAAAGCCTCCCAT
CCATCTAAGCTCATGCTAGGAGATAGAGCATGTCCAAGAGAGGAGGCGAGGCATCAGACCTAGAAGATATGGCTGGGCAT
CCAACCCAAATCTCCTTCCCGGAGAACAGACTCTAAGTCAGATCCAGCCACCTTGAGTAACCAGCTCAAGGTACACAGA
ACAAGAGAGTCTGTATACAGCAGGTGCTAAACAAATGCTTGTGGTAGCAAAAGCTATAGGTTTGGGTGAGAACTCCGA
CCCAAGTCGCGAGTGAAGAGCGAAAGGCCCTTACTCGCCACCGCCCCCGCCCCACCTGGGGTCTCTATAACAGATCACTT
TCACCCCTGCGGGAGCCAGAGAGCCCTGGCATCTAGGTAGCCCCCCCCCGCCCCCGCAAGCAGCCAGCCCTGCC
TTTGGGGCAAGTTCTTTCTCAGCCTGGACCTGTGATAATGAGGGGTGGACGCGCCGCTTTGGTCTGCTTCAAGTCT
AATGAATTCTTATCCCTACCACCTGCCCTTCTACCCGCTCCTCCACAGCAGCTGTCTGATTATTACCTTCAATTAAAC
CTCCACTCCTTTCTCCATCTCCTGGGATACCGCCCTGTCTCCAGTGGCTGGTAAAGGAGCTTAGGAAGGACCAGAGCCAG
GTGTGGCTAGAGGCTACCAGCAGGGCTGGGGATGAGGAGCTAAACTGGAAGAGTGTGTGGTTAGTAGGCACAAAGCCTT
GGGTGGGATCCCTAGTACCGGAGAAGTGGAGATGGGCGCTGAGAAGTTCAAGACCATCCATCTTAACTACACAGCCAGT
TTGAGGCCAGCCTGGGTACATAAAAAACCAATCTCAAAAGCTGCCAATTCGATTCTGTGCCACGTAGTGGCCGATGTA
ATAGTGGATGAAGTCGTTGAATCTGGGGCAACCTATTTTACAGATGTGGGAAAAGCAACTTTAAGTACCCTGCCACA
GATCACAAGAAAGTAAGTGACAGAGCTCCAGTGTTCATCCCTGGGTTCCAAGGACAGGAGAGAGAAAGCCAGGGTGGG
ATCTCACTGCTCCCCGGTGCCCTCTCTATAATCCATACAGATTGAAAGCGCAGGGCAGGTTTGGAAAAGAGAGAAG
GGTGAAGGAGCAGACAGTCTGGCCTAGGCTGCAGCCCCACGCATCCCTCTCTCCGCAGATGTGTCCGAGTACAGCT
GCCCGAGCTGCACTACACCGCTTCTGTACAGACGGCCATGCCGACGCCAAGCCGGTCAACCGAGTTGGTGTGCTCC
GGCCAGTGGCGCCCCCGCGGCTGTGCCAACGCCATCGGGCGCTGAAGTGGTGGCGCCGAACGGACCGGATTTCCG
CTGCATCCCGGATCGCTACCGCGCGCAGCGGTGCAGTGTGTGCCCCGGGGCGCGCGCCGCGCTCGCGCAAGGTGC
GTCTGGTGGCTCTGTGAAGTGAAGCGCTCACCCGCTTCCACAACAGTCCGAGCTCAAGGACTTCGGGCGGAGACC
GCGCGGCCGAGAAGGGTGCAGCGCGCGGCCGCGCGCGGGGAGCCAAAGCCAACAGCGGAGCTGGAGAACGCCCTA
CTAGAGCGAGCCCGCCCTATGACAGCCCCCGCGGATCCGATTCTGTTTCAAGTGTAAAGCTGCAGCCAGGCCAGGGGT
GCCAAACTTTCCAGACCGTGTGGAGTTCCAGCCAGTAGAGACCGCAGGTCTTCTGCCCCGCTGCGGGGATGGGGAGG
GGGTGGGGTTCCCGGGGCCAGGAGAGGAAGCTTGAGTCCAGACTCTGCCTAGCCCCGGGTGGGATGGGGGTCTTTCTA
CCCTCGCCGACCTATACAGACAAGGCAGTGTTCACCTTAAAGGGAAGGAGTGTGAACGAAAGACCTGGGACTGG
TTATGGAGCTACAGTAAGATCTACTCTCCACCCAAATGTAAAGCCTGCGTGGGCTAGATAGGGTTTCTGACCTGACC
TGGCCACTGAGTGTGATGTGGGCTACGTGTTCTCTTTGGTACGGTCTTCTTGTAAAAAGGGACCGGAACCTGCT
GAGATTCCAAGGATTGGGGTACCCCTGTAGACTGGTGAGAGAGAGGAGAACAGGGGAGGGTTAGGGGAGAGATTGTGG
TGGGCAACCGCTAGAAGAAGCTGTTTGTGGCTCCAGCCTCGCCGCTCAGAGGTTTGGCTTCCCCACTCTTCTCTC
TCAAACTCTGCCTTCAAACTCATATCTGGGATAGGGAAGGCCAGGGTCCGAGAGATGGTGAAGGGCCAGAAATCACACTC
CTGGCCCCCGAAGAGCAGTGTCCCGCCCCAACGCTTGTCTATATTGTAAAGGATTTTCTACACAACAGTTTAAGGT
CGTTGGAGGAAACTGGGCTTGCCAGTCACTCCATCTTGTCTCTTGCAGGACACCACTCTCTGCTGCCACCCACGG
ACACATTTCTGTCTAGAAAACAGAGCGTCTGTGTCTCTCTGAGACAGCATATCTTACATTAAAAAGAATAATACGGG

-continued

GGGGGGGGCGGAGGGCGCAAGTGTATACATATGCTGAGAAGCTGTCAGGCGCCACAGCACCCACAATCTTTTGT
AAATCATTTCCAGACACCTCTTACTTTCTGTGTAGATTTTAAATGTTAAAAGGGGAGGAGAGAGCGTTTGTAAACAGAA
GCACATGGAGGGGGGGTAGGGGGGTTGGGGCTGGTGACTTTGGCGAACTTTCCATGTGAGACTCATCCACAAAGACTGA
AAGCCGCGTTTTTTTTTTAAAGATTGAGTGACATATTTATTTCTCATTTAAGTTATTTATGCCAACATTTTCTTG
TAGAGAAAAGCAGTGTAAATATCGCTTTGTGAAGCACAAGTGTGTGGTTTTTGTGTTTTTGTGTTTTTCCCCGACCAGA
GGCATTTGTTAAATAAGACAATGAATCTCGAGCAGGAGGCTGTGGTCTTGTTTGTCAACCACACACAATGTCTCGCCACT
GTCACTCACCTCCCTTCCCTTGGTCACAAGACCCAAACCTTGACAACACCTCCGACTGCTCTCTGGTAGCCCTTGTGGCA
ATACGTGTTTCTTTGAAAAGTCACATTCATCCTTTCTTTGCAACCTGGCTCTCATTTCCCCAGCTGGGTCTATGTCAT
ACCCCTCACCCAGCCTCCCTTTAGCTGACCACTCTCCACACTGTCTTCCAAAAGTGACGTTTACCAGGCGAGTTCCCT
GGTCCAGGTCATCCCATGTCTCTCTGCTCCAGACCTTCTCCCAAGATGTTCACTCTCCACTCCATCAAGCCCC
AGTGGCCCTGCGGCTATCCCTGTCTCTCAAGTAGCTGAATCTACTGTGACACCACATGAATTCCTCCCTGTCTTA
AGGTTCAATGGAATCTTGCCTGCCCTGAACCTTCCAGGACTGTCCAGCGTCTGATGTCTCTCTCTTTGTAAGCCC
CACCCTCATTTGATTCCCAATCTAGATCTTCCCTTGTTCATTCCTTACGGGATAGTGTCTCATCTGGCCAAGTCTCT
GCTTGATATTGGGATAAATGCAAGCCAAGTACAATTGAGGACAGTTCATCATTTGGCCAAGCTTTTCAAAATGTGAA
TTTTACACCTATAGAAGTGTAAAAGCCTTCCAAAGCAGAGGCAATGCCTGGCTCTTCTTCAACATCAGGGCTCTGTCTT
TATGGGTCTGGTGGGTAGTACATTCATAAACCACACTAGGGGTGTGAAAGCAAGATGATTGGGAGTTCGAGGCCAAT
CTTGGCTATGAGGCCCTGTCTCAACCTCTCTCCCTCCCTCCAGGGTTTTGTTTTGTTTTGTTTTTGTATTGAACTG
CAACACTTTAAATCCAGTCAAGTGCATCTTTGCGTGAGGGGAATCTATCCCTAATATAAGCTTCCATCTTGATTGTGT
ATGTGCACACTGGGGGTGAACTTGGGCTTTGTACCTGCCGGCAAGCTCTCTACTGCTCTAAACCCAGCCCTCACTGG
CTTTCTGTTTCACTCCCAATGAATTCCTTAAATGAATTATCAATATCATGTCTTTGAAAAATACATTGAGTGTCTGT
GGTGTCCCTGTGGTTCCAGATTCAGGAAGGACTTTTCAGGGAATCCAGGCATCCTGAAGAAATGCTTTAGAGCAGGAGGC
CATGGAGACCTTGCCAGCCCCACAAGGCAGTGTGGTGACAGGGGTGAGGATGGAGGCAGGCTTGCAATGAACTGAGA
CAGGGTACTCAGGATTAAGAAAGCTTCCCCAAAACAATTCAGATCAGTTCCTGGTACTTGCACTGTTCAGCTATGCA
GAGCCAGTGGGCATAGGTGAAGACACCGTTGTACTGTCTACTTAAGTGTCTTCAAGCCGGCAGAGACAATAAT
GTTATGGTGACCCAGGGGACAGTGATTCCAGAGGAACACAGAAGAGAGTGTCTGTAGAGGCTGCCGAAGGAGAAGGG
GTCCAGACTCTCTAAGCAAGACTCCACTCACATAAGACACAGGCTGAGCAGAGCTGGCCGTGGATGCAGGGAGCCCA
TCCACCATCTTTAGCATGCCCTTGTATTCCCATCAGATGCCAGGATGAGGGGCATCAGAGAGTCCAAGTGTATGCCAA
ACCCAAACACCTAGGACTTGTCTTCTGGGACAGACAGATGCAGGAGAGCTAGGTTGGGCTGTGATCCCATTTACCACA
AAGAGGGAAAAACAAAAACAACAAACAAAAAACAACAAAAAACAACAAAAAACCAGGTCCAAATTGTA
GGTCAGGTTAGAGTTTATTTATGGAAGTTATATTCATCTCCATGGGGTCTACAAGGCTGGCGCCCATCAGAAAGAACA
AACAAAGGCTGATCTGGGAGGGGTGACTCTATGGCAGGAGCACGTGTCTTGGGGTACAGCCAGACACGGGGCTTG
TATTAATCAGGGCTTGTATTAATAGGCTGAGAGTCAAGCAGACAGAGAGACAGAAGGAAACACACACACACACACA
CACACACACACACACACATGCACACACCCTCACTTCTCACTCGAAGAGCCCCCTACTTACATTCTAAGAACAACCC
ATTCCTCTCTATAAGAGAGACAAAGTTGCAGAAACCCAAAGAGCCACAGGGTCCCCACTCTCTTTGAAATGACTTGGAC
TTGTTGCAGGGAAGACAGAGGGGTCTGCAGAGGCTTCTGGGTGACCCAGAGCCACAGACACTGAAATCTGGTGTGAGA
CCTGTATAAACCTCTTCCACAGGTTCCCTGAAAGGAGCCACATTCCTCAACCTGTCTCTGACCACTGAGGATGAGA
GCACTTGGGCTTCCCCATCTTGGAGTGCACCTGGTTTTCCCATCTGAGGGCACATGAGTCTCAGGTTCTGGGAAAG
TTCCACAAGTATTGAAAGTGTCTTGTGTTTGTGTGATTAAATTTAGGTGTATGAGTGTCTTGTGTTGAATATATGCCT
GTGTAGCATTTACAAGCCTGGTGCCTGAGGAGATCAGAAGATGGCATCAGATACCTTGAACCTGGACTTGCAGACAGTTA
TGAGCCACTGTGTGGGTGCTAGGAACAGAACCTGGATCTCCGGAAGAGCAGACAGCCAGCGCTCTTAGCCACTAAGCCA

[illegible]

-continued

TTCATGTATCTGAAAAGCGACACAACCATTTTCACATCATGGCATCTTCCTAACCCCATTCCTTTTGTGTTTGT
 TTGAGACAGGGTTTCTCTGTGTAGTCTGGCTGTCTGGAACCTCACTTTGTAGACCAGGCTGGCCTCGAACTCAGAAATC
 CTGGGATTAAAGGTGTGTGCCACCACGCCCGGCCCTAACCCCATTCCTTAATGGTGATCCAGTGGTTGAAATTCGGGCC
 ACACACATGTCCATTAGGGATTAGCTGTGTCTTCTGAGCTACCTGGTACAATCTTTATCCCTGGGGCTGGGCTCCTG
 ATCCCTGACTCGGGCCGATCAAGTCCAGTTCCTGGGCCGATCAAGTCCAGTTCCTGGGCCGAAACAGTCCAGTCCCT
 AGCTCGATTAGCTCATCTGGCTCCCTGGCCTGTCTTACTTACACTCTTCCCTTGCTCTGGACTGTGTGCTTTCTTTA
 CTCAGTTGTCTGCCACAGTCCCTAAGCCACCTCTGTAAGACAACCTAAGATAATACTTCCCTCAAGCACGGAAGTCTCTG
 AGTCACCACACCCTCTGGAGGTGTGTGGACACATGTTTCATGCGTGTGGTTGCGCTTACGTACGTGTGC

Sequence ID No. 18: Human Beer Genomic Sequence (This gene has two
 exons, at
 positions 161-427 and 3186-5219).

tagaggagaa gtccttgggg aggtttgct ctgagcacac cctttccct cctccgggg	60
ctgagggaac catgggacca gccctgcccc agcctgtcct cattggctgg catgaagcag	120
agaggggctt taaaaggcg accgtgtctc ggctggagac cagagcctgt gctactgga	180
ggtggcgtgc cctcctctgg ctggtaccat gcagctccca ctggccctgt gtctcgtctg	240
cctgctggtc cacacagcct tccgtgtagt ggagggccag ggggtggcagg cgttcaagaa	300
tgatgccacg gaaatcatcc ccgagctcgg agagtacccc gagcctccac cggagctgga	360
gaacaacaag accatgaacc gggcggagaa cggagggcgg cctccccacc accccttga	420
gaccaaaagg atgggttgga ggagagaatt cttagtaaaa gatcctgggg aggttttaga	480
aacttctctt tgggaggcct ggaagactgg ggtagaccca gtgaagattg ctggcctctg	540
ccagcactgg tcgaggaaca gtcttgcttg gaggtggggg aagaatggct cgtggtgca	600
gccttcaaat tcaggtgcag aggcctgagg caacagacgc tggtagagag ccagggcagg	660
gaggacgctg ggggtgtgag ggtatggcat cagggcatca gaacaggctc aggggctcag	720
aaaagaaaag gtttcaaga atctcctcct ggaatatag gagccacgtc cagctgctgg	780
taccactggg aagggaaaca ggtaaggagg cctcccatcc acagaacagc acctgtgggg	840
caccggacac tctatgtcgg tgggtgctgt ccccaaccaca cagaccaca tcatggaatc	900
cccaggaggt gaacccccag ctgaaagggg aagaaacagg ttccaggcac tcagtaactt	960
ggtagtgaag agagctgagg tgtgaacctg gtttgatcca actgcaagat agccctggtg	1020
tgtggggggg tgtgggggac agatctccac aaagcagtg ggaggaaggc cagagaggca	1080
ccctgcagtg gtgcattgcc catggcctgc ccagggagct ggcacttgaa ggaatgggag	1140
ttttcgacac agtttttagc cctgacatgg gtgcagctga gtccaggccc tggaggggag	1200
agcagcatcc tctgtgcagg agtagggaca tctgtcctca gcagccacc cagtcccaac	1260
cttgccctcat tccaggggag ggagaaggaa gaggaaccct gggttcctgg tcaggcctgc	1320
cacgagaagc ccaggtgaca gtgtgcatct ggtctataa ttggcaggaa tcctgaggcc	1380
atggggggcgt ctgaaatgac acttcagact aagagcttcc ctgtcctctg gccattatcc	1440
aggtggcaga gaagtccact gccaggctc ctggacccca gccctccccg cctcacaacc	1500
tgttgggact atgggtgtgt aaaaagggca actgcatggg aggccagcca ggaccctccg	1560
tcttcaaaat ggaggacaag ggcgcctccc cccacagctc cctttctagg caaggtcagc	1620
tgggctccag cgactgcctg aagggctgta aggaacccaa acacaaaatg tccaccttgc	1680
tggactccca cgagaggcca cagcccctga ggaagccaca tgctcaaac aaagtcatga	1740
tctgcagagg aagtgcctgg cctagggggc ctattctcga aaagccgcaa aatgccccct	1800

-continued

tccctgggca aatgcccc tgaccacaca cacattccag cctgcagag gtgaggatgc	1860
aaaccagccc acagaccaga aagcagcccc agacgatggc agtggccaca tctccctgc	1920
tgtgcttgct cttcagagtg ggggtggggg gtggccttct ctgtcccctc tctggtttgg	1980
tcttaagact attttccatt ctttcttgtc acattggaac tatccccatg aaacctttgg	2040
gggtggactg gtactcacac gacgaccagc tatttaaaaa gctccccacc atctaagtcc	2100
caactaggag acatgggtcaa ggtgtgtgca ggggatcagg ccaggcctcg gagcccaatc	2160
tctgcctgcc cagggagtat caccatgagg cgcccattca gataacacag aacaagaaat	2220
gtgcccagca gagagccagg tcaatgtttg tggcagctga acctgtagggt tttgggtcag	2280
agctcagggc ccctatggta ggaagtaac gacagtaaaa agcagccctc agctccatcc	2340
cccagcccag cctcccatgg atgctcgaaac gcagagcctc cactcttgcc ggagccaaaa	2400
ggtgctggga cccaggggaa gtggagtccg gagatgcagc ccagcctttt gggcaagtcc	2460
ttttctctgg ctgggcctca gtattctcat tgataatgag ggggttggaac aactgcctt	2520
tgattccttt caagtctaata gaattcctgt cctgatcacc tccccttcag tccctcgctt	2580
ccacagcagc tgccctgatt tattaccctc aattaacctc tactccttcc tccatccctt	2640
gtccaccctt cccaagtggc tggaaaagga atttgggaga agccagagcc aggcagaagg	2700
tgtgctgagt acttaacctg cccaggccag ggaccctgag gcacaagtgt ggettaaatc	2760
ataagaagac cccagaagag aaatgataat aataatacat aacagccgac gctttcagct	2820
atatgtgcca aatgggtattt tctgcattgc gtgtgtaatg gattaactcg caatgcttgg	2880
ggcgcccatc ttgcagaca ggaagaagag agaggttaag gaacttgccc aagatgacac	2940
ctgcagttag cgatggagcc ctggtgtttg aaccccagca gtcatttggc tccgagggga	3000
cagggtgccg agggagagctt tccaccagct ctagagcacc tgggaccttc ctgcaataga	3060
tgttcagggg caaaagcctc tggagacagg cttggcaaaa gcagggctgg ggtggagaga	3120
gacgggcccg tccagggcag gggtagccag gcgggcccgc accctcacgc gcgcctctct	3180
ccacagagct gtccgagtac agctgcccgc agctgcactt caccgcctac gtgaccgatg	3240
ggcctgtgcc cagcgccaag ccgggtaccg agctgggtgtg ctccggccag tgcggcccgc	3300
cgccctgctt gcccacgccc atcgggccgc gcaagtgggt gcgacctagt gggcccgaat	3360
tccgctgcat ccccgaccgc taccgcgcgc agcgcgtgca gctgctgtgt cccggtgggt	3420
aggcgccgcg cgcgcgcaag gtgcgcctgg tggcctcgtg caagtgcag gcctcacc	3480
gcttccacaa ccagtcggag ctcaaggact tcgggaccga ggcgctcgg ccgcagaagg	3540
gcccgaagcc gcggccccgc gcccgagcgc ccaagccaa ccaggccgag ctggagaacg	3600
cctaetagag ccgcgccgcg cccctcccca ccggcgggcg ccccgccctt gaaccgcgc	3660
cccacatttc tgcctctgc gcgtgggttg attgtttata ttctattgta aatgcctgca	3720
acccagggca gggggctgag accttccagg ccctgaggaa tcccgggcgc cggcaaggcc	3780
cccctcagcc cgcagctga ggggtccac ggggcagggg aggggaattga ggtcacaga	3840
caactgagca cgcagcccg cctctggggc cgctacctt tgctggtccc acttcagagg	3900
aggcagaat ggaagcattt tcaccgccct ggggttttaa gggagcgggt tgggagtggg	3960
aaagtccagg gactggttaa gaaagtggga taagattccc ccttgacctt cgctgccat	4020
cagaaagcct gaggcgtgcc cagagcaca gactgggggc aactgtagat gtggtttcta	4080
gtcctggctc tgccactaac ttgctgtgta accttgaaat acacaattct ccttcgggac	4140
ctcaatttcc actttgtaaa atgagggtgg aggtgggaat aggatctcga ggagactatt	4200

-continued

ggcatatgat tccaaggact ccagtgcctt ttgaatgggc agaggtgaga gagagagaga	4260
gaaagagaga gaatgaatgc agttgcattg attcagtgcc aaggtcactt ccagaattca	4320
gagttgtgat gctctcttct gacagccaaa agtgaaaaac aaacagaaaa aaaaaagtaa	4380
agagtctatt tatggctgac atatttacgg ctgacaaact cctggaagaa gctatgctgc	4440
ttcccagcct ggcttccccg gatgtttggc tacctccacc cctccatctc aaagaaataa	4500
catcatccat tggggtagaa aaggagaggg tccgaggggtg gtgggagggg tagaaatcac	4560
atccgcccc aactcccaaa gagcagcatc cctccccga cccatagcca tgttttaaag	4620
tcaccttccg aagagaagtg aaagggtcaa ggacactggc cttgcaggcc cgaggggagca	4680
gccatcacaa actcacagac cagcacatcc cttttgagac accgccttct gccaccact	4740
cacggacaca ttctgccta gaaaacagct tcttactgct ctacatgtg atggcatatc	4800
ttacactaaa agaataattat tgggggaaaa actacaagtg ctgtacatat gctgagaaac	4860
tgacagagcat aatagctgac acccaaaaat ctttttgaaa atcatttcca gacaacctct	4920
tactttctgt gtagttttta attgttaaaa aaaaaaagtt ttaaacagaa gcacatgaca	4980
tatgaaagcc tgacaggactg gtcgtttttt tggcaattct tccacgtggg acttgctcac	5040
aagaatgaaa gtagtggttt ttaaaagatt aagttacata tttattttct cacttaagtt	5100
atttatgcaa aagtttttct tgtagagaat gacaatgtta atattgcttt atgaattaac	5160
agtcgttctc tccagagtcc agagacattg ttaataaaga caatgaatca tgaccgaaag	5220
gatgtggtct cattttgtca accacacatg acgtcatttc tgtcaaagtt gacacccttc	5280
tcttggtcac tagagctcca accttgagca cacttttgac tgctctctgg tggcccttgt	5340
ggcaattatg tcttcttttg aaaagtcatg tttatccctt cctttccaaa cccagaccgc	5400
atttcttcc caggggcatg gtaataacct cagccttgta tcccttttagc agcctccct	5460
ccatgctggc ttccaaaatg ctgttctcat tgtatcactc ccctgctcaa aagccttcca	5520
tagctcccc ttgccagga tcaagtgcag tttccctatc tgacatggga ggccttctct	5580
gcttgactcc cactccccc tccaccaagc ttcctactga ctccaaatgg tcatgcagat	5640
ccctgcttcc ttagtttgcc atccacactt agcaccacca ataactaatc ctctttcttt	5700
aggattcaca ttacttgtca tctcttcccc taaccttcca gagatgttcc aatctcccat	5760
gatccctctc tctctgagg ttccagcccc ttttgtctac accactactt tggttcctaa	5820
ttctgttttc catttgacag tcattcatgg aggaccagcc tggccaagtc ctgcttagta	5880
ctggcataga caacacaaag ccaagtacaa ttcaggacca gctcacagga aacttcactc	5940
tcttcgaagt gtggatttga tgcctcctgg gtagaaatgt aggatcttca aaagtgggcc	6000
agcctcctgc acttctctca aagtctcgcc tccccaaggt gtcttaatag tgctggatgc	6060
tagctgagtt agcatcttca gatgaagagt aaccctaaag ttactcttca gttgccctaa	6120
ggtgggatgg tcaactggaa agcttttaaa taagtccagc ctaccttggg ggaaccacc	6180
cccacaaaga aagctgaggt ccctcctgat gacttgtag tttaactacc aataaccac	6240
ttgaattaat catcatcatc aagtctttga taggtgtgag tgggtatcag tggccggtcc	6300
cttctgtggg ctccagcccc cgaggaggcc tcagttagcc cctgcagaaa atccatgcat	6360
catgagtgtc tcagggccca gaatatgaga gcaggtagga aacagagaca tcttccatcc	6420
ctgagaggca gtgcggtcca gtgggtgggg acacgggctc tgggtcaggt ttgtgtgtgt	6480
tgtttgtttg ttttgagaca gagtctcgct ctattgcccc ggctggagtg cagtgtcaca	6540
atctcggtct actgcaactt ctgccttccc ggattcaagt gattctcctg cctcagcctc	6600

-continued

cagagtagct gggattacag gtgctgcca	6660
agagacgggg ttccaccatg ttggccaggc tagtctcga ctcttgacct caagtgatct	6720
gctgctctcg gcctcccaaa gtgctgggat tacaggcgtg agccaccaca cccagcccca	6780
ggttggtgtt tgaatctgag gagactgaag caccaagggg ttaaagtgtt tgcccacagc	6840
catacttggg ctgagttcct tgccctaccc ctgacttgag ctgcttagaa cctggtgggc	6900
acatgggcaa taaccaggtc acaactgttt gtaccaagtg ttatgggaat ccaagatagg	6960
agtaatttgc tctgtggagg ggatgaggga tagtggttag ggaaagcttc acaaagtggg	7020
tggtgcttag agattttcca ggtggagaag ggggcttcta ggcagaaggc atagcccaag	7080
caaagactgc aagtgcattg ctgctcatgg gtagaagaga atccaccatt cctcaacatg	7140
taccgagtc ttgccatgtg caaggcaaca tgggggtacc aggaattcca agcaatgtcc	7200
aaacctaggg tctgctttct gggacctgaa gatacaggat ggatcagccc aggctgcaat	7260
cccattacca cgagggggaa aaaaacctga aggctaaatt gtaggctggg ttagaggtta	7320
tttatggaaa gttatatctt acctacatgg ggtctataag cctggcgcca atcagaaaaag	7380
gaacaaaaca cagacctagc tggaggggc agcattttgt ttagaggggc ggggcacatg	7440
ttctgggggt acagccagac tcaggggctt tattaatagt ctgagagtaa gacagacaga	7500
gggatagaa gaaataggtc ctttctctc tctctctctc tctctctctc actctctctc	7560
tctctcacac acacacacag acacacacac acgctctgta ggggtctact tatgctcaa	7620
gtacaaatca ggccacattt acacaaggag gtaaaggaaa agaactgttg aggagccaca	7680
ggaccccaaa attccctggt ttccttgat caggcaggac ttacgcagct gggaggggtg	7740
agagcctgca gaagccacct gcgagtaagc caagtccaga gtcacagaca ccaaagctg	7800
gtgccatgtc ccacaccgc ccacctcca ctgctcctt gacacagccc tgtgctccac	7860
aacccggctc ccagatcatt gattatagct ctggggcctg caccgtcctt cctgccacat	7920
ccccaccca ttcttggaac ctgccctctg tcttctcctt tgtccaaggg caggcaaggg	7980
ctcagctatt gggcagcttt gaccaacagc tgaggctcct tttgtggctg gagatgcagg	8040
aggcagggga atattcctct tagtcaatgc gaccatgtgc ctggtttgcc cagggtgggc	8100
tcgtttacac ctgtaggcca agcgtaat ttaacagctc ccacttctac tctaaaaaat	8160
gacccaatct gggcagtaaa ttatatggtg cccatgctat taagagctgc aacttgctgg	8220
gcgtggtggc tcacacctgt aatcccagta ctttgggacg tcaaggcggg tggatcacct	8280
gaggtcacga gttagagact ggcctggcca gcatggcaaa accccatctt tactaaaaat	8340
acaaaaatta gcaaggcatg gtggcatgca cctgtaatcc cagggtactg ggaggctgag	8400
acaggagaat ggcttgaaac caggaggcag aggttgagct gagccaagat tgtgccactg	8460
ccctccagcc ctggcaacag agcaagactt catctcaaaa gaaaaaggat actgtcaatc	8520
actgcaggaa gaacccaggt aatgaatgag gagaagagag gggctgagtc accatagtgg	8580
cagcaccgac tcctgcagga aaggcgagac actgggtcat gggactgaa ggggtccctg	8640
aatgacgttc tgcttttagag accgaacctg agccctgaaa gtgcatgcct gttcatgggt	8700
gagagactaa attcatcatt ccttggcagg tactgaatcc tttcttacgg ctgccctcca	8760
atgcccaatt tccctacaat tgtctgggtt gcctaagctt ctgccacca agagggccag	8820
agctggcagc gagcagctgc aggtaggaga gataggtagc cataaggag gtgggaaaga	8880
gagatggaag gagaggggtg cagagcacac acctccctg cctgacaact tcctgagggc	8940
tggtcatgcc agcagattta aggcggaggc aggggagatg gggcgggaga ggaagtgaat	9000

-continued

aaggagaggg tggggatgga gaggaagaga gggatgatcat tcattcattc cattgctact	9060
gactggatgc cagctgtgag ccaggcacca ccttagctct gggcatgtgg ttgtaatctt	9120
ggagcctcat ggagctcaca gggagtgtctg gcaaggagat ggataatgga cggataacaa	9180
ataaacattt agtacaatgt ccgggaatgg aaagtctctg aaagaaaaat aaagctgggtg	9240
agcatataga cagccctgaa ggcggccagg ccaggcattt ctgaggaggt ggcatttgag	9300
c	9301

From the foregoing, it will be appreciated that although specific embodiments of the invention have been described herein for purposes of illustration, various modifications

may be made without deviating from the spirit and scope of the invention. Accordingly, the invention is not limited except as by the appended claims.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 41

<210> SEQ ID NO 1

<211> LENGTH: 2301

<212> TYPE: DNA

<213> ORGANISM: Homo sapien

<400> SEQUENCE: 1

agagcctgtg ctactggaag gtggcgtgcc ctctctgtgc tggtagcatg cagctccac	60
tggccctgtg tctcgtctgc ctgctgttac acacagcctt ccgtgtagtg gagggccagg	120
ggtggcaggc gttcaagaat gatgccagg aaatcatccc cagctcggga gagtacccg	180
agcctccacc ggagctggag aacaacaaga ccatgaaccg ggcggagaac ggagggcggc	240
ctccccacca cccctttgag accaaagacg tgtccgagta cagctgccgc gactgcact	300
tcaccgcta cgtgaccgat gggccgtgcc gcagcgcaa gccggtcacc gactggtgt	360
gctccggcca gtggggcccg gcgcgcctgc tgcccaacgc catcgccgcg ggcaagtgt	420
ggcgacctag tggggcccgac ttccgctgca tccccgaccg ctaccgcgcg cagcgctgc	480
agctgctgtg tcccggtggt gaggcgccgc gcgcgcgcaa ggtgcgcctg gtggcctcgt	540
gcaagtgcga gcgcctcacc cgcttcaca accagtcgga gctcaaggac ttcgggaccg	600
aggccgctcg gccgcagaag ggcgggaagc cgcggccccc cgcccgagag gccaaagcca	660
accaggccga gctggagaac gcctactaga gcccgcgcgc gccctcccc accggcgggc	720
gccccggccc tgaaccgcgc ccccacattt ctgtcctctg cgcgtggttt gattgtttat	780
atttcattgt aaatgcctgc aaccagggc aggggggtga gaccttccag gccctgagga	840
atccccggcg ccggcaaggc cccctcagc ccgccagctg aggggtccca cggggcaggg	900
gagggaattg agagtcacag acactgagcc acgcagcccc gcctctgggg ccgectacct	960
ttgctggtcc cacttcagag gaggcagaaa tggaagcatt ttcaccgccc tgggggttta	1020
agggagcggg gtgggagtgg gaaagtccag ggactgggta agaaagtggg ataagattcc	1080
cccttgacc tcgctgcccc tcagaaagcc tgaggcgtgc ccagagcaca agactggggg	1140
caactgtaga tgtggtttct agtcctggct ctgccactaa cttgctgtgt aacctgaac	1200
tacacaattc tccttcggga cctcaatttc cactttgtaa aatgaggggt gagggtggaa	1260
taggatctcg aggagactat tggcatatga ttccaaggac tccagtgcct ttgtaatggg	1320
cagaggtgag agagagagag agaaagagag agaatgaatg cagttgcatt gattcagtc	1380
caaggtcact tccagaattc agagttgtga tgctctcttc tgacagccaa agatgaaaaa	1440

-continued

```

caaacagaaa aaaaaaagta aagagtctat ttatggctga catatttacg gctgacaaac 1500
tcctggaaga agctatgctg cttcccagcc tggcttcccc ggatgttttg ctacctccac 1560
ccctccatct caaagaaata acatcatcca ttggggtaga aaaggagagg gtccgagggt 1620
ggtgggaggg atagaaatca catccgcccc aacttcccaa agagcagcat ccctcccccg 1680
accatagcc atgttttaaa gtcaccttc gaagagaagt gaaagggtca aggacactgg 1740
ccttgaggc ccgaggaggc agccatcaca aactcacaga ccagcacatc ctttttgaga 1800
caccgccttc tgcccaccac tcacggacac atttctgcct agaaaacagc ttcttactgc 1860
tcttacatgt gatggcatat cttacactaa aagaatatta ttgggggaaa aactacaagt 1920
gctgtacata tgctgagaaa ctgcagagca taatagctgc caccacaaaa tctttttgaa 1980
aatcatttcc agacaacctc ttactttctg tgtagttttt aattgttaaa aaaaaaagt 2040
tttaaacaga agcacatgac atatgaaagc ctgcaggact ggtcgttttt ttggcaattc 2100
ttccacgtgg gacttgtcca caagaatgaa agtagtggtt tttaagagt taagtacat 2160
atttatttcc tcacttaagt tatttatgca aaagttttcc ttgtagagaa tgacaatgtt 2220
aatattgctt tatgaattaa cagtctgttc ttccagagtc cagagacatt gtaataaag 2280
acaatgaatc atgaccgaaa g 2301

```

```

<210> SEQ ID NO 2
<211> LENGTH: 213
<212> TYPE: PRT
<213> ORGANISM: Homo sapien

```

```

<400> SEQUENCE: 2

```

```

Met Gln Leu Pro Leu Ala Leu Cys Leu Val Cys Leu Leu Val His Thr
1           5           10           15

Ala Phe Arg Val Val Glu Gly Gln Gly Trp Gln Ala Phe Lys Asn Asp
20           25           30

Ala Thr Glu Ile Ile Pro Glu Leu Gly Glu Tyr Pro Glu Pro Pro Pro
35           40           45

Glu Leu Glu Asn Asn Lys Thr Met Asn Arg Ala Glu Asn Gly Gly Arg
50           55           60

Pro Pro His His Pro Phe Glu Thr Lys Asp Val Ser Glu Tyr Ser Cys
65           70           75           80

Arg Glu Leu His Phe Thr Arg Tyr Val Thr Asp Gly Pro Cys Arg Ser
85           90           95

Ala Lys Pro Val Thr Glu Leu Val Cys Ser Gly Gln Cys Gly Pro Ala
100          105          110

Arg Leu Leu Pro Asn Ala Ile Gly Arg Gly Lys Trp Trp Arg Pro Ser
115          120          125

Gly Pro Asp Phe Arg Cys Ile Pro Asp Arg Tyr Arg Ala Gln Arg Val
130          135          140

Gln Leu Leu Cys Pro Gly Gly Glu Ala Pro Arg Ala Arg Lys Val Arg
145          150          155          160

Leu Val Ala Ser Cys Lys Cys Lys Arg Leu Thr Arg Phe His Asn Gln
165          170          175

Ser Glu Leu Lys Asp Phe Gly Thr Glu Ala Ala Arg Pro Gln Lys Gly
180          185          190

```


-continued

Arg Lys Pro Arg Pro Arg Ala Arg Ser Ala Lys Ala Asn Gln Ala Glu
 195 200 205
 Leu Glu Asn Ala Tyr
 210

<210> SEQ ID NO 3
 <211> LENGTH: 2301
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapien

<400> SEQUENCE: 3

```

agagcctgtg ctactggaag gtggcgtgcc ctctcttgge tggtagcatg cagctccac 60
tggccctgtg tctcgtctgc ctgctggtac acacagcctt ccgtgtagtg gagggctagg 120
ggtggcaggc gttcaagaat gatgccacgy aaatcatccc cgagctcgga gagtaccccg 180
agcctccacc ggagctggag aacaacaaga ccatgaaccg ggcggagaac ggagggcggc 240
ctccccacca ccccttttag accaaagacg tgtccgagta cagctgccgc gagctgcact 300
tcacccgcta cgtgaccgat gggccgtgcc gcagcgccaa gccggtcacc gagctggtgt 360
gctccggcca gtgcggcccg gcgcgcctgc tgcccaacgc catcggccgc ggcaagtgtt 420
ggcgacctag tgggcccgcac ttccgctgca tcccgcaccg ctaccgcgcg cagcgctgac 480
agctgctgtg tcccggttgt gagcgccgc gcgcgcgcaa ggtgcgcctg gtggcctcgt 540
gcaagtgcac gcgcctcacc cgcttcacca accagtcgga gctcaaggac ttccgggaccg 600
aggccgctcg gccgcagaag ggcgcgaagc gcgcggcccg cgcgcggagc gccaaagcca 660
accaggccga gctggagaac gcctactaga gccgcgccgc gccctcctcc accggcgggc 720
gccccggccc tgaaccgcgc cccacattt ctgtcctctg cgcgtggttt gattgtttat 780
atttcattgt aaatgcctgc aaccaggggc agggggctga gaccttcag gccctgagga 840
atcccgggcg ccggcaaggc cccctcagc ccgccagctg aggggtccca cggggcaggg 900
gagggaattg agagtacag acactgagcc acgcagcccc gcctctgggg ccgcctacct 960
ttgctgtgcc cacttcagag gaggcagaaa tggaagcatt ttcaccgcc tggggtttta 1020
agggagcggg gtgggagtgg gaaagtccag ggactggtta agaaagtgg ataagattcc 1080
cccttgacac tcgtgcccc tcagaaagcc tgaggcgtgc ccagagcaca agactggggg 1140
caactgtaga tgtggtttct agtcctggct ctgccaactaa cttgctgtgt aacctgaac 1200
tacacaattc tccttcggga cctcaatttc cactttgtaa aatgagggtg gaggtgggaa 1260
taggatctcg aggagactat tggcatatga ttccaaggac tccagtgcct ttggaatggg 1320
cagaggtgag agagagagag agaaagagag agaataatg cagttgcatt gattcagtgc 1380
caaggtcact tccagaattc agagtgtgta tgctctcttc tgacagccaa agatgaaaaa 1440
caaacagaaa aaaaaagta aagagtctat ttatggctga catatttacg gctgacaaac 1500
tcctggaaga agctatgctg cttcccagcc tggcttcccc ggatgttttg ctacctccac 1560
ccctccatct caaagaata acatcatcca ttggggtaga aaaggagagg gtccgagggt 1620
ggtggggagg atagaaatca catccgcccc aacttcccaa agagcagcat ccctcccccg 1680
accatagacc atgttttaaa gtcaccttcc gaagagaagt gaaaggttca aggacactgg 1740
ccttgacagg ccgagggagc agccatcaca aactcacaga ccagcacatc ccttttgaga 1800
caccgccttc tgcccaccac tcacggacac atttctgcct agaaaacagc ttcttactgc 1860
tcttacatgt gatggcatat cttacactaa aagaatatta ttgggggaaa aactacaagt 1920

```

-continued

gctgtacata tgctgagaaa ctgcagagca taatagctgc caccacaaaa tctttttgaa	1980
aatcattttcc agacaacctc ttactttctg tgtagttttt aattgttaaa aaaaaaagt	2040
tttaaacaga agccatgac atatgaaagc ctgcaggact ggctgttttt ttggcaattc	2100
ttccacgtgg gacttgtcca caagaatgaa agtagtggtt tttaaagagt taagttacat	2160
atttattttc tcacttaagt tatttatgca aaagtttttc ttgtagagaa tgacaatgtt	2220
aattattgctt tatgaattaa cagtctgttc ttccagagtc cagagacatt gttaataaag	2280
acaatgaatc atgaccgaaa g	2301

<210> SEQ ID NO 4
 <211> LENGTH: 23
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapien

<400> SEQUENCE: 4

Met	Gln	Leu	Pro	Leu	Ala	Leu	Cys	Leu	Val	Cys	Leu	Leu	Val	His	Thr
1			5						10					15	

Ala	Phe	Arg	Val	Val	Glu	Gly
			20			

<210> SEQ ID NO 5
 <211> LENGTH: 2301
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapien

<400> SEQUENCE: 5

agagcctgtg ctactggaag gtggcgtgcc ctccctctggc tggtagcatg cagctccac	60
tggccctgtg tctcatctgc ctgctggtac acacagcctt ccgtgtagtg gagggccagg	120
ggtggcaggc gttcaagaat gatgccacgg aaatcatccg cgagctcgga gagtaccocg	180
agcctccacc ggagctggag aacaacaaga ccatgaaccg ggcggagaac ggagggcggc	240
ctccccacca ccccttttag accaaagacg tgtccgagta cagctgccgc gagctgcact	300
tcaccgccta cgtgaccgat gggccgtgcc gcagcgccaa gccggtcacc gagctggtgt	360
gtcccgccca gtgcggcccg gcgcgcctgc tgcccaacgc catcggccgc ggcaagtgtt	420
ggcgacctag tggggccgac ttccgctgca tccccgaccg ctaccgcgcg cagcgctgc	480
agctgctgtg tcccggtgtt gaggcgccgc gcgcgcgcaa ggtgcgcctg gtggcctcgt	540
gcaagtgcaa gcgcctcacc cgcttcaca accagtcgga gctcaaggac ttcgggaccg	600
aggccgctcg gccgcagaag ggcgggaagc cgcggccccc cgcgggagc gccaaagcca	660
accaggccga gctggagaac gcctactaga gcccgccccg gcccctcccc accggcgggc	720
gccccggccc tgaaccgcgc cccacattt ctgtcctctg cgcgtggttt gattgtttat	780
atttcattgt aaatgcctgc aaccagggc agggggctga gacctccag gccctgaggc	840
atcccgggcg ccggcaaggc cccctcagc ccgccaagctg aggggtccca cggggcaggc	900
gaggggaattg agagtcacag aactgagcc acgcagcccc gcctctgggg ccgcctacct	960
ttgctgggtc cacttcagag gaggcagaaa tggaagcatt ttcaccgcc ttgggtttta	1020
agggagcggg gtgggagtg gaaagtccag ggactggta agaaagtgg ataagattcc	1080
cccttgacac tcgctgcccc tcagaaagcc tgaggcgtgc ccagagcaca agactggggg	1140
caactgtaga tgtgttttct agtccctgct ctgccaactaa cttgctgtgt aacctgaac	1200
tacacaattc tccttcggga cctcaatttc cactttgtaa aatgagggtg gaggtgggaa	1260
taggatctcg aggagactat tggcatatga ttccaaggac tccagtgcct tttgaatggg	1320

-continued

```

cagagggtgag agagagagag agaaagagag agaataatg cagttgcatt gattcagtgc 1380
caaggctact tccagaattc agagttgtga tgctctcttc tgacagccaa agatgaaaaa 1440
caaacagaaa aaaaaaagta aagagtctat ttatggctga catatttacg gctgacaaac 1500
tcctggaaga agctatgtg cttccagcc tggcttcccc ggatgtttgg ctacctccac 1560
ccctccatct caaagaaata acatcatcca ttggggtaga aaaggagagg gtccgagggg 1620
ggtgggaggg atagaaatca catccgcccc aacttcccaa agagcagcat ccotcccccg 1680
acccatagcc atgttttaaa gtcaccttc gaagagaagt gaaagggtca aggacactgg 1740
ccttgagggc ccgagggagc agccatcaca aactcacaga ccagcacatc ccttttgaga 1800
caccgccttc tgcccaccac tcacggacac atttctgcct agaaaacagc ttcttactgc 1860
tcttacatgt gatggcatat cttacactaa aagaatatta ttgggggaaa aactacaagt 1920
gctgtacata tgctgagaaa ctgcagagca taatagctgc caccacaaaa tctttttgaa 1980
aatcatttcc agacaacctc ttactttctg tgtagttttt aattgtttaa aaaaaaagt 2040
tttaaacaga agcacatgac atatgaaagc ctgcaggact ggtcgttttt ttggcaattc 2100
ttccacgtgg gacttgtcca caagaatgaa agtagtggtt tttaaagagt taagtacat 2160
atatttttct tcaacttaagt tatttatgca aaagtttttc ttgtagagaa tgacaatgtt 2220
aatattgctt tatgaattaa cagtctgttc ttccagagtc cagagacatt gtaataaag 2280
acaatgaatc atgaccgaaa g 2301

```

<210> SEQ ID NO 6

<211> LENGTH: 213

<212> TYPE: PRT

<213> ORGANISM: Homo sapien

<400> SEQUENCE: 6

```

Met Gln Leu Pro Leu Ala Leu Cys Leu Ile Cys Leu Leu Val His Thr
1           5           10           15
Ala Phe Arg Val Val Glu Gly Gln Gly Trp Gln Ala Phe Lys Asn Asp
20           25           30
Ala Thr Glu Ile Ile Arg Glu Leu Gly Glu Tyr Pro Glu Pro Pro Pro
35           40           45
Glu Leu Glu Asn Asn Lys Thr Met Asn Arg Ala Glu Asn Gly Gly Arg
50           55           60
Pro Pro His His Pro Phe Glu Thr Lys Asp Val Ser Glu Tyr Ser Cys
65           70           75           80
Arg Glu Leu His Phe Thr Arg Tyr Val Thr Asp Gly Pro Cys Arg Ser
85           90           95
Ala Lys Pro Val Thr Glu Leu Val Cys Ser Gly Gln Cys Gly Pro Ala
100          105          110
Arg Leu Leu Pro Asn Ala Ile Gly Arg Gly Lys Trp Trp Arg Pro Ser
115          120          125
Gly Pro Asp Phe Arg Cys Ile Pro Asp Arg Tyr Arg Ala Gln Arg Val
130          135          140
Gln Leu Leu Cys Pro Gly Gly Glu Ala Pro Arg Ala Arg Lys Val Arg
145          150          155          160
Leu Val Ala Ser Cys Lys Cys Lys Arg Leu Thr Arg Phe His Asn Gln
165          170          175
Ser Glu Leu Lys Asp Phe Gly Thr Glu Ala Ala Arg Pro Gln Lys Gly
180          185          190

```


-continued

```

aatcatttcc agacaacctc ttactttctg tgtagttttt aattgtttaa aaaaaaaagt 2040
tttaaacaga agcacatgac atatgaaagc ctgcaggact ggtcgttttt ttggcaattc 2100
ttccacgtgg gacttgtcca caagaatgaa agtagtggtt tttaagagat taagttacat 2160
atttattttc tcacttaagt tatttatgca aaagtttttc ttgtagagaa tgacaatgtt 2220
aatattgctt tatgaattaa cagtctgttc ttccagagtc cagagacatt gttaataaag 2280
acaatgaatc atgaccgaaa g 2301

```

<210> SEQ ID NO 8
 <211> LENGTH: 213
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapien

<400> SEQUENCE: 8

```

Met Gln Leu Pro Leu Ala Leu Cys Leu Val Cys Leu Leu Val His Thr
1      5      10      15
Ala Phe Arg Val Val Glu Gly Gln Gly Trp Gln Ala Phe Lys Asn Asp
20     25     30
Ala Thr Glu Ile Ile Arg Glu Leu Gly Glu Tyr Pro Glu Pro Pro Pro
35     40     45
Glu Leu Glu Asn Asn Lys Thr Met Asn Arg Ala Glu Asn Gly Gly Arg
50     55     60
Pro Pro His His Pro Phe Glu Thr Lys Asp Val Ser Glu Tyr Ser Cys
65     70     75     80
Arg Glu Leu His Phe Thr Arg Tyr Val Thr Asp Gly Pro Cys Arg Ser
85     90     95
Ala Lys Pro Val Thr Glu Leu Val Cys Ser Gly Gln Cys Gly Pro Ala
100    105    110
Arg Leu Leu Pro Asn Ala Ile Gly Arg Gly Lys Trp Trp Arg Pro Ser
115    120    125
Gly Pro Asp Phe Arg Cys Ile Pro Asp Arg Tyr Arg Ala Gln Arg Val
130    135    140
Gln Leu Leu Cys Pro Gly Gly Glu Ala Pro Arg Ala Arg Lys Val Arg
145    150    155    160
Leu Val Ala Ser Cys Lys Cys Lys Arg Leu Thr Arg Phe His Asn Gln
165    170    175
Ser Glu Leu Lys Asp Phe Gly Thr Glu Ala Ala Arg Pro Gln Lys Gly
180    185    190
Arg Lys Pro Arg Pro Arg Ala Arg Ser Ala Lys Ala Asn Gln Ala Glu
195    200    205
Leu Glu Asn Ala Tyr
210

```

<210> SEQ ID NO 9
 <211> LENGTH: 642
 <212> TYPE: DNA
 <213> ORGANISM: Cercopithecus pygerythrus

<400> SEQUENCE: 9

```

atgcagctcc cactggccct gtgtcttgtc tgctgtgtgg tacacgcagc ctccgtgta 60
gtggaggggc aggggtggca ggccttcaag aatgatgcc aaggaaatcat ccccgagctc 120
ggagagtacc ccgagcctcc accggagctg gagaacaaca agaccatgaa ccgggcggag 180
aatggagggc ggcctcccca ccacccctt gagaccaaag acgtgtccga gtacagctgc 240

```

-continued

cgagagctgc acttcacccg ctacgtgacc gatgggcccgt gccgcagcgc caagccagtc	300
accgagttgg tgtgtcccg ccagtgcggc ccggcacgccc tgetgcccaa cgccatcggc	360
cgcggcaagt ggtggcgccc gagtggccc gacttccgct gcatcccca ccgtaccgc	420
gcgcagcgtg tgcagtgct gtgtcccggt ggtgccgcgc cgcgcgcgcg caaggtgcgc	480
ctgggtggcct cgtgcaagt caagcgctc acccgcttc acaaccagtc ggagctcaag	540
gacttcggtc ccgagggccc tcggcccgag aagggccgga agccgcggcc ccgcgcccgg	600
ggggccaaag ccaatcaggc cgagctggag aacgcctact ag	642

<210> SEQ ID NO 10
 <211> LENGTH: 213
 <212> TYPE: PRT
 <213> ORGANISM: Cercopithecus pygerythrus

<400> SEQUENCE: 10

Met Gln Leu Pro Leu Ala Leu Cys Leu Val Cys Leu Leu Val His Ala	
1 5 10 15	
Ala Phe Arg Val Val Glu Gly Gln Gly Trp Gln Ala Phe Lys Asn Asp	
20 25 30	
Ala Thr Glu Ile Ile Pro Glu Leu Gly Glu Tyr Pro Glu Pro Pro Pro	
35 40 45	
Glu Leu Glu Asn Asn Lys Thr Met Asn Arg Ala Glu Asn Gly Gly Arg	
50 55 60	
Pro Pro His His Pro Phe Glu Thr Lys Asp Val Ser Glu Tyr Ser Cys	
65 70 75 80	
Arg Glu Leu His Phe Thr Arg Tyr Val Thr Asp Gly Pro Cys Arg Ser	
85 90 95	
Ala Lys Pro Val Thr Glu Leu Val Cys Ser Gly Gln Cys Gly Pro Ala	
100 105 110	
Arg Leu Leu Pro Asn Ala Ile Gly Arg Gly Lys Trp Trp Arg Pro Ser	
115 120 125	
Gly Pro Asp Phe Arg Cys Ile Pro Asp Arg Tyr Arg Ala Gln Arg Val	
130 135 140	
Gln Leu Leu Cys Pro Gly Gly Ala Ala Pro Arg Ala Arg Lys Val Arg	
145 150 155 160	
Leu Val Ala Ser Cys Lys Cys Lys Arg Leu Thr Arg Phe His Asn Gln	
165 170 175	
Ser Glu Leu Lys Asp Phe Gly Pro Glu Ala Ala Arg Pro Gln Lys Gly	
180 185 190	
Arg Lys Pro Arg Pro Arg Ala Arg Gly Ala Lys Ala Asn Gln Ala Glu	
195 200 205	
Leu Glu Asn Ala Tyr	
210	

<210> SEQ ID NO 11
 <211> LENGTH: 638
 <212> TYPE: DNA
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 11

atgcagccct cactagcccc gtgcctcacc tgcctacttg tgcacgtgc cttctgtgct	60
gtggaggggc aggggtggca agccttcagg aatgatgcca cagaggtcat cccagggctt	120
ggagagtacc ccgagcctcc tcctgagAAC aaccagacca tgaaccgggc ggagaatgga	180
ggcagacctc cccaccatcc ctatgacgcc aaaggtgtgt ccgagtacag ctgccgcgag	240

-continued

```

ctgcactaca cccgcttcct gacagacggc ccatgccgca gcgccaagcc ggtcaccgag    300
ttggtgtgct cgggccagtg cggccccgcy cggctgctgc ccaacgccat cgggcgcgtg    360
aagtgtgtgg gcccgaaagg accggatttc cgtgcatcc cggatcgcta ccgcgcgcag    420
cgggtgcagc tgctgtgccc cggggggcgc gcgccgcgct cgcgcaaggt gcgtctgttg    480
gcctcgtgca agtgcaagcg cctcaccgcg ttccacaacc agtcggagct caaggacttc    540
gggccggaga ccgcgcgggc gcagaagggt cgcaagccgc ggcccggcgc ccggggagcc    600
aaagccaacc aggcggagct ggagaacgcc tactagag                                638

```

<210> SEQ ID NO 12
 <211> LENGTH: 211
 <212> TYPE: PRT
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 12

```

Met Gln Pro Ser Leu Ala Pro Cys Leu Ile Cys Leu Leu Val His Ala
1          5          10          15
Ala Phe Cys Ala Val Glu Gly Gln Gly Trp Gln Ala Phe Arg Asn Asp
20          25          30
Ala Thr Glu Val Ile Pro Gly Leu Gly Glu Tyr Pro Glu Pro Pro Pro
35          40          45
Glu Asn Asn Gln Thr Met Asn Arg Ala Glu Asn Gly Gly Arg Pro Pro
50          55          60
His His Pro Tyr Asp Ala Lys Asp Val Ser Glu Tyr Ser Cys Arg Glu
65          70          75          80
Leu His Tyr Thr Arg Phe Leu Thr Asp Gly Pro Cys Arg Ser Ala Lys
85          90          95
Pro Val Thr Glu Leu Val Cys Ser Gly Gln Cys Gly Pro Ala Arg Leu
100         105         110
Leu Pro Asn Ala Ile Gly Arg Val Lys Trp Trp Arg Pro Asn Gly Pro
115         120         125
Asp Phe Arg Cys Ile Pro Asp Arg Tyr Arg Ala Gln Arg Val Gln Leu
130         135         140
Leu Cys Pro Gly Gly Ala Ala Pro Arg Ser Arg Lys Val Arg Leu Val
145         150         155         160
Ala Ser Cys Lys Cys Lys Arg Leu Thr Arg Phe His Asn Gln Ser Glu
165         170         175
Leu Lys Asp Phe Gly Pro Glu Thr Ala Arg Pro Gln Lys Gly Arg Lys
180         185         190
Pro Arg Pro Gly Ala Arg Gly Ala Lys Ala Asn Gln Ala Glu Leu Glu
195         200         205
Asn Ala Tyr
210

```

<210> SEQ ID NO 13
 <211> LENGTH: 674
 <212> TYPE: DNA
 <213> ORGANISM: Rattus norvegicus

<400> SEQUENCE: 13

```

gaggaccgag tgcccttcct cctctggca ccatgcagct ctactagcc ccttgcttg    60
cctgcctgct tgtacatgca gccttcgttg ctgtggagag ccagggttg caagccttca    120
agaatgatgc cacagaaatc atccggggac tcagagagta cccagagcct cctcaggaac    180

```

-continued

tagagaacaa ccagaccatg aaccgggccg agaacggagg cagacccccc caccatcctt	240
atgacaccaa agacgtgtcc gactacagct gccgcgagct gcactacacc cgcttcgtga	300
ccgacggccc gtgccgcagt gccaaagccg tcaccgagtt ggtgtgctcg ggccagtgcg	360
gccccgcgcg gctgctgccc aacgccatcg ggcgcgtgaa gtggtggcgc ccgaacggac	420
ccgacttcg ctgcatcccg gatcgctacc gcgcgcagcg ggtgcagctg ctgtgccccg	480
gcggcgcgcc gccgcgctcg cgcaagggtc gtctggtggc ctgctgcaag tgcaagcgcc	540
tcaccgcgtt ccacaaccag tcggagctca aggacttcg acctgagacc gcgcggccgc	600
agaagggtcg caagccgcgg ccccgcgccc ggggagccaa agccaaccag gcggagctgg	660
agaacgccta ctacg	674

<210> SEQ ID NO 14
 <211> LENGTH: 213
 <212> TYPE: PRT
 <213> ORGANISM: Rattus norvegicus

<400> SEQUENCE: 14

Met	Gln	Leu	Ser	Leu	Ala	Pro	Cys	Leu	Ala	Cys	Leu	Leu	Val	His	Ala
1				5					10					15	
Ala	Phe	Val	Ala	Val	Glu	Ser	Gln	Gly	Trp	Gln	Ala	Phe	Lys	Asn	Asp
		20						25					30		
Ala	Thr	Glu	Ile	Ile	Pro	Gly	Leu	Arg	Glu	Tyr	Pro	Glu	Pro	Pro	Gln
		35					40				45				
Glu	Leu	Glu	Asn	Asn	Gln	Thr	Met	Asn	Arg	Ala	Glu	Asn	Gly	Gly	Arg
	50					55					60				
Pro	Pro	His	His	Pro	Tyr	Asp	Thr	Lys	Asp	Val	Ser	Glu	Tyr	Ser	Cys
	65				70					75				80	
Arg	Glu	Leu	His	Tyr	Thr	Arg	Phe	Val	Thr	Asp	Gly	Pro	Cys	Arg	Ser
			85						90					95	
Ala	Lys	Pro	Val	Thr	Glu	Leu	Val	Cys	Ser	Gly	Gln	Cys	Gly	Pro	Ala
		100						105					110		
Arg	Leu	Leu	Pro	Asn	Ala	Ile	Gly	Arg	Val	Lys	Trp	Trp	Arg	Pro	Asn
	115					120					125				
Gly	Pro	Asp	Phe	Arg	Cys	Ile	Pro	Asp	Arg	Tyr	Arg	Ala	Gln	Arg	Val
	130				135						140				
Gln	Leu	Leu	Cys	Pro	Gly	Gly	Ala	Ala	Pro	Arg	Ser	Arg	Lys	Val	Arg
	145				150				155					160	
Leu	Val	Ala	Ser	Cys	Lys	Cys	Lys	Arg	Leu	Thr	Arg	Phe	His	Asn	Gln
		165						170						175	
Ser	Glu	Leu	Lys	Asp	Phe	Gly	Pro	Glu	Thr	Ala	Arg	Pro	Gln	Lys	Gly
		180						185						190	
Arg	Lys	Pro	Arg	Pro	Arg	Ala	Arg	Gly	Ala	Lys	Ala	Asn	Gln	Ala	Glu
	195					200						205			
Leu	Glu	Asn	Ala	Tyr											
	210														

<210> SEQ ID NO 15
 <211> LENGTH: 532
 <212> TYPE: DNA
 <213> ORGANISM: Bos torus

<400> SEQUENCE: 15

agaatgatgc cacagaaatc atccccgagc tgggcgagta ccccgagcct ctgccagagc	60
tgaacaacaa gaccatgaac cgggcggaga acggaggagg acctccccac caccctttg	120

-continued

```

agaccaaaga cgccctccgag tacagctgcc gggagctgca cttcaccgcg tacgtgaccg 180
atggggccgtg ccgcagcgcc aagccgggtca ccgagctggt gtgctcgggc cagtgcggcc 240
cggcgcgccct gctgcccac gccatcggcc gcggcaagtg gtggcgccca agcgggcccg 300
acttcgctg catcccccag cgtaccgcg cgcagcgggt gcagctgttg tgctctggcg 360
gcgcggcgcc gcgcgcgcgc aaggtgcgcc tggtagcctc gtgcaagtgc aagcgccctca 420
ctcgcttcca caaccagtcc gagctcaagg acttcggggc cgaggccgcg cggccgaaa 480
cgggcccga gctgcggccc gcgcgccggg gcaccaaagc cagccgggcc ga 532

```

<210> SEQ ID NO 16

<211> LENGTH: 176

<212> TYPE: PRT

<213> ORGANISM: Bos torus

<400> SEQUENCE: 16

```

Asn Asp Ala Thr Glu Ile Ile Pro Glu Leu Gly Glu Tyr Pro Glu Pro
1           5           10           15
Leu Pro Glu Leu Asn Asn Lys Thr Met Asn Arg Ala Glu Asn Gly Gly
20          25          30
Arg Pro Pro His His Pro Phe Glu Thr Lys Asp Ala Ser Glu Tyr Ser
35          40          45
Cys Arg Glu Leu His Phe Thr Arg Tyr Val Thr Asp Gly Pro Cys Arg
50          55          60
Ser Ala Lys Pro Val Thr Glu Leu Val Cys Ser Gly Gln Cys Gly Pro
65          70          75          80
Ala Arg Leu Leu Pro Asn Ala Ile Gly Arg Gly Lys Trp Trp Arg Pro
85          90          95
Ser Gly Pro Asp Phe Arg Cys Ile Pro Asp Arg Tyr Arg Ala Gln Arg
100         105         110
Val Gln Leu Leu Cys Pro Gly Gly Ala Ala Pro Arg Ala Arg Lys Val
115         120         125
Arg Leu Val Ala Ser Cys Lys Cys Lys Arg Leu Thr Arg Phe His Asn
130         135         140
Gln Ser Glu Leu Lys Asp Phe Gly Pro Glu Ala Ala Arg Pro Gln Thr
145         150         155         160
Gly Arg Lys Leu Arg Pro Arg Ala Arg Gly Thr Lys Ala Ser Arg Ala
165         170         175

```

<210> SEQ ID NO 17

<211> LENGTH: 35828

<212> TYPE: DNA

<213> ORGANISM: Mus musculus

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (1)...(35828)

<223> OTHER INFORMATION: n = A,T,C or G

<400> SEQUENCE: 17

```

cgcgttttgg tgagcagcaa tattgcgctt cgatgagcct tggcgttgag attgatacct 60
ctgctgcaca aaaggcaatc gaccgagctg gaccagcgca ttcgtgacac cgtctccttc 120
gaacttattc gcaatggagt gtcattcatc aaggacngcc tgatcgcaaa tgggtgctatc 180
cacgcagcgg caatgaaaaa ccctcagccg gtgaccaata tctacaacat cagccttggt 240
atcctgcgtg atgagccagc gcagaacaag gtaaccgtca gtgccgataa gttcaaagtt 300
aaacctggtg ttgataccaa cattgaaacg ttgatcgaaa acgcgctgaa aaacgctgct 360

```

-continued

gaatgtgcgg cgctggatgt cacaagcaa atggcagcag acaagaaagc gatggatgaa	420
ctggcttcct atgtccgcac ggccatcatg atggaatgtt tccccggtgg tgttatctgg	480
cagcagtgcc gtcgatatga tgcaattgat aattattatc atttgcgggt cctttccggc	540
gatccgcctt gttacggggc ggcgacctcg cgggttttcg ctatttatga aaattttccg	600
gtttaaggcg ttccggttct tcttcgtcat aacttaatgt ttttatttaa aataccctct	660
gaaaagaaag gaaacgacag gtgctgaaag cgagcttttt ggcctctgtc gtttccttcc	720
tctgtttttg tccgtggaat gaacaatgga agtcaacaaa aagcagagct tatcgatgat	780
aagcgggtcaa acatgagaat tcgcgggccgc ataatacgac tcaactatagg gatcgacgcc	840
tactccccgc gcatgaagcg gaggagctgg actccgcatg cccagagacg cccccaacc	900
cccaaaagtc ctgacctcag cctctaccag ctctggttg ggcttgggcg gggcaaggc	960
taccacgttc tcttaacagg tggctgggct gtctcttggc cgcgcgtcat gtgacagctg	1020
cctagtctcg cagtgaagtc accgtggaat gtctgccttc gttgccatgg caacgggatg	1080
acgttacaat ctgggtgtgg agcttttcct gtccgtgtca ggaaatccaa ataccctaaa	1140
ataccctaga agaggaagta gctgagccaa ggctttcctg gcttctccag ataaagtgtg	1200
acttagatgg aaaaaacaa atgataaag acccgagcca tctgaaaatt cctcctaatt	1260
gcaccactag gaaatgtgta tattattgag ctcgatgtg ttcttatttt aaaaagaaaa	1320
ctttagtcat gttattaata agaatttctc agcagtggga gagaaccaat attaacacca	1380
agataaaagt tggcatgatc cacattgcag gaagatccac gttgggtttt catgaatgtg	1440
aagaccccat ttattaagtc cctaagctct gtttttgcac actaggaagc gatggccggg	1500
atggctgagg ggctgtaagg atctttcaat gtcttacctg tgtgtttcct gtctgcacc	1560
taggacctgc tgcctagcct gcagcagagc cagagggggt tcacatgatt agtctcagac	1620
acttgggggc aggttgcatg tactgcacg cttatttcca tacggagcac ctactatgtg	1680
tcaaacacca tatgggtgtc actcttcaga acggtggtgg tcatcatggt gcatttgctg	1740
acggttgatg tgggtgtaga gagctgagat atatggacgc actcttcagc attctgtcaa	1800
cgtggctgtg cattcttgct cctgagcaag tggctaaca gactcacagg gtcagcctcc	1860
agctcagtcg ctgcatagtc ttagggaacc tctcccagtc ctccctacct caactatcca	1920
agaagccagg gggcttgggc gtctcaggag cctgcttgct gggggacagg ttgttgagtt	1980
ttatctgcag taggttgctt aggcatagtg tcaggactga tggctgcctt ggagaacaca	2040
tcctttgccc tctatgcaaa tctgaccttg acatgggggc gctgctcagc tgggagatc	2100
aactgcatac ctaaaagccaa gcctaaagct tcttcgtcca cctgaaactc ctggaccaag	2160
gggtctccgg cacatcctct caggccagtg agggagtctg tgtgagctgc actttccaat	2220
ctcagggcgt gagaggcaga gggaggtggg ggcagagcct tgcagctctt tcctccatc	2280
tggacagcgc tctggctcag cagcccatat gagcacaggc acatcccccac cccaccccca	2340
cctttcctgt cctgcagaat ttaggtctctg ttacgggggg gggggggggg ggggcagtc	2400
tatcctctct taggtagaca ggactctgca ggagacactg ctttgtaaga tactgcagtt	2460
taaatattga tgtgtgagg ggaagcgaa gggcctctt gaccattcag tcaaggtacc	2520
ttctaactcc catcgtattg gggggctact ctagtctag acattgcaga gagcctcaga	2580
actgtagtta ccagtgtggt aggattgatc cttcaggggag cctgacatgt gacagttcca	2640
ttcttaccac agtcaccgaa catttattca gtacctacc cgtaacaggc accgtagcag	2700

-continued

gtactgaggg acggaccact caaagaactg acagaccgaa gccttggaat ataaacacca	2760
aagcatcagg ctctgccaac agaacactct ttaacactca ggccctttaa cactcaggac	2820
ccccacccc accccaagca gttggcactg ctatccacat ttacagaga ggaaaaacta	2880
ggcacaggac gatataagtg gcttgcttaa gcttgctgac atggtaaatg gcagggctgg	2940
attgagaccc agacattcca actctagggt ctatttttct tttttctcgt tgttcgaatc	3000
tgggtcttac tgggtaaact caggctagcc tcacactcat atccttctcc catggcttac	3060
gagtgtagg attccagggt tgtgctacca tgtctgactc cctgtagctt gtctatacca	3120
tcctcacaac ataggaattg tgatagcagc acacacaccg gaaggagctg gggaaatccc	3180
acagagggct ccgcaggatg acaggcgaat gcctacacag aagggtggga agggaagcag	3240
agggaacagc atgggcgtgg gaccacaagt ctatttgggg aagctgccgg taaccgtata	3300
tggctggggt gaggggagag gtcattgagat gaggcaggaa gagccacagc aggcagcggg	3360
tacgggtccc ttattgcca gaggtctgga tcttctctct cttctctctt ccggggctgc	3420
ctgttcattt tccaccactg cctcccatcc aggtctgtgg ctcaggacat caccagctg	3480
cagaaactgg gcatcaccca cgtctgaat gctgccgagg gcaggtcctt catgcacgtc	3540
aacaccagtg ctagcttcta cgaggattct ggcatcacct acttgggcat caaggccaat	3600
gatacgagg agttcaacct cagtgtctac ttgaaaggg ccacagattt cattgaccag	3660
gcgctggccc ataaaaatgg taaggaaagt acattccggc acccatggag cgtaagccct	3720
ctgggacctg cttctccaa agaggccccc acttgaaaaa ggttcagaa agatcccaaa	3780
atatgccacc aactaggat taagtgtcct acatgtgagc cgatgggggc cactgcata	3840
agtctgtgcc atagacatga caatggataa taatatttca gacagagagc aggagttagg	3900
tagctgtgct cctttccctt taattgagtg tgcccatttt tttattcatg tatgtgtata	3960
catgtgtgtg cacacatgcc ataggttgat actgaacacc gtcttcaatc gttcccccac	4020
ccaccttatt ttttaggca ggtctcttc cctgatcctg gggctcattg gtttatctag	4080
gctgctggcc agtgagctct ggagttctgc tttctctac ctcctagcc ctgggactgc	4140
aggggcattg gctgggcccag gcttttatgt cgcgttgggg atctgaactt aggtccctag	4200
gcctgagcac cgtaaaactg ctgccacatc ccagcctgt ttgagcaagt gaaccattcc	4260
ccagaattcc ccagtgagg ctttctctacc cttttatttg ctaggcattc atgagtggtc	4320
acctcgccag aggaatgagt ggccacgact ggctcagggt cagcagccta gagatactgg	4380
gttaagtctt cctgcgcctc gctccctgca gccgcagaca gaaagtagga ctgaatgaga	4440
gctggctagt ggtcagacag gacagaaggc tgagaggggc acagggcaga tgtcagcaga	4500
gcagacagg tctccctctg tgggggaggg gtggccact gcaggtgtaa ttggccttct	4560
ttgtgctcca tagaggcttc ctgggtacac agcagcttcc ctgtcctggt gattcccaaa	4620
gagaactccc taccactgga cttacagaag ttctattgac tgggtgaacg gttcaacagc	4680
tttggtctct ggtggacggt gcatactgt gtatcagctc aagagctcat tcacgaatga	4740
acacacacac acacacacac acacacacac acacaagcta attttgatat gccttaacta	4800
gctcagtgac tgggcatttc tgaacatccc tgaagttagc acacatttcc ctctggtgtt	4860
cctggcttaa cacttctaa atctatattt tatctttgct gccctgttac cttctgagaa	4920
gcccctaggg cacttccctc tcgcacctac attgctggat ggtttctctc ctgcagctct	4980
taaatctgat ccctctgct ctgagccatg ggaacagccc aataactgag ttagacataa	5040
aaacgtctct agccaaaact tcagctaaat ttagacaata aatcttactg gttgtggaat	5100

-continued

ccttaagatt cttcatgacc tccttcacat ggcacgagta tgaagcttta ttacaattgt	5160
ttattgatca aactaactca taaaaagcca gttgtctttc acctgctcaa ggaaggaaca	5220
aaattcatcc ttaactgacg tgtgcacctt gcacaatcca tacgaatata ttaagagtac	5280
taagattttg gttgtgagag tcacatgtta cagaatgtac agctttgaca aggtgcatcc	5340
ttgggatgcc gaagtacact gctgttccag cccctacat tctgaggctg ttttggaaagc	5400
aatgctctgg aagcaacttt agggagtagg atgctggaac agcgggtcac ttcagcatcc	5460
cgatgacgaa tcccgtaaaa gctgtacatt ctgtaacaga ctgggaaagc tgcagacttt	5520
aaggccaggc ccctatggtc cctcttaac cctgtcacac ccaacccgag cccttctcct	5580
ccagccgttc tgtgcttctc actctggata gatggagaac acggccttgc tagttaaagg	5640
agtgaggctt cacccttctc acatggcagt ggttggtcat cctcattcag ggaactctgg	5700
ggcattctgc ctttacttcc tctttttgga ctacagggaa tatatgtga cttgttttga	5760
ccttgtgtat ggggagactg gatctttggt ctggaatgtt tcctgctagt ttttcccat	5820
cctttggcaa accctatcta tatcttacca ctaggcatag tggccctcgt tctggagcct	5880
gccttcaggc tggttctcgg ggaccatgtc cctggtttct cccagcata tgggtttcac	5940
agtgttcact gcgggtggtt gctgaacaaa gcggggattg catcccagag ctccggtgcc	6000
ttgtgggtac actgctaaga taaaatggat actggcctct ctctgaccac ttgcagagct	6060
ctgggtgcct gtgggtacac tgctaagata aaatggatac tggcctctct ctatccactt	6120
gcaggactct agggaacagg aatccattac tgagaaaacc aggggctagg agcagggagg	6180
tagctgggca gctgaagtgc ttggcgacta accaatgaat accagagttt ggaatctctag	6240
aataactcta aaatctgggt gggcagagtg gcctgcctgt aatcccagaa ctccgggaggc	6300
ggagacaggc aatcatcaga gcaactggc taaccagaat agcaaaacac tgagctctgg	6360
gtctgtgag agatcctgcc ttaacatata agagagagaa taaaacattg aagaagacag	6420
tagatgccaa ttttaagccc ccacatgcac atggacaagt gtgctgttga acacacatat	6480
gcactcatgt gaaccaggca tgcacactcg ggcttatcac acacataatt tgaagagag	6540
agtgaagag gagagtgcac attagagttc acaggaaagt gtgagtgagc acacccatgc	6600
acacagacat gtgtgccagg gagtaggaaa ggagcctggg tttgtgtata agagggagcc	6660
atcatgtgtt tctaaggagg gcgtgtgaag gaggcgttgt gtgggctggg actggagcat	6720
ggttgaact gagcatgctc cctgtgggaa acaggagggg gccaccctg cagagggtcc	6780
cactgtccag cgggatcagt aaaagcccct gctgagaact ttaggtaata gccagagaga	6840
gaaaggtagg aaagtggggg gactcccatc tctgatgtag gaggatctgg gcaagtagag	6900
gtgctgttga ggtagaaga ggggtgcaga ggagatgctc ttaattctgg gtcagcagtt	6960
tctttcaaaa taatgcctgt gaggaggtgt aggtggtggc cattactca ctacagagag	7020
ggatgatgat gcccggtgga tgctggaat gccgagcat caaccctggc tctggaagaa	7080
ctccatcttt cagaaggaga gtggtatctgt gtatggccag cggggtcaca ggtgcttggg	7140
gcccctgggg gactcctagc actgggtgat gtttatcgag tgctcttggtg tgcaggcac	7200
tggcctgggg ctttgtttct gtctctgttt tgtttcgttt tttgagacag actcttgcta	7260
tgtatccgtg tcaatcttgg aatctcactg catagcccag gctgcggaga gaggggaggg	7320
caataggcct tgtaagcaag ccacacttca gagactagac tccaccctgc gaatgatgac	7380
aggtcagagc tgagtccggg aagatttttt ttccagctgc caggtggagt gtggagtggc	7440

-continued

agctagcggc	aagggtagag	ggcgagctcc	ctgtgcagga	gaaatgcaag	caagagatgg	7500
caagccagtg	agttaagcat	tctgtgtggg	gagcaggtgg	atgaagagag	aggctgggct	7560
ttcgccctcg	gggggggggt	gaggggtggg	gatgaggtga	gaggagggca	gctccctgca	7620
gtgtgatgag	atttttcctg	acagtgacct	ttggcctctc	cctccccac	ttcccttctt	7680
tcctttcttc	ccaccattgc	tttcttggtc	cttgagaaat	tctgagtttc	cacttcactg	7740
gtgatgcaga	cggaaacaga	agccgtgtgt	gtgtgtgtgt	gtgtgtgtgt	gtgtgtgtgt	7800
gtgtgtgtgt	ttgtgtgtat	gtgtgtgtgt	gtgtttgtgt	gtatgtgtgt	cagtgggaat	7860
ggctcatagt	ctgcaggaa	gtgggcagga	aggaataagc	tgtaggctga	ggcagtgtgg	7920
gatgcaggga	gagagagag	gagggatacc	agagaaggaa	attaagggag	ctacaagagg	7980
gcattgttgg	ggtgtgtgtg	tgtgtgtgtt	gtttatat	gtattggaaa	tacattcttt	8040
taaaaaat	ttatccattt	attttttttt	atgtgcacgt	gtgtgtgcct	gcatgagttc	8100
atgtgtgcca	cgtgtgtg	ggaacccttg	gaggccacaa	gggggcatct	gatccctgg	8160
aactggagtt	ggaggaggtt	gtgagtcctc	tgacatgttt	gctgggaact	gaaccccggt	8220
cctatgcaag	agcaggaa	gcagttatct	gctgagccat	ctctccagtc	ctgaaatcca	8280
ttctcttaaa	atacagtg	cagagacatg	atgggattta	cgtatggatt	taatgtggcg	8340
gtcattaagt	tccggcacag	gcaagcacct	gtaaagccat	caccacaacc	gcaacagtga	8400
atgtgacct	cacccccatg	ttcttcatgt	cccctgtccc	ctccatcctc	cattctcaag	8460
cacctcttgc	tctgcctctg	tcgctggaga	acagtgtgca	tctgcacact	cttatgtcag	8520
tgaagtacaa	cagcctgcac	cccttcctgg	tctgagtatt	tgggttctga	ctctgctatc	8580
acacactact	gtactgcatt	ctctcgctct	ctttttttaa	acataatttt	atttgtttgt	8640
gtgtatgcac	atgtgccaca	tgtgtacaga	tactatggag	gccagaagag	gccatggccg	8700
tccttgagag	tggagttaca	ggcagcgtgt	gagctgcctg	gtgtgggtgc	tgggaaccaa	8760
acttgaatct	aaagcaagca	cttttaactg	ctgaggcagc	tctcagtacc	cttcttcatt	8820
tctccgcctg	ggttccattg	tatggacaca	tgtagctaga	atatcttget	tatctaatta	8880
tgtacattgt	tttgtgtcaa	gagagagtaa	tgctctatag	cctgagctgg	cctcaacctt	8940
gccatcctcc	tgcctcagcc	tcctcctcct	gagtgttagg	atgacaggcg	agtgttaact	9000
tacatgtgtt	catgttttgt	tcaagactga	aggataacat	tcatcacagag	aaggctctggg	9060
tcacaaagt	tgcagttcac	tgaatggcac	aacccgtgat	caagaaacaa	aactcagggg	9120
ctggagagat	ggcactgact	gctcttccag	aggtcgggag	ttcaattccc	agcaaccaca	9180
tgggtggctca	cagccatcta	taacgagatc	tgacgccttc	ttctggtgtg	tctgaagaca	9240
gctacagtgt	actcacataa	aataaataaa	tctttaaaac	acacacacac	acacaattac	9300
cacccacaga	agccactccc	atgttccctc	ccacgtctct	gcctacagta	ctcccaggtt	9360
accactgttc	aggcttctaa	caacctgggt	tacttggggc	tcttttctgc	tctgtggagc	9420
cacacatttg	tgtgcctcat	acacgttctt	tctagtaagt	tgcataattac	tctgcgtttt	9480
tacatgtatt	tattttattgt	agttgtgtgt	gcgtgtgggc	ccatgcatgg	cacagtgtgt	9540
ggggatgtca	gagtattgtg	aacaggggac	agttcttttc	ttcaatcatg	tgggttccag	9600
aggttgaact	cagggtcatca	tgtgtggcag	caaatgcctt	tacccactga	gacatctcca	9660
tattcttttt	ttttccctcg	aggtaggggc	ttgttccata	gcccacactg	gctttgcact	9720
tgcagttcaa	agtactctcc	tgtctccacc	tcttagagta	ttggaattac	gatgtgtact	9780
accacacctg	actggatcat	taattctttg	atgggggcgg	ggaagcgcac	atgctgcagg	9840

-continued

tgaagggatg	actggactgg	acatgagcgt	ggaagccaga	gaacagcttc	agtctaagtc	9900
tctcccaact	gagctatttc	ggtttgccag	agaacaactt	acagaaagt	ctcagtgcca	9960
tgtggattcg	gggttgaggt	tcaactcatc	agcttgacat	tggctcctct	acccactgag	10020
ccttctcact	actctctacc	tagatcatta	attctttttt	aaaaagactt	attagggggc	10080
tggagagatg	gctcagccgt	taagagcacc	gaatgccctt	ccagaggtcc	tgagttcaat	10140
tcccagcatg	ccattgctgg	gcagtagggg	gcgcaggtgt	tcaacgtgag	tagctgttgc	10200
cagttttccg	cggtaggaaa	cctcttgaca	cctgctgtc	cctggtcatt	ctgggtgggt	10260
gcatggtgat	atgcttgttg	tatggaagac	tttgactgtt	acagtgaagt	tgggcttcca	10320
cagttaccac	gtctcccctg	tttcttgacg	gccgggtgct	tgtccattgc	cgcgagggct	10380
acagccgcgc	cccaacgcta	gttatcgctt	acctcatgat	gcggcagaag	atggacgtca	10440
agtctgctct	gagtactgtg	aggcagaatc	gtgagatcgg	ccccaacgat	ggcttctctg	10500
cccaactctg	ccagctcaat	gacagactag	ccaaggaggg	caagtgaaa	ctctaggggtg	10560
cccacagcct	cttttgacga	ggctgactg	ggagggccct	ggcagccatg	tttaggaaac	10620
acagtatacc	cactcccctg	accaccagac	acgtgcccac	atctgtccca	ctctggctct	10680
cggggggccc	tccaccctta	gggagcacat	gaagaagctc	cctaagaagt	tctgctcctt	10740
agccatcctt	tctgttaatt	tatgtctctc	cctgaggtga	ggttcagggt	tatgtccctg	10800
tctgtggcat	agatacatct	cagtgaacca	gggtgggagg	gctatcaggg	tgcatggccc	10860
gggacacggg	cactcttcat	gacccctccc	ccacctgggt	tcttctctgt	tgggccagaa	10920
ccacgagcct	ggtaaaggaa	ctatgcaaac	acaggccctg	acctcccat	gtctgttctt	10980
ggctctcaca	gcccagacag	ccctgctgag	gcagacgaat	gacattaagt	tctgaagcag	11040
agtggagata	gattagtgc	tagatttcca	aaaagaagga	aaaaaaaggc	tgcattttaa	11100
aattatttcc	ttagaattaa	agatactaca	tagggggccct	tgggtaagca	aatccatttt	11160
tcccagaggc	tatcttgatt	ctttggaatg	tttaaagtgt	gccttgccag	agagcttacg	11220
atctatatct	gctgcttcag	agccttccct	gaggatggct	ctgttctctt	gcttggttaga	11280
agagcgatgc	cttgggcagg	gtttccccct	tttcagaata	caggggtgaa	agtccagcct	11340
attacaacaa	aacaaacaaa	caaacaacaa	aaggacctcc	atttggagaa	ttgcaaggat	11400
tttatcctga	attatagtgt	tggtaggttc	aagtcacac	gccaaagtgt	tgccatcctg	11460
gttgctatcc	taagaataat	taggaggagg	aacctagcca	attgcagctc	atgtccgtgg	11520
gtgtgtgcac	gggtgcatat	gttggaaggg	gtgcctgtcc	ccttggggac	agaaggaaaa	11580
tgaaggggcc	ctctgctcac	cctggccatt	tacgggaggg	tctgctgggt	ccacgggtgtc	11640
tgtgcaggat	cctgaaaactg	actcgctgga	cagaaacgag	acttggcggc	accatgagaa	11700
tggagagaga	gagagcaaa	aaagaaacag	cctttaaagg	aactttctaa	gggtgggtttt	11760
tgaacctcgc	tggaccttgt	atgtgtgcac	atttgccaga	gattgaacat	aatcctcttg	11820
ggacttcacg	ttctcattat	ttgtatgtct	ccggggtcac	gcagagccgt	cagccaccac	11880
cccagcacc	ggcacatag	cgtctcataa	aagccattt	tatgagaacc	agagctgttt	11940
gagtaccccg	tgtatagaga	gagttgttgt	cgtggggcac	ccggatccca	gcagcctgggt	12000
tgctgcctg	taggatgtct	tacaggagtt	tgacagagaa	ccttccttgg	agggaaagaa	12060
atatcagggg	tttttgttga	atatttcaaa	ttcagcttta	agtgtaagac	tcagcagtgt	12120
tcatgggttaa	ggtaagggaac	atgccttttc	cagagctgct	gcaagaggca	ggagaagcag	12180

-continued

acctgtctta	ggatgtcact	cccagggtaa	agacctctga	tcacagcagg	agcagagctg	12240
tgcagcctgg	atggtcattg	tccctatttc	tgtgtgacca	cagcaaccct	ggcacatag	12300
ggctggtcac	cctttttttt	tttttttttt	tttttttttg	gcccagaatg	aagtgaccat	12360
agccaagttg	tgtacctcag	tctttagttt	ccaagcggct	ctcttgctca	atacaatgtg	12420
catttcaaaa	taacactgta	gagttgacag	aactgggtca	tgtgttatga	gagaggaaaa	12480
gagaggaaaa	aacaaaacaa	aacaaaacac	cacaaaccaa	aaacatctgg	gctagccagg	12540
catgattgca	atgtctacag	gcccagttca	tgagaggcag	agacagggaag	accgccgaaa	12600
ggccaaggat	agcatgggtc	acgtatcgag	actccagcca	gggtctacgt	cccaagatcc	12660
taggtttttg	attttgggct	ttggtttttg	agacagggtt	tctctgtgta	gccttggtcg	12720
tcctggaaac	cgctctgtag	accaggctgg	cctcaaactt	agagatctgc	ctgactctgc	12780
ctttgagggc	tgggacgaat	gccaccactg	cccaactaag	attccattaa	aaaaaaaaaa	12840
agttcaagat	aattaagagt	tgccagctcg	ttaaagctaa	gtagaagcag	tctcaggcct	12900
gctgcttgag	gctgttcttg	gcttgacact	gaaatctgcc	cccaacagtg	tccaagtcca	12960
catgactttg	agccatctcc	agagaaggaa	gtgaaaattg	tggctcccca	gtcgattggg	13020
acacagtctc	tctttgtcta	ggtaacacat	ggtgacacat	agcattgaac	tctccactct	13080
gagggtgggt	ttccctcccc	ctgcctcttc	tgggttggtc	accccatagg	acagccacag	13140
gacagtcact	agcacctact	ggaaacctct	ttgtgggaac	atgaagaag	agcctttggg	13200
agattcctgg	ctttccatta	gggctgaaag	tacaacgggt	cttggttggtc	tttgctctgt	13260
gtttataaaa	ctagctacta	ttcttcaggt	aaaataccga	tgttggtgaa	aagccaaccc	13320
cgtggctgcc	cgtgagtagg	gggtgggggt	gggaatcctg	gatagtgttc	tatccatgga	13380
aagtgggtgga	ataggaatta	agggtgttcc	ccccccccc	aacctcttcc	tcagacccag	13440
ccactttcta	tgacttataa	acatccaggt	aaaaattaca	aacataaaaa	tggtttctct	13500
tctcaatctt	ctaaagtctg	cctgcctttt	ccaggggtag	gtctgtttct	ttgctgttct	13560
attgtcttga	gagcacagac	taacacttac	caaatgaggg	aactcttggc	ccatactaag	13620
gctcttctgg	gtccagcacc	tcttaagtta	ttttaagaat	tctcacttgg	cctttagcac	13680
acccgccacc	cccaagtggt	tgtggataat	gccatggcca	gcagggggca	ctgttgaggc	13740
gggtgccttt	ccaccttaag	ttgcttatag	tatttaagat	gctaaatgtt	ttaatcaaga	13800
gaagcactga	tcttataata	cgaggataag	agattttctc	acaggaaaatt	gtctttttca	13860
taattctttt	acaggctttg	tcctgatcgt	agcatagaga	gaatagctgg	atatttaact	13920
tgtattccat	tttctctctc	cagcgttagg	ttaactccgt	aaaaagtgat	tcagtggacc	13980
gaagaggctc	agagggcagg	ggatgggtgg	gtgaggcaga	gcactgtcac	ctgccaggca	14040
tgggaggtcc	tgccatccgg	gaggaaaagg	aaagttagc	ctctagtcta	ccaccagtgt	14100
taacgcactc	taagttgtta	acaaaataa	atgtcttaca	ttacaaagac	gtctgttttg	14160
tgtttccttt	tgtgtgtttg	ggctttttat	gtgtgcttta	taactgctgt	gggtgtgctg	14220
ttgttagttt	tgaggtagga	tctcagctg	gccttgaact	tctgatcgcc	tgccctgcc	14280
cctgccccctg	cccctgtccc	tgctccaag	tgttaggaact	aaaagcacat	gccaccacac	14340
cagtacagca	tttttctaac	atttaaaaat	aatcacctag	gggctggaga	gagggttcca	14400
gctaagagtg	cacactgtct	ttgggttagga	cctgagttta	gttcccagaa	cctatactgg	14460
gtggctccag	gtccagagga	tccaggacct	ctggcctcca	tgggcatctg	ctcttagcac	14520
ataccacacat	acagatacac	acataaaaaat	aaaatgaagc	ctttaaaaaac	ctcctaaaaac	14580

-continued

ctagcccttg gaggtacgac tctggaaagc tggcatactg tgtaagtcca tctcatggtg	14640
ttctggctaa cgtaagactt acagagacag aaaagaactc aggggtgtgct ggggggttggg	14700
atggagggaag agggatgagt aggggggagca cggggaactt gggcagtga aattccttgc	14760
aggacactag agggaggataa ataccagtca ttgcacccac tactggacaa ctccaggga	14820
ttatgtctgg tgaaaagaga agggcccaggt tattggctgc attggctgca ttgctgtaac	14880
atTTTTTTaa attgaaaaga aaaagatgta aatcaagggt agatgagtgg ttgctgtgag	14940
ctgagagctg gggtagtgta gacatgtgga caactccatc aaaaagcgac agaagaacg	15000
ggctgtgggt acagctacct ctaatctcca cctccgggag gtgatcaagg ttagccctca	15060
gctagcctgt ggtgcatgag accctgtttc aaaaacttta ataaagaaat aatgaaaaa	15120
gacatcaggg cagatccttg gggccaaagg cggacaggcg agtctcgtg taaggctcgtg	15180
tagaagcggg tgcagtggca cgtgcccgag gcatcatgag agagccctag gtaagtaagg	15240
atggatgtga gtgtgtcggc gtcggcgcac tgcacgtcct ggctgtgggt ctggactggc	15300
atctttgttg agctgtggag gggaaatggg tagggagatc ataaatccc tccgaattat	15360
ttcaagaact gtctattaca attatctcaa aatattaaaa aaaaagaaga attaaaaaac	15420
aaaaaaccta tccaggtgtg gtggtgtgca cctatagcca cgggcacttg gaaagctgga	15480
gcaagaggat ggcgagtttg aaggtatctg gggctgtaca gcaagaccgt cgtcccaaa	15540
ccaaaccaa cagcaaaccc attatgtcac acaagagtgt ttatagttag cggcctcgt	15600
gagagcatgg ggtgggggtg ggggtggggg acagaaatat ctaaactgca gtcaataggg	15660
atccactgag accctggggc ttgactgcag cttaaccttg ggaaatgata aggggtttgt	15720
gttgagttaa agcatcgatt actgacttaa cctcaaatga agaaaaagaa aaaaagaaa	15780
caacaaaagc caaaccaagg ggctggtgag atggctcagt gggtaagagc acccgactgc	15840
tcttccgaag gtccagagtt caaatcccag caaccacatg gtggctcaca accatctgta	15900
acgagatatg atgccctctt ctggtgtgtc tgaagacagc tacagtgtac ttacatataa	15960
taataaatac ttaaaaaaaa aaaaaaaaaa aaagcctaaa ccgagcaaac caggcccca	16020
aacagaaggc aggcacgagc gcaggcacca cgagccatcc tgtgaaaagg cagggtacc	16080
catgggcca ggagggtcca gagagatagg ctggtaagct cagtttctct gtataaccctt	16140
ttcttgtgtg aactacttcc aattacagat aaaataacaa ataaacaaa tctagagcct	16200
ggccactctc tgctcgtctg atttttcttg ttacgtccag cagggtggcg aagtgttcca	16260
aggacagatc gcatcattaa ggtggccagc ataactctcc atcagcaggt ggtgctgtga	16320
gaaccattat ggtgctcaca gaatcccggg cccaggagct gccctctccc aagtctggag	16380
caataggaaa gctttctggc ccagacaggg ttaacagtcc acattccaga gcaggggaaa	16440
aggagactgg aggtcacaga caaaagggcc agcttctaac aacttcacag ctctggtagg	16500
agagatagat ccccccaac aatggccaca gctgggtttg tctgcccga aggaaactga	16560
cttaggaagc aggtatcaga gtccccttcc tgaggggact tctgtctgcc ttgtaaagct	16620
gtcagagcag ctgcattgat gtgtgggtga cagaagatga aaaggaggac ccaggcagat	16680
cgccacagat ggaccggcca cttacaagtc gaggcagggt gcagagcctt gcagaagctc	16740
tgagggtgga cgacactgat tcattaccca gttagcatac cacagcgggc taggcggacc	16800
acagcctcct tcccagtcct cctccagggc tggggagtc tccaaccttc tgtctcagtg	16860
cagcttccgc cagccctccc tcttttgc cctcaggtgt gaacctccc tcctctcctt	16920

-continued

ctccctgtgg	catggccctc	ctgctactgc	aggctgagca	ttggatttct	ttgtgcttag	16980
atagacctga	gatggctttc	tgatttatat	atatatatcc	atcccttgga	tcttacatct	17040
aggaccaga	gctgtttgtg	ataccataag	aggctgggga	gatgatatgg	taagagtgcct	17100
tgctgtacaa	gcataagac	atgagttcga	atccccagca	accatgtgga	aaaataacct	17160
tctaacctca	gagttgaggg	gaaaggcag	tggtattctg	gggcttactg	gccagctagc	17220
cagcctaacc	taaatgtctc	agtcagagat	cctgtctcag	ggaataactt	gggagaatga	17280
ctgagaaaga	cacctcctca	ggctctccat	gcacccacac	agacacacgg	ggggggggta	17340
atgtaataag	ctaagaaata	atgagggaaa	tgattttttg	ctaagaaatg	aaattctgtg	17400
ttggccgcaa	gaagcctggc	cagggaagga	actgcctttg	gcacaccagc	ctataagtca	17460
ccatgagttc	cctggctaag	aatcacatgt	aatggagccc	aggtccctct	tgcttgggtg	17520
ttgcctctcc	cactggtttt	gaagagaaat	tcaagagaga	tctccttggg	cagaattgta	17580
ggtgctgagc	aatgtggagc	tggggccaat	gggattcctt	taaaggcatc	cttcccaggg	17640
ctgggtcata	cttcaatagt	agggtgcttg	cacagcaagc	gtgagaccct	aggttagagt	17700
ccccagaatc	tgcccccaac	cccccaaaaa	ggcatccttc	tgctctggg	tgggtggggg	17760
gagcaaacac	ctttaactaa	gaccattagc	tggcaggggg	aacaaatgac	cttggtctaga	17820
ggaatttggg	caagctggat	tccgccttct	gtagaagccc	cacttgtttc	ctttgttaag	17880
ctggcccaca	gtttgttttg	agaatgcctg	agggggccag	ggagccagac	aattaaagc	17940
caagctcatt	ttgatatctg	aaaaccacag	cctgactgcc	ctgcccgtgg	gaggtactgg	18000
gagagctggc	tggtctcctg	cctcaccaac	gccccccccc	ccaacacaca	ctcctcggtt	18060
cacctgggag	gtgccagcag	caatttgga	gtttactgag	cttgagaagt	cttggggagg	18120
ctgacgtcaa	gcacacccct	tctccacccc	ccccccccc	accccctga	ggaggagggt	18180
gaggaaacat	gggaccagcc	ctgctccagc	ccgtccttat	tggctggcat	gaggcagagg	18240
gggctttaaa	aaggcaaccg	tatctaggct	ggacactgga	gcctgtgcta	ccgagtgcct	18300
tcctccacct	ggcagcatgc	agccctcact	agccccgtgc	ctcatctgcc	tacttggtga	18360
cgtgcctct	tggtgctggg	agggccaggg	gtggcaagcc	ttcaggaatg	atgccacaga	18420
ggtcatccca	gggcttgagg	agtaccccca	gcctcctcct	gagaacaacc	agacctgaa	18480
ccgggcggag	aatggaggga	gacctcccca	ccatccctat	gacgccaaag	gtacgggatg	18540
aagaagcaca	ttagtggggg	gggggggtcct	gggaggtgac	tgggttggtt	ttagcatctt	18600
cttcagaggt	ttgtgtgggt	ggctagcctc	tgctacatca	gggcagggac	acatttgcct	18660
ggaagaatac	tagcacagca	ttagaacctg	gagggcagca	ttggggggct	ggtagagagc	18720
acccaaggca	gggtggaggc	tgaggtcagc	cgaagctggc	attaacacgg	gcatgggctt	18780
gtatgatggg	ccagagaatc	tcctcctaag	gatgaggaca	caggtcagat	ctagctgctg	18840
accagtgggg	aagtgatatg	gtgaggctgg	atgccagatg	ccatccatgg	ctgtactata	18900
tcccacatga	ccaccacatg	aggtaaaaga	ggccccagct	tgaagatgga	gaaaccgaga	18960
ggctcctgag	ataaagtcac	ctgggagtaa	gaagagctga	gactgggaagc	tggtttgac	19020
cagatgcaag	gcaaccctag	attgggtttg	ggtgggaacc	tgaagccagg	aggaatccct	19080
ttagttcccc	cttgcccagg	gtctgtctaa	tgagcccaga	gggttagcat	taaaagaaca	19140
gggtttgtag	gtggcatgtg	acatgagggg	cagctgagtg	aaatgtcccc	tgtatgagca	19200
cagggtggc	cacttgccct	gagcttgac	cctgacccca	gctttgcctc	attcctgagg	19260
acagcagaaa	ctgtggaggc	agagccagca	cagagagatg	cctgggggtg	gggtgggggt	19320

-continued

atcacgcacg gaactagcag caatgaatgg ggtgggggtgg cagctggagg gacactccag 19380
 agaaatgacc ttgctgggtca ccatttgtgt gggaggagag ctcattttcc agcttgccac 19440
 cacatgctgt ccctcctgtc tcctagccag taagggatgt ggaggaaagg gccaccccaa 19500
 aggagcatgc aatgcagtca cgtttttgca gaggaagtgc ttgacctaaq ggcactattc 19560
 ttggaaagcc ccaaaactag tccttccttg gcaaacagg cctccccac ataccacctc 19620
 tgcaggggtg agtaaattaa gccagccaca gaagggtggc aaggcctaca cctccccct 19680
 gttgtgcccc ccccccccc gtgaagggtc atcctggcct ctgcccctct ggctttggtg 19740
 ctgggatttt ttttttcctt ttatgtcata ttgatcctga caccatggaa cttttggagg 19800
 tagacaggac ccacacatgg attagttaa agcctcccat ccatctaagc tcatggtagg 19860
 agatagagca tgtccaagag agggaggcag gcatcagacc tagaagatat ggctgggcat 19920
 ccaacccaat ctcttcccc ggagaacaga ctctaagtc gatccagcca ccttgagta 19980
 accagctcaa ggtacacaga acaagagagt ctggtataca gcagggtgcta aacaaatgct 20040
 tgtggtagca aaagctatag gttttgggtc agaactccga cccaagtcgc gagtgaagag 20100
 cgaaaggccc tctactgcc accgccccgc cccacactgg ggtcctataa cagatcactt 20160
 tcacccttgc gggagccaga gagccctggc atcctaggta gcccccccg ccccccccc 20220
 gcaagcagcc cagccctgcc tttggggcaa gttcttttct cagcctggac ctgtgataat 20280
 gaggggggtg gacgcgccgc ctttggctgc tttcaagtct aatgaattct tatccctacc 20340
 acctgcccc ctaccccgct cctccacagc agctgtcctg atttattacc ttcaattaa 20400
 ctccactcct ttctccatct cctgggatac cgcccctgct ccagtggctg gtaaaggagc 20460
 ttaggaagga ccagagccag gtgtggctag aggtctaccg gcagggtctg ggtgaggag 20520
 ctaaaactga agagtgtttg gttagtaggc acaaacctt ggggtgggac ctagtagccg 20580
 gagaagtga gatgggcgct gagaagtca agaccatcca tccttaacta cacagccagt 20640
 ttgaggccag cctgggtac ataaaaacc aatctcaaaa gctgccaatt ctgattctgt 20700
 gccacgtagt gcccgatgta atagtggatg aagtcgttga atcctggggc aaacctat 20760
 acagatgtgg ggaanaagca ctttaagtac cctgcccaca gatcacaag aaagtaagt 20820
 acagagctcc agtgtttcat ccctgggttc caaggacagg gagagagaag ccagggtggg 20880
 atctcactgc tccccggtgc ctcttcccta taatccatac agattcgaaa gcgcagggca 20940
 ggtttgaaa aaagagagaag ggtggaagga gcagaccagt ctggcctagg ctgcagcccc 21000
 tcacgcatec ctctctccgc agatgtgtcc gagtacagct gccgcgagct gcactacacc 21060
 cgcttctcta cagacggccc atgccgagc gccaaagccg tcaccgagtt ggtgtgctcc 21120
 ggccagtgcg gccccgcgcg gctgctgccc aacgccatcg ggcgcgtgaa gtggtggcgc 21180
 ccgaacggac cggatttccg ctgcatcccg gatcgctacc gcgcgcagcg ggtgcagctg 21240
 ctgtgccccg ggggcgcgcg gccgcgctcg cgcaaggtgc gtctggtggc ctctgcaaag 21300
 tgcaagcgcc tcaccgctt ccacaaccag tcggagctca aggacttcgg gccggagacc 21360
 gcgcggcgcc agaagggtcg caagccgcgg cccggcgccc ggggagccaa agccaaccag 21420
 gcggagctgg agaagccta ctagagcgag cccgcgccta tgcaacccc gcgcgacccg 21480
 attcgttttc agtgtaaagc ctgcagccca ggccaggggt gccaaacttt ccagaccgtg 21540
 tggagttccc agcccagtag agaccgagg tccttctgcc cgctcgggg gatggggagg 21600
 ggggtggggt cccgcgggcc aggagaggaa gcttgagtcc cagactctgc ctgaccccg 21660

-continued

gtgggatggg ggtctttcta ccctcgccgg acctatacag gacaaggcag tgtttccacc	21720
ttaaaggga gggagtgtgg aacgaaagac ctgggactgg ttatggacgt acagtaagat	21780
ctactccttc caccacaatg taaagcctgc gtgggctaga tagggtttct gaccctgacc	21840
tggccactga gtgtgatgtt gggctacgtg gttctctttt ggtacggtct tctttgtaa	21900
atagggaccg gaactctgct gagattccaa ggattggggt accccgtgta gactggtgag	21960
agagaggaga acagggggagg ggttagggga gagattgtgg tgggcaaccg cctagaagaa	22020
gctgtttgtt ggctcccagc ctccgcccct cagagggttg gcttccccc ctccttcctc	22080
tcaaatctgc cttcaaatcc atatctggga tagggaaggc cagggccga gagatggtgg	22140
aagggccaga aatcacactc ctggccccc gaagagcagt gtccgcccc caactgcctt	22200
gtcatattgt aaagggattt tctacacaac agtttaaggt cgttgaggga aactgggctt	22260
gccagtccac tccatcctt gtcccttgcc aggacaccac ctccctgcctg ccaccacgg	22320
acacatttct gtctagaaac agagcgtcgt cgtgctgtcc tctgagacag catatcttac	22380
attaaaaaga ataatacggg gggggggggc ggagggcgca agtgttatac atatgctgag	22440
aagctgtcag gcgccacagc accaccaca atctttttgt aaatcatttc cagacacctc	22500
ttactttctg tgtagatttt aattgttaaa aggggaggag agagagcgtt tgtaacagaa	22560
gcacatggag gggggggtag ggggggtggg gctggtagt ttggcgaact ttccatgtga	22620
gactcatcca caaagactga aagccgcgtt ttttttttta agagttcagt gacatattta	22680
ttttctcatt taagttattt atgccaacat ttttttcttg tagagaaagg cagtgttaat	22740
atcgctttgt gaagcacaag tgtgtgtgtt tttttgtttt ttgttttttc cccgaccaga	22800
ggcattgtta ataaagacaa tgaatctcga gcaggaggct gtggtcctgt tttgtcaacc	22860
acacacaatg tctcgccact gtcattctac tcccttcctt tggtcacaag acccaaacct	22920
tgacaacacc tccgactgct ctctggtagc ccttggtgga atacgtgttt cctttgaaa	22980
gtcacattca tcccttcctt tgcaaacctg gctctcattc cccagctggg tcatcgtcat	23040
accctcacc cagcctccct ttagctgacc actctccaca ctgtcttcca aaagtgcacg	23100
tttcaccgag ccagttccct ggtccaggtc atcccattgc tccctcctgc tccagaccct	23160
tctcccacaa agatgttcat ctcccactcc atcaagcccc agtggccctg cggctatccc	23220
tgtctcttca gttagctgaa tctactgtct gacaccacat gaattccttc cctgtctta	23280
aggttcatgg aactcttgcc tgcccctgaa ccttccagga ctgtcccagc gtctgatgtg	23340
tcctctctct tgtaaagccc caccacca tttgattccc aattctagat ctcccttgt	23400
tcattccttc acgggatagt gtctcatctg gccaaagcct gcttgatatt gggataaatg	23460
caaagccaag tacaattgag gaccagttca tcattgggccc aagctttttc aaatgtgaa	23520
ttttacacct atagaagtgt aaaagccttc caaagcagag gcaatgcctg gctcttcctt	23580
caacatcagg gctcctgctt tatgggtctg gtgggtagt acattcataa acccaacact	23640
aggggtgtga aagcaagatg attgggagtt cgaggccaat cttggctatg aggcctgtc	23700
tcaacctctc ctccctccct ccagggtttt gttttgtttt gtttttttga tttgaaactg	23760
caacacttta aatccagtca agtgcatctt tgcgtgaggg gaactctatc cctaataata	23820
gcttccatct tgatttgtgt atgtgcacac tgggggttga acctgggcct ttgtacctgc	23880
cgggcaagct ctctactgct ctaaaaccag ccctcactgg ctttctgttt caactcccaa	23940
tgaattcccc taaatgaatt atcaatatca tgtctttgaa aaataccatt gagtgctgct	24000
ggtgtccctg tggttccaga ttccaggaag gacttttcag ggaatccagg catcctgaag	24060

-continued

aatgtcttag	agcaggagggc	catggagacc	ttggccagcc	ccacaaggca	gtgtggtgca	24120
gaggggtgag	atggaggcag	gcttgcaatt	gaagctgaga	cagggctactc	aggattaaaa	24180
agcttcccc	aaaacaattc	caagatcagt	tcctgggtact	tgacactgtt	cagctatgca	24240
gagcccagt	ggcataggt	aagacaccgg	ttgtactgtc	atgtactaac	tgtgcttcag	24300
agccggcaga	gacaaataat	gttatggtga	ccccagggga	cagtgattcc	agaaggaaaca	24360
cagaagagag	tgtgtctaga	ggctgcctga	aggagaaggg	gtcccagact	ctctaagcaa	24420
agactccact	cacataaaga	cacaggctga	gcagagctgg	ccgtggatgc	agggagccca	24480
tccaccatcc	tttagcatgc	ccttgtattc	ccatcacatg	ccagggatga	ggggcatcag	24540
agagtccaag	tgatgcccc	acccaaacac	acctaggact	tgctttcttg	gacagacaga	24600
tgaggagag	actaggttgg	gctgtgatcc	cattaccaca	aagagggaaa	aaacaaaaaa	24660
caaaacaaca	aaacaaaaaa	aacaaaacaa	aaacaaaaaa	aacccaaggt	ccaatttgta	24720
ggtcaggtta	gagtttattt	atggaagtt	atattctacc	tccatgggg	ctacaaggct	24780
ggcgcccatc	agaaagaaca	aacaacaggc	tgatctggga	ggggtggtac	tctatggcag	24840
ggagcacagt	tgcttgggg	acagccagac	acggggcttg	tattaatcac	agggcttgta	24900
ttaataggct	gagagtcaag	cagacagaga	gacagaagga	aacacacaca	cacacacaca	24960
cacacacaca	cacacacaca	catgcacaca	ccactcactt	ctcactcgaa	gagcccctac	25020
ttacattcta	agaacaaacc	attcctcttc	ataaaggaga	caaagtgtca	gaaacccaaa	25080
agagccacag	ggtccccact	ctctttgaaa	tgacttggac	ttgttgacag	gaagacagag	25140
gggtctgcag	aggcttcctg	ggtgacccag	agccacagac	actgaaatct	ggtgctgaga	25200
cctgtataaa	ccctcttcca	caggttccct	gaaaggagcc	cacattcccc	aacctgtct	25260
cctgaccact	gaggatgaga	gcacttgggc	cttccccatt	cttgagatgc	accttggttt	25320
ccccatctga	gggcacatga	ggtctcaggt	cttgggaaag	ttccacaagt	attgaaagt	25380
ttcttgtttt	gtttgtgatt	taatttaggt	gtatgagtgc	ttttgcttga	atatatgcct	25440
gtgtagcatt	tacaagcctg	gtgcctgagg	agatcagaag	atggcatcag	ataccctgga	25500
actggacttg	cagacagtta	tgagccactg	tgtgggtgct	aggaacagaa	cctggatcct	25560
ccggaagagc	agacagccag	cgctcttagc	cactaagcca	tactgaggt	tctttctgtg	25620
gctaaagaga	caggagacaa	aggagagttt	cttttagtca	ataggacat	gaatgttcct	25680
cgtaacgtga	gactagggca	gggtgatccc	ccagtgcac	cgatggccct	gtgtagttat	25740
tagcagctct	agtccttattc	cttaataagt	cccagtttgg	ggcaggagat	atgtattccc	25800
tgctttgaag	tggctgaggt	ccagttatct	acttccaagt	acttgtttct	ctttctggag	25860
ttggggaagc	tcctgcctg	cctgtaaatg	tgctccattct	tcaaccttag	acaagatcac	25920
tttccctgag	cagtcaggcc	agtcacaaagc	ccttcaattt	agctttcata	aggaacaccc	25980
cttttgttgg	gtggaggtag	cacttgccct	gaatcccagc	attaagaagg	cagagacagt	26040
cggatctctg	tgagttcaca	gccagcctgg	tctacggagt	gagttccaag	acagccaggc	26100
ctacacagag	aaacctgtgc	tcgaaaaaaa	caaaaaacaa	agaaataaag	aaaaagaaaa	26160
caaaaaacgaa	caaacagaaa	aacaagccag	agtgtttgc	cccgtatttt	attaatcata	26220
tttttgtccc	tttgccattt	tagactaaaa	gactcgggaa	agcagggtctc	tctctgtttc	26280
tcacccggac	acaccagaa	ccagatgtat	ggaagatggc	taatgtgctg	cagttgcaca	26340
tctggggctg	ggtggattgg	ttagatggca	tgggctgggt	gtggttacga	tgactgcagg	26400

-continued

agcaaggagt atgtggtgca tagcaaacga ggaagtttgc acagaacaac actgtgtgta	26460
ctgatgtgca ggtatgggca catgcaagca gaagccaagg gacagcctta gggtagtggt	26520
tccacagacc cctccccctt tttaacatgg gcatctctca ttggcctgga gcttgccaac	26580
tgggctgggc tggctagctt gtaggtccca gggatctgca tatctctgcc tccctagtgc	26640
tgggattaca gtcatatatg agcacacctg gcttttttat gtgggttctg gctttgaac	26700
ccagatctga gtgcttgcaa ggcaatcggg tgaatgactg cttcatctcc ccagaccctg	26760
ggattctact ttctattaaa gtatttctat taaatcaatg agcccctgcc cctgcaactca	26820
gcagttctta ggcctgctga gagtcaagtg gggagtgaga gcaagcctcg agaccccatc	26880
agcgaagcag aggacaaga aatgaaaact tgggattcga ggctcgggat atggagatac	26940
agaaagggtc agggaaggaa atgaaccaga tgaatagagg caggaaagggt agggccctgc	27000
atacatggaa cctggtgtac atgttatctg catgggggtt gcattgcaat ggctcttcag	27060
caggttcacc aactctggaa acagaagcca aaaagaagag taggtggtgt tggagtcaga	27120
tactgtcagt catgcctgaa gaaatggaag caattaacga tgcgccgcaa ttaggatatt	27180
agctccctga agaaaggcaa gaagctgggc tgtgggcact gaaggagact ttgaatgatg	27240
tcacattctc tgtatgccta gcagggcagt attggagact gagacttgac ttgtgtgtcc	27300
atatgattcc tccttttctc acagtcactt ggggctcctg agcttcgtcc ttgtccaaga	27360
acctggagct ggcagtgggc agctgcagtg atagatgtct gcaagaaaga tctgaaaga	27420
gggaggaaga tgaaggaccc agaggaccac cgacctctgc tgcctgacaa agctgcagga	27480
ccagctcttc ctacagatgg gagacagagg cgagagatga atggtcaggg gagagtcag	27540
agaaaggaga gggtagggca gagaccaaag gagggaaaca cttgtgtctc acagctactg	27600
actgagtacc agctgcgtgg cagacagcca atgccaaggc tcggctgac atggcacctc	27660
gtgggaactc tagccagtg ctggcagagg ggaagtctga atgggtcatg gtttggtat	27720
gatctgaatg tgggccagcc ctagtctcct tccagttgct gggataaagc accctgacca	27780
aaactacttt tttgtttgtt tgttttggtt tggttttgtt tggttttctg aggcagggtt	27840
ttctgtatc accctagctg tcctggaact cactctgtag accaggctgg cctcgaactc	27900
agaaatcccc ctgcctctgc ctccctaagt ctggaattaa aggcctgcgc caccactgcc	27960
ggcccaaagc tactttaaga gagagagagg aatgtataag tattataatt ccaggttata	28020
gttcattgct gtagaattgg agtcttcata ttccaggtaa tctccacag acatgccaca	28080
aaacaacctg ttctacgaaa tctctcatgg actccctcc ccagtaattc taaactgtgt	28140
caaatctaca agaaatagtg acagtcacag tctctaactg tttgggcatg agtctgaagt	28200
ctcattgcta agtactggga agatgaaaac tttacctagt gtcagcattt ggagcagagc	28260
ctttgggatt tgagatgggc ttttgagag ctccctaatg ctacatggag agagggggcc	28320
tgggagagac ccatacacct tttgctgcct tatgtcacct gacctgctcc ttgggaagct	28380
ctagcaagaa ggccttcctt ggatcaccca ccaccttgca cctccagaac tcagagccaa	28440
attaaacttt cttgttactg tcgtcaaac acagtcgggc tgggttgtat cactgtcaat	28500
gggaaacaga cttgcctgga tggataactt gtacattgca taatgtctag aaatgaaaag	28560
tcctatagag aaaaagaaaa ttactgtggc cacagataga ggccttgag gaggctggct	28620
ttgtctctcc cgaggaggtg gcgagtaagg tgtaaatgtt catggatgta aatgggcccc	28680
tatatgaggg tctggggtaa caagaaggcc tgtgaatata aagcactgaa ggtatgtcta	28740
gtctggagaa ggtcactaca gagagttctc caactcagt cccatacaca cacacacaca	28800

-continued

```

cacacacaca cacacacaca cacacacaca ccacaaagaa aaaaaggaag aaaaatctga 28860
gagcaagtac agtacttaaa attgtgtgat tgtgtgtgtg actctgatgt cacatgctca 28920
tcttgcccta tgagtgaag accaaatggc cctgagagg cataacaacc acactgttgg 28980
ctgtgtgctc acgtttttct taaagcgtct gtctggtttg ctgctagcat caggcagact 29040
tgacgacagac tacatatgct cagccctgaa gtccttctag ggtgcatgct tcttcagaat 29100
ttcagaaagt catctgtggc tccaggaccg cctgcactct ccctctgccg cgaggctgca 29160
gactctaggc tgggggtgaa gcaacgctta cctctgggac aagtataaca tgttggtttt 29220
tctttccctc tgtggctcca acctggacat aaaatagatg caagctgtgt aataaatatt 29280
tcctcccgct cacttagtgc tcaacaataa ctactctgag agcacttatt aatagggtggc 29340
ttagacataa gctttggctc attccccac tagctcttac ttctttaact ctttcaaacc 29400
attctgtgtc tccacatgg ttagttacct ctccttccat cctggttcgc ttcttcttc 29460
gagtcgccct cagtgtctct aggtgatgct tgtaagatat tctttctaca aagctgagag 29520
tggtggcact ctgggagttc aaagccagcc tgatctacac agcaagctcc aggatatcca 29580
gggcaatgtt gggaaaaacct ttctcaaca aaaagagggg ttcagttgtc aggaggagac 29640
ccatgggcta agaagctag acgagccatg gtgatgcata cctttcatcc aagcacttag 29700
gaggcaaaga aaggtgaac tctttgactt tgaggccagc taggttacat agtgataccc 29760
tgcttagtgt gtgtgtgtgt gtgtgtgtgt gtgtgtgtgt gtgtgtaatt taaaagtcta 29820
aaaatgcatt cttttaaaaa tatgtataag tatttgcctg cacatatgta tgtatgtatg 29880
tataccatgt gtgtgtctgg tgctgaagga ctaggcatag actccctaga actagagtca 29940
tagacagttg tgacactccc caacccccca ccatgtgggt gcttgaagct aaactcctgt 30000
cctttgtaaa gcagcaggtg tctatgaacc ctgaaccatc tctccagtct ccagatgtgc 30060
attctcaag aggagtcctt catatttccc taaactgaac atccttatca gtgagcatcc 30120
tcgagtcacc aaagctactg caaacctctt tagggaacat tcactattca cttctacttg 30180
gctcatgaaa cttaagtaca cacacacaaa cacacacaca cacacagagt catgcaactca 30240
caaaagcatg catgtacacc attcttatta gactatgctt tgctaaaaga ctttcctaga 30300
tactttaaaa catcacttct gccttttggg gggcaggttc caagattggg actggcgtac 30360
tggaactga acaaggtaga gatctagaaa tcacagcagg tcagaagggc cagcctgtac 30420
aagagagagt tccacacctt ccaggaacac tgagcagggg gctgggacct tgcctctcag 30480
cccaagaaac tagtgcgttt cctgtatgca tgcctctcag agattccata agatctgcct 30540
tctgccataa gatctcctgc atccagacaa gcctagggga agttgagagg ctgcctgagt 30600
ctctcccaca ggcccttctt tgcctggcag tattttttta tctggaggag aggaatcagg 30660
gtgggaatga tcaaatcaaa ttatcaagga aaaagtaaaa aacatatata tatatatatt 30720
aactgatcta gggagctggc tcagcagtta agagttctgg ctgcccttgc ttcagatctt 30780
gctttgatcc ccagcaccac catgatggct ttcaactgta tctctgcttc caggggatcc 30840
aacagcctct tctgacctcc atagacaaga cctagtcctc tgcaagagca ccaaatgctc 30900
ttatctgttg atccatctct ctagcctcat gccagatcat ttaaaactac tggacactgt 30960
cccattttac gaagatgtca ctgccagtc atttgccatg agtggatatt tcgattcttt 31020
ctatgttctc acccttgcaa tttataagaa agatatctgc atttgtctcc tgagagaaca 31080
aagggtggag ggctactgag atggctctag gggtaaaagt gcttgccaca aaatctgaca 31140

```

-continued

acttaagttt ggtcttggaa tccacatggt ggagagagag aagagattcc cgtaagttgt	31200
cctcaaaactt cccacacatg tgctgtggct tatgtgtaac cccaataagt aaagatagtt	31260
ttaaactacta cataaggtag ggtttcttca tgaccccaag gaatgatgcc cctgatagag	31320
cttatgctga aaccccatct ccattgtgcc atctggaaaag agacaattgc atcccggaaa	31380
cagaatcttc atgaatggat taatgagcta ttaagaaaagt ggcttgggta ttgcacatgc	31440
tgcgggcgta atgacctcca ccatgatgtt atccagcatg aaggctctca ccagaagtca	31500
tacaaatctt cttaggcttc cagagtcgtg agcaaaaaaa gcacacctct aaataaatta	31560
actagcctca ggtagttaac caccgaaaat gaaccaaggc agttctaata caaaaccact	31620
tccttccctt gttcaaaacca cagtgcctta ttatctaaaa gataaacttc aagccaagct	31680
tttaggttg cagtatttat gtaacaacaa ggcccgttga cacacatctg taactcctag	31740
tactgggcct caggggcaga gacaggtgga gccctggagt ttgaattcca ggttctgtga	31800
gaaactctgt ctgaaaagac aatatggtga gtgacccggg aggatatctg atattgactt	31860
ctggccaaca cacagccatc tctgcacatc tgtagttgca agccttttgc actaagtttg	31920
gccagagta cagtttgcaa gtgtttgttg actgaatgca cgtgttgctg gtgactctaca	31980
aagtcacctt ccttctcaag ctgacgacac tggcttcggc cagctgctca ttcaagcctc	32040
tttgagagt catcacgggg atggggggagc agggccctc cctagaacac caagcctgtg	32100
gttgtttatt caggacatta ttgaggcca agatgacaga taactctatc acttggccaa	32160
cagtcgggtg ttgcggtgtt aggttatttc tgtgtctgca gaaaacagtg caacctggac	32220
aaaagaaata aatgatata tttttcattc aggcactag attccgtggt acaaaaggct	32280
ccctggggaa cgaggccggg acagcgccgc tcctgagtcg ctatttccgt ctgtcaactt	32340
ctctaacttc ttgatttctt ccctctgtct gtttccctcc tcttgctggg gccagtgga	32400
gtctgtgtac tcacagggag gaggttgga aagccctggt cctctacggg ctgggggaaag	32460
gggggaagct gtcggcccaag tgacttttcc ccttttctct ttttcttaga aaccagtctc	32520
aatttaagat aatgagtctc ctcatccacg tgtgctcact attcataggg acttatccac	32580
ccccgccctg tcaatctgac taagtaagac aagtcanaatt taaaagggaa cgtttttcta	32640
aaaatgtggc tggaccgtgt gccggcacga aaccagggat ggcggtctaa gttacatgct	32700
ctctgccagc cccggtgcct tttcctttcg gaaaggagac ccggaggtaa aacgaagtgt	32760
ccaacttttg atgatggtgt gcgcggggtg actctttaaa atgtcatcca tacctgggat	32820
aggggaaggct cttcagggag tcatctagcc ctcccttcag gaaaagattc cacttccggt	32880
ttagttagct tccacctggt cccttatccg ctgtctctgc ccactagtcc tcatccatcc	32940
ggtttccgcc ctcatccacc ttgccctttt agttcctaga aagcagcacc gtagtcttgg	33000
caggtgggcc attggtcact ccgctaccac tggtaccatg gccaccaagg tgtcatttaa	33060
atatgagctc actgagtcct gcgggatggc ttggttggtg atatgcttgc tgcaaaatcg	33120
tgagaactgg agttcaatcc ccagcacatg gatgtatttc cagcacctgg aaggcaggga	33180
gcagagatct taaagctcct ggccagacag cccagcctaa ttagtaatca gtgagagacc	33240
ctgtctcaag aaacaagatg gaacatcaaa ggtcaacctc ttgtctccac acacacaaat	33300
acacacatgc acatacatcc acacacaggc aaacacatgc acacacctga acacctcca	33360
caaatacata cataaaaaaa taaatacata cacacatata tacatacacc aacattccct	33420
ctccttagtc tcctggctac gctcttgcac cccccactaa ggcttcaact tcttctatct	33480
cttcatcttg actcctctgt actttgcatg ccttttccag caaaggcttt tctttaaatc	33540

-continued

tccgtcatto ataaactccc tctaaatttc tccccctgcc cttttctttc tctctaggga	33600
gataaagaca cactactacaa agtcaccgtg ggaccagttt attcaccacac ccacccttgc	33660
ttctgttcat ccggccagct aagtagtcca acctctctgg tgctgtaccc tggaccctgg	33720
cttcaccaca gctcctccat gctaccacgc cctgcaaacc ttcagcctag cctctggttc	33780
tccaaccagc acaggcccgag tctggcttct atgtcctaga aatctccttc attctctcca	33840
tttccctcct gaattctacca ccttctttct ccttctcct gacctctaat gtcttggtca	33900
aacgattaca aggaagccaa tgaatttagc agtttggggg acctcagagt cagcagggga	33960
gctgggatga attcacattt ccaggccttt gctttgctcc ccggattctg acaggcagtt	34020
ccgaagctga gtccaggaag ctgaatttaa aatcacactc cagctgggtt ctgaggcagc	34080
cctaccacat cagctggccc tgactgagct gtgtctgggt gccagtgggt ctggtgggtc	34140
tggtgggtgt ggtgggtggg gtgggtgggt tggtgggtgt ggtgggtgtg tgtgtgtgtg	34200
ttttctgctt ttacaaaact tttctaattc ttatacaaa gacaaatctg cctcatatag	34260
gcagaaagat gacttatgcc tatataagat ataaagatga ctttatgcc cttattagca	34320
atagttactg tcaaaagtaa ttctatttat acacccttat acatgggtatt gcttttgttg	34380
gagactctaa aatccagatt atgtatttaa aaaaaattc cccagtcctt aaaaggtgaa	34440
gaatggaccc agatagaagg tcacggcaca agtatggagt cggagtgtgg agtcctgcc	34500
atggctctga cagaagcctc cagagagggt ccaagacaaa tgcctcgctt cctaaggaa	34560
actggcagcc ctgatgaggt accagagatt gctaagtga ggaatacagg atcagaccca	34620
tgagggggct taaagcgtga ctgtagcagc cctccgctga ggggctccag gtgggcgccc	34680
aagggtctgc agtgggagcc acatgagagg tgatgtcttg gagtcacctc gggtaaccatt	34740
gtttagggag gtggggattt gtggtgtgga gacaggcagc ctcaaggatg cttttcaaca	34800
atgggttgat agttggaact aaaacagggg ccatcacact ggctcccata gctctgggct	34860
tgccagcttc cacatctgcc cccacacccc tgtctggcac cagctcaagc tctgtgatcc	34920
tacacatcca aaagaggaag agtagcctac tgggcctgcc acctctctg gaacctcagg	34980
tgagagtgtg gcaagcccta ggctcctgtc caggatgcag ggctgccaga taggatgtc	35040
agctatctcc tgagctggaa ctatttttagg aataaggatt atgcccgcgc ggggttggcc	35100
agcacccccag cagcctgtgc ttgcgtaaaa gcaagtgtctg ttgatttatc taaaaacaga	35160
gccgtggacc caccacaggg acaagtatgt atgcatctgt tcatgtatc tgaanaagcga	35220
cacaaccatt ttccacatca tggcatcttc ctacccccca ttcttttttg ttttgtttt	35280
ttgagacagg gtttctctgt gtagtcctgg ctgtcctgga actcactttg tagaccaggc	35340
tggcctcgaa ctcaaaaatc ctgggattaa aggtgtgtgc caccacgccc gccctaacc	35400
cccattctta atggtgatcc agtggttgaa atttcgggcc acacacatgt ccattaggga	35460
ttagctgtct tcttctgagc tacctggtag aatctttatc ccctggggcc tgggtcctg	35520
atccctgact cgggcccgat caagtccagt tccctggccc gatcaagtcc agttcctggg	35580
cccgaacaa tccagtcctt agctcgatta gctcatcctg gctccctggc ctgttcttac	35640
ttacactctt ccccttgctc tggacttggt gctttcttta ctcaagttgt ctgccacagt	35700
ccctaagcca cctctgtaag acaactaaga taatacttcc ctcaagcacg gaaagtcctg	35760
agtcaccaca cctctggag gtgtgtggac acatgttcat gcgtgtggtt gcgcttacgt	35820
acgtgtgc	35828

-continued

<210> SEQ ID NO 18
 <211> LENGTH: 9301
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapien

<400> SEQUENCE: 18

tagaggagaa gtctttgggg agggtttgct ctgagcacac ccctttccct ccctccgggg	60
ctgagggaaa catgggacca gccctgcccc agcctgtcct cattggctgg catgaagcag	120
agaggggctt taaaaaggcg accgtgtctc ggctggagac cagagcctgt gctactggaa	180
gggtggcgtgc cctcctctgg ctggtaccat gcagctccca ctggccctgt gtctcgtctg	240
cctgctggta cacacagcct tccgtgtagt ggagggccag ggggtggcagg cgttcaagaa	300
tgatgccacg gaaatcatcc ccgagctcgg agagtacccc gagcctccac cggagctgga	360
gaacaacaa accatgaacc gggcgagaa cggaggcgcg cctccccacc accccttga	420
gaccaaagg atgggggtga ggagagaatt cttagtaaaa gatcctgggg aggttttaga	480
aacttctctt tgggaggctt ggaagactgg ggtagacca gtgaagattg ctggcctctg	540
ccagcactgg tccaggaaaca gtcttgcttg gaggtggggg aagaatggct cgctggtgca	600
gccttcaaat tcagggtcag aggcattgag caacagacgc tggtagagc ccagggcagg	660
gaggacgctg ggggtgtgag ggtatggcat cagggcacga gaacaggctc aggggctcag	720
aaaagaaaag gtttcaaaag atctcctcct gggaatatag gagccacgtc cagctgctgg	780
taccactggg aagggaacaa ggtaaggag cctcccatcc acagaacagc acctgtgggg	840
caccggacac tctatgctgg tgggtgctgt cccaccaca cagaccaca tcatggaatc	900
cccaggaggt gaacccccag ctgaaagggg aagaaacagg ttccaggcac tcagtaactt	960
ggtagtgaag agagctgagg tgtgaacctg gtttgatcca actgcaagat agccctgggt	1020
tgtggggggg tgtgggggac agatctccac aaagcagtgg ggaggaaggc cagagaggca	1080
ccctgcagt gtgcattgcc catggcctgc ccagggagct ggcactgaa ggaatggag	1140
tttccggcac agtttttagc cctgacatgg gtgcagctga gtccaggccc tggaggggag	1200
agcagcatcc tctgtgcagg agtagggaca tctgtcctca gcagccacc cagtcccaac	1260
cttgccatcc tccaggggag ggagaaggaa gaggaacctt gggttcctgg tcaggcctgc	1320
acagagaagc ccaggtgaca gtgtgcatct ggctctataa ttggcaggaa tcctgaggcc	1380
atgggggctg ctgaaatgac acttcagact aagagcttcc ctgtcctctg gccattatcc	1440
aggtggcaga gaagtccact gccagggctc ctggacccca gccctccccg cctcacaacc	1500
tgttgggact atggggtgct aaaaagggca actgcatggg aggcagacca ggaacctccg	1560
tcttcaaaat ggaggacaag ggcgcctccc cccacagctc cccttctagg caaggtcagc	1620
tgggctccag cgaactgcctg aagggtgta aggaacccaa acacaaaatg tccaccttgc	1680
tggactccca cgagaggcca cagcccctga ggaagccaca tgctcaaac aaagtcatga	1740
tctgcagagg aagtgcctgg cctaggggcg ctattctcga aaagccgcaa aatgccccct	1800
tccttgggca atgcccccc tgaccacaca cacattccag ccctgcagag gtgaggatgc	1860
aaaccagccc acagaccaga aagcagcccc agacgatggc agtggccaca tctccccctg	1920
tgtgcttctg cttcagatg ggggtggggg gtggccttct ctgtccccct tctggtttgg	1980
tcttaagact atttttcatt ctttcttctg acattggaac tatccccatg aaacctttgg	2040
gggtggactg gtactcacac gacgaccagc tatttaaaaa gctccccacc atctaagtcc	2100
accataggag acatggtcaa ggtgtgtgca ggggatcagg ccaggcctcg gagcccaatc	2160

-continued

tctgcctgcc	cagggagtat	caccatgag	cgcccatcca	gataacacag	aacaagaaat	2220
gtgcccagca	gagagccagg	tcaatgtttg	tggcagctga	acctgtaggt	tttgggtcag	2280
agctcagggc	ccctatggta	ggaagtaac	gacagtaaaa	agcagccctc	agctccatcc	2340
cccagcccag	cctcccatgg	atgctcgaac	gcagagcctc	cactcttgcc	ggagccaaaa	2400
ggtgctggga	ccccaggga	gtggagtccg	gagatgcagc	ccagcctttt	gggcaagttc	2460
ttttctctgg	ctgggacctca	gtattctcat	tgataatgag	ggggttgagc	acactgcctt	2520
tgattccttt	caagtcta	gaattcctgt	cctgatcacc	tccccttcag	tcctctgcct	2580
ccacagcagc	tgccctgatt	tattaccttc	aattaacctc	tactcctttc	tccatccctt	2640
gtccaccctt	cccaagtggc	tggaaaagga	atttgggaga	agccagagcc	aggcagaagg	2700
tgtgctgagt	acttaccctg	cccaggccag	ggaccctgcg	gcacaagtgt	ggcttaaatc	2760
ataagaagac	cccagaagag	aatgataat	aataatacat	aacagccgac	gctttcagct	2820
atatgtgcca	aatggatttt	tctgcattgc	gtgtgtaatg	gattaactcg	caatgcttgg	2880
ggcggcccat	tttgacagca	ggaagaagag	agagggttaa	gaacttgccc	aagatgacac	2940
ctgcagttag	cgatggagcc	ctggtgtttg	aaccccagca	gtcattttggc	tccgagggga	3000
cagggtgcgc	aggagagctt	tccaccagct	ctagagcacc	tgggaccttc	ctgcaataga	3060
tgttcagggg	caaaagcctc	tggagacagg	cttggcaaaa	gcagggtctg	ggtggagaga	3120
gacgggcccg	tccaggggcag	gggtggccag	gcgggcgggc	accctcacgc	gcgcctctct	3180
ccacagacgt	gtccgagtac	agctgcgcgc	agctgcactt	caccgcgtac	gtgaccgatg	3240
ggccgtgcgc	cagcgccaag	ccggtcaccg	agctggtgtg	ctccggccag	tgcggcccgg	3300
cgcgcctgct	gccccaacgc	atcgccgcgc	gcaagtgggtg	gcgacctagt	gggcccgaact	3360
tccgctgcat	ccccgaccgc	taccgcgcgc	agcgcgtgca	gctgctgtgt	cccggtggtg	3420
aggcgcccg	cgcgcgcaag	gtgcgcctgg	tggcctcgtg	caagtgcagg	cgcctcacc	3480
gcttccacaa	ccagtcggag	ctcaaggact	tccgggaccga	ggccgctcgg	ccgcagaagg	3540
gccggaagcc	gcggccccgc	gcccgagagc	ccaaagccaa	ccaggccgag	ctggagaacg	3600
cctactagag	cccgccgcgc	cccctcccca	ccggcggggc	cccggccctt	gaaccgcgc	3660
cccacatttc	tgctctctgc	gcgtggtttg	attgtttata	tttcattgta	aatgcctgca	3720
acccaggggc	gggggctgag	accttccagg	ccctgaggaa	tcccgggcgc	cggaaggcc	3780
cccctcagcc	cgcagctga	gggtgccac	ggggcagggg	aggggaattga	gagtcacaga	3840
cactgagcca	cgcagcccg	cctctggggc	cgcctacctt	tgctgggtccc	acttcagagg	3900
aggcagaagt	ggaagcattt	tcaccgccct	gggtttttaa	gggagcgggtg	tgggagtggg	3960
aaagtccagg	gactgggtta	gaaagtggga	taagattccc	ccttgacact	cgtgcccct	4020
cagaaagcct	gaggcgtgcc	cagagcacia	gactgggggc	aactgtagat	gtggtttcta	4080
gtcctggctc	tgccactaac	ttgctgtgta	accttgaaact	acacaattct	ccttcgggac	4140
ctcaatttcc	actttgtaaa	atgagggtgg	aggtgggaat	aggatctcga	ggagactatt	4200
ggcatatgat	tccaaggact	ccagtgcctt	ttgaatgggc	agagggtgaga	gagagagaga	4260
gaaagagaga	gaatgaatgc	agttgcattg	attcagtgcc	aaggtaactt	ccagaattca	4320
gagttgtgat	gctctcttct	gacagccaaa	gatgaaaaac	aaacagaaaa	aaaaaagtaa	4380
agagtctatt	tatggctgac	atatttacgg	ctgacaaaact	cctggaagaa	gctatgctgc	4440
ttcccagcct	ggcttcccgc	gatgtttggc	tacctccacc	cctccatctc	aaagaaataa	4500

-continued

catcatccat	tggggtagaa	aaggagaggg	tccgaggggtg	gtgggagggga	tagaaatcac	4560
atccgcccc	acttcccaaa	gagcagcacc	cctccccga	cccatagcca	tgttttaaag	4620
tcaccttcg	aagagaagtg	aaagggtcaa	ggacactggc	cttgaggcc	cgaggagca	4680
gccatcaca	actcacagac	cagcacatcc	cttttgagac	accgccttct	gcccaccact	4740
cacggacaca	ttctgccta	gaaaacagct	tcttactgct	cttacatgtg	atggcataatc	4800
ttacactaaa	agaatattat	tgggggaaaa	actacaagtg	ctgtacatat	gctgagaaac	4860
tgacagcat	aatagctgcc	acccaaaaat	ctttttgaaa	atcattttcca	gacaacctct	4920
tactttctgt	gtagttttta	attgttaaaa	aaaaaaagtt	ttaaacagaa	gcacatgaca	4980
tatgaaagcc	tgacgagctg	gtcgtttttt	tggcaattct	tccacgtggg	acttgtccac	5040
aagaatgaaa	gtagtgggtt	ttaaagaggt	aagttacata	tttattttct	cacttaagtt	5100
atttatgcaa	aagtttttct	tgtagagaat	gacaatgtta	atattgcttt	atgaattaac	5160
agtcgttct	tccagagtc	agagacattg	ttaataaaga	caatgaatca	tgaccgaaag	5220
gatgtggct	cattttgtca	accacacatg	acgtcatttc	tgtcaaagtt	gacacccttc	5280
tcttggtcac	tagagctcca	accttgga	caccttgac	tgctctctgg	tgcccttctg	5340
ggcaattatg	tcttctttg	aaaagtcacg	tttatccctt	cctttccaaa	cccagaccgc	5400
atttcttcac	ccagggcacg	gtaataacct	cagccttgta	tccttttagc	agcctccctt	5460
ccatgtggc	ttccaaaatg	ctgttctcat	tgtatcactc	ccctgctcaa	aagccttcca	5520
tagctcccc	tgcccagga	tcaagtgcag	tttccctatc	tgacatggga	ggccttctct	5580
gcttgactcc	cacctccac	tccaccaagc	ttcctactga	ctccaaatgg	tcatgcagat	5640
ccctgcttcc	ttagtttgcc	atccacactt	agcaccacca	ataactaatc	ctctttcttt	5700
aggattcaca	ttacttgtca	tctcttcccc	taaccttcca	gagatgttcc	aatctcccat	5760
gatccctctc	tctctgagg	ttccagcccc	ttttgtctac	accactactt	tggttcttaa	5820
ttctgttttc	catttgacag	tcattcatgg	aggaccagcc	tgccaagtc	ctgcttagta	5880
ctggcataga	caacacaaag	ccaagtacaa	ttcaggacca	gctcacagga	aacttcactc	5940
tcttcgaagt	gtggatttga	tgcctcctgg	gtagaaatgt	aggatcttca	aaagtgggcc	6000
agcctcctgc	acttctctca	aagtcctgcc	tccccaaggt	gtcttaatag	tgctggatgc	6060
tagctgagtt	agcatcttca	gatgaagagt	aaccctaag	ttactcttca	gttgccctaa	6120
ggtgggatgg	tcaactggaa	agctttaaat	taagtccagc	ctaccttggg	ggaaccacc	6180
cccacaaaga	aagctgaggt	ccctcctgat	gactgtcag	tttaactacc	aataaccac	6240
ttgaattaat	catcatcacc	aagtccttga	taggtgtgag	tgggtatcag	tgccgggtcc	6300
cttctgggg	ctccagcccc	cgaggaggcc	tcagttagcc	cctgcagaaa	atccatgcat	6360
catgagtgtc	tcagggccca	gaatatgaga	gcaggtagga	aacagagaca	tcttccatcc	6420
ctgagaggca	gtgcgttcca	gtgggtgggg	acacgggctc	tgggtcaggt	ttgtgttgtt	6480
tgttttgttg	ttttgagaca	gagtcctgct	ctattgcccc	ggctggagtg	cagtgtcaca	6540
atctcggtt	actgcaactt	ctgccttccc	ggattcaagt	gattctcctg	cctcagcctc	6600
cagagttagt	gggattacag	gtgcgtgcca	ccacgcctgg	ctaatttttg	tatttttgat	6660
agagacgggg	tttcaccatg	ttggccaggc	tagtctcgaa	ctcttgacct	caagtgatct	6720
gcctgctctg	gcctcccaaa	gtgctgggat	tacaggcgtg	agccaccaca	cccagcccca	6780
ggttggtgtt	tgaatctgag	gagactgaag	caccaagggg	ttaaattgtt	tgcccacagc	6840
catacttggg	ctcagttcct	tgccctaccc	ctcacttgag	ctgcttagaa	cctggtgggc	6900

-continued

acatgggcaa taaccagggtc acactgtttt gtaccaagtgt ttatgggaat ccaagatagg	6960
agtaatttgc tctgtggagg ggatgagggg tagtggttag ggaaagcttc acaaagtggg	7020
tgttgcttag agattttcca ggtggagaag ggggcttcta ggcagaaggc atagcccaag	7080
caaagactgc aagtgcattg ctgctcatgg gtagaagaga atccaccatt cctcaacatg	7140
taccgagtc ttgccattgt caaggcaaca tgggggtacc aggaattcca agcaatgtcc	7200
aaacctaggg tctgttttct gggacctgaa gatacaggat ggatcagccc aggtcgcaat	7260
cccattacca cgagggggaa aaaaacctga aggctaaatt gtaggctggg ttagaggtta	7320
tttatggaaa gttatatctt acctacatgg ggtctataag cctggcgcca atcagaaaaa	7380
gaacaaacaa cagacctagc tgggaggggc agcattttgt ttaggggggc ggggcacatg	7440
ttctgggggt acagccagac tcagggcttg tattaatagt ctgagagtaa gacagacaga	7500
gggatagaag gaaatagggt cctttctctc tctctctctc tctctctctc actctctctc	7560
tctctcacac acacacacag acacacacac acgctctgta ggggtctact tatgctcaa	7620
gtacaaatca ggccacattt acacaaggag gtaaaggaaa agaactgttg aggagccaca	7680
ggaccccaaa attccctggt ttccttgaat caggcaggac ttacgcagct gggagggtg	7740
agagcctgca gaagccacct gcgagtaagc caagttcaga gtcacagaca ccaaaagctg	7800
gtgccatgct ccacaccgcg ccacctccca cctgctcctt gacacagccc tgtgctccac	7860
aaccgggctc ccagatcatt gattatagct ctggggcctg caccgtcctt cctgccacat	7920
ccccacccca ttcttggaac ctgccctctg tcttctccct tgtccaaggg caggcaaggg	7980
ctcagctatt gggcagcttt gaccaacagc tgaggctcct tttgtggctg gagatgcagg	8040
aggcagggga atattcctct tagtcaatgc gacctgtgc ctggtttgcc cagggtggtc	8100
tcgttttacac ctgtaggcca agcgtaatga ttaacagctc ccacttctac tctaaaaaat	8160
gacccaatct gggcagtaaa ttatatggtg cccatgctat taagagctgc aacttgctg	8220
gcgtggtggc tcacacctgt aatccagta ctttgggacg tcaaggcggg tggatcacct	8280
gaggtcacga gttagagact ggcctggcca gcatggcaaa acccatctt tactaaaaat	8340
acaaaaatta gcaaggcatg gtggcatgca cctgtaatcc caggtactcg ggaggctgag	8400
acaggagaaat ggcttgaaac caggaggcag aggttgagct gagccaagat tgtgccactg	8460
ccctccagcc ctggcaacag agcaagactt catctcaaaa gaaaaggat actgtcaatc	8520
actgcaggaa gaacccagggt aatgaatgag gagaagagag gggctgagtc accatagtg	8580
cagcacccgac tcctgcagga aaggcgagac actgggtcat gggtagtgaa gggcgccctg	8640
aatgacgttc tgccttagag accgaacctg agccctgaaa gtgcatgcct gttcatgggt	8700
gagagactaa attcatcatt ccttggcagg tactgaatcc tttcttacgg ctgccctcca	8760
atgccaat tccctacaat tgtctgggggt gcctaagctt ctgcccacca agagggccag	8820
agctggcagc gagcagctgc aggtaggaga gataggtagc cataaggagag gtgggaaaga	8880
gagatgggaa gagaggggtg cagagcacac acctccctg cctgacaact tcctgagggc	8940
tggtcatgcc agcagattta aggcggaggc aggggagatg gggcgggaga ggaagtga	9000
aaggagaggg tggggatgga gaggaagaga gggtagatcat tcattcatc cattgctact	9060
gactggatgc cagctgtgag ccaggcacca ccctagctct gggcatgtg ttgtaatctt	9120
ggagcctcat ggagctcaca gggagtgctg gcaaggagat ggataatgga cggataacaa	9180
ataaacattt agtacaatgt ccgggaatgg aaagtctctg aaagaaaaat aaagctgggtg	9240

-continued

agcatataga cagccctgaa ggcgccagg ccaggcattt ctgaggaggt ggcatttgag 9300

c 9301

<210> SEQ ID NO 19
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer for PCR

<400> SEQUENCE: 19

ccggagctgg agaacaacaa g 21

<210> SEQ ID NO 20
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer for PCR

<400> SEQUENCE: 20

gcactggccg gagcacacc 19

<210> SEQ ID NO 21
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer for PCR

<400> SEQUENCE: 21

aggccaaccg cgagaagatg acc 23

<210> SEQ ID NO 22
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer for PCR

<400> SEQUENCE: 22

gaagtccagg gcgacgtagc a 21

<210> SEQ ID NO 23
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer for PCR

<400> SEQUENCE: 23

aagcttggtta ccatgcagct cccac 25

<210> SEQ ID NO 24
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer for PCR

<400> SEQUENCE: 24

aagcttctac ttgtcatcgt cgtccttgta gtcgtaggcg ttctccagct 50

<210> SEQ ID NO 25
<211> LENGTH: 19

-continued

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer for PCR

<400> SEQUENCE: 25

gcactggccg gagcacacc 19

<210> SEQ ID NO 26
<211> LENGTH: 39
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer for PCR

<400> SEQUENCE: 26

gtcgtcggat ccatggggtg gcaggcggtc aagaatgat 39

<210> SEQ ID NO 27
<211> LENGTH: 57
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer for PCR

<400> SEQUENCE: 27

gtcgtcaagc ttctacttgt catcgctctt gtagtcgtag gcgttctcca gtcgggc 57

<210> SEQ ID NO 28
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer for PCR

<400> SEQUENCE: 28

gacttgatc ccagggggtg caggcggtc 29

<210> SEQ ID NO 29
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer for PCR

<400> SEQUENCE: 29

agcataagct tctagtaggc gttctccag 29

<210> SEQ ID NO 30
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer for PCR

<400> SEQUENCE: 30

gacttgatc cgaagggaag aagaaagg 29

<210> SEQ ID NO 31
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer for PCR

<400> SEQUENCE: 31

-continued

agcataagct tttaatccaa atcgatgga 29

<210> SEQ ID NO 32
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer for PCR

<400> SEQUENCE: 32

actacgagct cggccccacc acccatcaac aag 33

<210> SEQ ID NO 33
<211> LENGTH: 34
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer for PCR

<400> SEQUENCE: 33

acttagaagc tttcagtcct cagccccctc ttcc 34

<210> SEQ ID NO 34
<211> LENGTH: 66
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer for PCR

<400> SEQUENCE: 34

aatctggatc cataacttcg tatagcatac attatacgaa gttatctgca ggattcgagg 60
gccccct 66

<210> SEQ ID NO 35
<211> LENGTH: 82
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer for PCR

<400> SEQUENCE: 35

aatctgaatt ccaccgggtg taattaaata acttcgtata atgtatgcta tacgaagtta 60
tagatctaga gtcagcttct ga 82

<210> SEQ ID NO 36
<211> LENGTH: 62
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer for PCR

<400> SEQUENCE: 36

atttaggtga cactatagaa ctcgagcagc tgaagcttaa ccacatggtg gctcacaacc 60
at 62

<210> SEQ ID NO 37
<211> LENGTH: 54
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer for PCR

<400> SEQUENCE: 37

aacgacggcc agtgaatccg taatcatggt catgctgcca ggtggaggag ggca 54

-continued

<210> SEQ ID NO 38
 <211> LENGTH: 31
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer for PCR

<400> SEQUENCE: 38

attaccaccg gtgacaccg cttcctgaca g

31

<210> SEQ ID NO 39
 <211> LENGTH: 61
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer for PCR

<400> SEQUENCE: 39

attacttaat taaacatggc gcgccatag gccggccct aattgcggcg catcgtaaat

60

t

61

<210> SEQ ID NO 40
 <211> LENGTH: 34
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer for PCR

<400> SEQUENCE: 40

attacggcgc gccgcaaagg aattcaagat ctga

34

<210> SEQ ID NO 41
 <211> LENGTH: 34
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer for PCR

<400> SEQUENCE: 41

attacggcgc gcccttcaca ggccgcaccc agct

34

We claim:

1. An isolate antibody or antigen binding fragment thereof which specifically binds to a TGF-beta binding protein, wherein said binding protein is selected from the group consisting of:

- (a) a polypeptide encoded by a polynucleotide that comprises a nucleotide sequence selected from SEQ ID NOs: 1, 5, 9, 11, 13, and 15, or a complementary sequence thereto which encodes a TGF-beta binding protein, and
 - (b) a polypeptide that comprises an amino acid sequence selected from SEQ ID NOs: 2, 6, 10, 12, 14, and 16.
2. An isolated antibody or antigen binding fragment thereof which specifically binds to a TGF-beta binding protein, wherein said binding protein is selected from the group consisting of:

- (a) a polypeptide encoded by a polynucleotide that comprises a nucleotide sequence selected from SEQ ID NOs: 1, 5, 7, 9, 11, 13, and 15, or a complementary sequence thereto which encodes a TGF-beta binding protein; and
- (b) a polypeptide that comprises an amino acid sequence selected from SEQ ID NOs: 2, 6, 3, 10, 12, 14, and 16.

45

3. The isolated antibody or binding fragment thereof of claim 1, wherein the isolated antibody or binding fragment thereof is a polyclonal antibody.

50

4. The isolated antibody or binding fragment thereof of claim 1, wherein the isolated antibody or binding fragment thereof is a monoclonal antibody.

55

5. The isolated antibody or binding fragment thereof of claim 1, wherein the isolated antibody or binding fragment thereof is a humanized antibody.

55

6. The isolated antibody or binding fragment thereof of any one of claims 3-5, and 1, wherein the antibody has an affinity of at least 10^{-7} M.

60

7. The isolated antibody or binding fragment thereof of any one of claims 3-5 and 1, wherein the antibody has an affinity of at least 10^{-8} M.

65

8. A hybridoma that produces an antibody according to any one of claims 3-5 and 1.

9. A method of producing monoclonal antibodies, comprising:

- (a) immunizing an animal with a TGF-beta binding protein or portion thereof, wherein said binding protein is selected from the group consisting of:

claims

157

- (i) a polypeptide encoded by a polynucleotide that comprises a nucleotide sequence selected from SEQ ID NOs:1, 5, 7, 9, 11, 13, and 15, or a complementary sequence thereto which encodes a TGF-beta binding protein, and
 - (ii) a polypeptide that comprises an amino acid sequence selected from SEQ ID NOs: 2, 6, 8, 10, 12, 14, and 16;
 - (b) harvesting spleen cells from said animal;
 - (c) fusing said spleen cells with a myeloma cell line; and
 - (d) culturing said fused cells under conditions that allow the production of said antibody.
10. A method for the production of an antibody of any one of claims 3-5 and 1 comprising culturing hybridoma cells under conditions that allow the production of said antibody.
11. A method for the production of an antibody, or binding fragment thereof, of any one of claims 3-5 and 1 comprising:

158

- (a) providing a recombinant host cell capable of producing said antibody; and
 - (b) culturing said cell under conditions that allow the production of said antibody.
12. A method for immunizing an animal comprising injecting an animal with a TGF-beta binding protein or portion thereof, wherein said binding protein is selected from the group consisting of:
- (i) a polypeptide encoded by a polynucleotide that comprises a nucleotide sequence selected from SEQ ID NOs:1, 5, 7, 9, 11, 13, and 15, or a complementary sequence thereto, and
 - (ii) a polypeptide that comprises an amino acid sequence selected from SEQ ID NOs: 2, 6, 8, 10, 12, 14, and 16.

* * * * *